

**Proprietary Information of United Healthcare:** The information contained in this document is proprietary and the sole property of United HealthCare Services, Inc. Unauthorized copying, use and distribution of this information are strictly prohibited. Copyright 2020 United HealthCare Services, Inc.



UnitedHealthcare® Community Plan  
**Medical Benefit Drug Policy**

## CRYSVITA® (BUROSUMAB-TWZA)

Policy Number: CS2020D007I

Effective Date: TBD

Formatted: Font: Bold

Table of Contents	Page
<a href="#">APPLICATION</a>	1
<a href="#">COVERAGE RATIONALE</a>	1
<a href="#">APPLICABLE CODES</a>	2
<a href="#">BACKGROUND</a>	2
<a href="#">CLINICAL EVIDENCE</a>	3
<a href="#">U.S. FOOD AND DRUG ADMINISTRATION</a>	5
<a href="#">CENTERS FOR MEDICARE AND MEDICAID SERVICES</a>	5
<a href="#">REFERENCES</a>	5
<a href="#">POLICY HISTORY/REVISION INFORMATION</a>	5
<a href="#">INSTRUCTIONS FOR USE</a>	<b>ERROR! BOOKMARK NOT DEFINED.</b>

### Commercial Policy

- [Crysvita® \(Burosomab-Twza\)](#)

### APPLICATION

This Medical Benefit Drug Policy only applies to the state of Louisiana

### COVERAGE RATIONALE

**Crysvita (burosomab) is proven and medically necessary for the treatment of X-linked hypophosphatemia (XLH) when the following criteria are met:**

- For **initial therapy**, all of the following:
  - Diagnosis of XLH, confirmed by **one** of the following:
    - Genetic testing (e.g., confirmed PHEX gene mutation in patient or first-degree relative)
    - Elevated Serum fibroblast growth factor 23 (FGF23) level > 30 pg/mL;
  - and**
  - Patient is age 6 months or greater; **and**
  - One** of the following:
    - Patient epiphyseal plate has **not** fused; **or**
    - All** of the following:
      - Patients' epiphyseal plate has fused; **and**
      - Patient is experiencing clinical signs and symptoms of the disease (e.g., limited mobility, musculoskeletal pain, bone fractures); **and**
      - Failure, contraindication, or intolerance to therapy with **vitamin D analog therapy (e.g., calcitriol, paricalcitol, doxercalciferol)** in combination with an oral phosphate agent (e.g., K-Phos®, K-Phos Neutra®);
  - and**
  - Prescribed by, or in consultation with, an endocrinologist or specialist experienced in the treatment of metabolic bone disorders; **and**
  - Fasting serum phosphorus is below the normal range for age; **and**
  - Dosing is in accordance with the United States Food and Drug Administration approved labeling; **and**
  - Initial authorization will be for no more than 12 months
- For **continuation therapy**, all of the following:
  - Patient has previously received treatment with burosomab; **and**

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

- Prescribed by, or in consultation with, an endocrinologist or specialist experienced in the treatment of metabolic bone disorders; **and**
- Patient has experienced normalization of serum phosphate while on therapy; **and**
- Patient has experienced a positive clinical response to burosomab (e.g., enhanced height velocity, improvement in skeletal deformities, reduction of fractures, reduction of generalized bone pain); **and**
- Dosing is in accordance with the United States Food and Drug Administration approved labeling; **and**
- Reauthorization will be for no more than 12 months.

**Crysvita (burosomab) is proven and medically necessary for the treatment of Fibroblast Growth Factor 23 (FGF23)-related hypophosphatemia in tumor-induced osteomalacia (TIO) when the following criteria are met:<sup>1</sup>**

- **For initial therapy, all of the following:**
  - [Diagnosis of FGF23-related hypophosphatemia in TIO associated with phosphaturic mesenchymal tumors; and](#)
  - [Disease cannot be curatively resected or localized; and](#)
  - [Patient is age 2 years or greater; and](#)
  - [Prescribed by, or in consultation with, an oncologist, an endocrinologist, or specialist experienced in the treatment of metabolic bone disorders; and](#)
  - [History of failure, contraindication, or intolerance to therapy with vitamin D analog therapy \(e.g., calcitriol, paricalcitol, doxercalciferol\) in combination with an oral phosphate agent \(e.g., K-Phos®, K-Phos Neutra®\); and](#)
  - [Fasting serum phosphorus is below the normal range for age; and](#)
  - [Dosing is in accordance with the United States Food and Drug Administration approved labeling; and](#)
  - [Initial authorization will be for no more than 12 months.](#)
- **For continuation therapy, all of the following:**
  - [Patient has previously received treatment with burosomab; and](#)
  - [Prescribed by, or in consultation with, an oncologist, an endocrinologist or specialist experienced in the treatment of metabolic bone disorders; and](#)
  - [Patient has experienced an increase of serum phosphate while on therapy; and](#)
  - [Patient has experienced a positive clinical response to burosomab \(e.g., enhanced height velocity, improvement in skeletal deformities, reduction of fractures, reduction of generalized bone pain\); and](#)
  - [Dosing is in accordance with the United States Food and Drug Administration approved labeling; and](#)
  - [Reauthorization will be for no more than 12 months.](#)

Formatted: Font: Bold

## APPLICABLE CODES

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Coverage Determination Guidelines may apply.

HCPCS Code	Description
J0584	Injection, burosomab-twza, 1 mg
ICD-10 Diagnosis Code	Description
E83.31	Familial hypophosphatemia
<a href="#">M83.8</a>	<a href="#">Other adult osteomalacia</a>

Formatted: Font: Bold

## BACKGROUND

XLH is a heritable disorder of renal phosphate transport, which results in abnormal phosphate hemostasis, resulting in hypophosphatemia and abnormal bone mineralization. Elevated serum FGF23 levels are observed in patients with XLH, and believed to be associated with phosphate level abnormalities. Burosomab inhibits excess FGF23 levels, which

Formatted: Font: Bold

results in normalization of serum phosphate.<sup>1-3</sup> Combining active vitamin D metabolites with a balanced dose of phosphate has been the mainstay of therapy for XHL. Most affected children are candidates for treatment. In adults, the role of treatment has not been well studied; treatment is generally reserved for individuals with symptoms such as skeletal pain, upcoming orthopedic surgery, biochemical evidence of osteomalacia with an elevated serum alkaline phosphatase (ALP) level, or recurrent pseudofractures or stress fractures. The primary goals of treatment in children are to correct or minimize rickets/osteomalacia, as assessed by radiographic abnormalities and resolution of skeletal abnormalities. In contrast with children, once a patient reaches adult height and the epiphyses have fused, the goal of therapy is simply to manage generalized bone pain and enhance limited mobility, if either occurs, and to cure any non-union fractures.<sup>6</sup>

**Tumor-induced osteomalacia (TIO) is a rare acquired paraneoplastic syndrome in which the biochemical and bone mineralization abnormalities closely resemble those in genetic forms of hypophosphatemic rickets. Clinical and experimental studies have documented that tumors produce humoral factor(s) that cause the abnormalities that occur in TIO. The tumors, typically benign, often are small, slow-growing polymorphous neoplasms, most commonly, phosphaturic mesenchymal tumors of the mixed connective tissue type.<sup>8,9</sup> The mesenchymal tumors associated with TIO ectopically express and secrete fibroblast growth factor 23 (FGF23) and other phosphaturic proteins. Most affected patients have increased circulating FGF23 levels.**

**Formatted:** Font: Bold

## CLINICAL EVIDENCE

A randomized, active-controlled, open-label, multicenter, phase 3 clinical trial (NCT02915705) compared the efficacy and safety of continuing conventional therapy (oral phosphate and active vitamin D), versus switching to burosumab, in pediatric patients with X-linked hypophosphatemia.<sup>7</sup> Children aged 1 to 12 years were enrolled who had a total Thacher rickets severity score of at least 2.0, fasting serum phosphorus lower than 0.97 mmol/L (3.0 mg/dL), confirmed PHEX mutation or variant of unknown significance in the patient or a family member with appropriate X-linked dominant inheritance, and received conventional therapy for at least 6 consecutive months for children younger than 3 years or at least 12 consecutive months for children older than 3 years. Patients were randomly assigned (1:1) to receive either subcutaneous burosumab, starting at 0.8 mg/kg every 4 weeks or conventional therapy for 64 weeks. The primary endpoint was change in rickets severity at week 40, assessed by the Radiographic Global Impression of Change global score. All patients who received at least one dose of treatment were included in the primary and safety analyses. Only 61 patients were enrolled out of 122 patients who were assessed. Of these, 32 (18 girls, 14 boys) were randomly assigned to continue receiving conventional therapy and 29 (16 girls, 13 boys) to receive burosumab. For the primary endpoint at week 40, patients in the burosumab group had significantly greater improvement in Radiographic Global Impression of Change global score than did patients in the conventional therapy group (least squares mean +1.9 [SE 0.1] with burosumab vs +0.8 [0.1] with conventional therapy; difference 1.1, 95% CI 0.8-1.5; p<0.0001). Treatment-emergent adverse events considered possibly, probably, or definitely related to treatment by the investigator occurred more frequently with burosumab (17 [59%] of 29 patients in the burosumab group vs 7 [22%] of 32 patients in the conventional therapy group). The investigators concluded that significantly greater clinical improvements were shown in rickets severity, growth, and biochemistries among children treated with burosumab compared to conventional therapy.

A randomized, open-label study (NCT 02163577) in 52 prepubescent XLH patients compared burosumab administered every 2 weeks versus every 4 weeks. Upon completion of a 16-week dose titration, patients were administered burosumab every 2 weeks for 48-weeks. No study patients discontinued burosumab and completed at least 64 weeks of the study. Patient dosing was individualized to achieve a target fasting serum phosphorus concentration of 3.5 to 5.0 mg/dL based on the fasting phosphorus level the day of dosing. Twenty-six of 52 patients received burosumab every two weeks up to a maximum dose of 2 mg/kg. The average dose was 0.73 mg/kg (range: 0.3, 1.5) at week 16, 0.98 mg/kg (range: 0.4, 2.0) at week 40 and 1.04 mg/kg (range: 0.4, 2.0) at week 60. The other 26 patients received burosumab every four weeks. At the beginning of the study, the mean age of patients was 8.5 years with 46% male. Regarding treatment with oral phosphate and active vitamin D analogs, 96% of study participants had received these for a mean (SD) duration of 7 (2.4) years. In addition, discontinuation of oral phosphate and active vitamin D analogs occurred prior to study enrollment. Radiographic evidence of rickets was observed in 94% of patients at baseline. In this study, patients receiving burosumab experienced a mean (SD) increase in serum phosphorus levels from 2.4 (0.40) at baseline to 3.3 (0.40) and 3.4 (0.45) mg/dL at week 40 and week 64 in the patients who received burosumab every 2 weeks. The 10-point Thacher Rickets Severity Score (RSS) and the 7-point Radiographic Global Impression of Change (RGI-C) were used to evaluate rickets. After 40 weeks of therapy, mean total RSS decreased from 1.9 to 0.8 and the mean RGI-C Global score was +1.7 in patients receiving burosumab every two weeks. Eighteen out of 26 patients achieved an RGI-C score of  $\geq +2.0$ . These findings were maintained at week 64.<sup>1,4</sup>

A 64-week open-label study (NCT 02750618) was conducted in 13 XLH patients age 1 to 4 years old. Study patients received burosumab at a dose of 0.8 mg/kg every two weeks with titration up to 1.2 mg/kg based on serum

**Formatted:** Font: Bold

phosphorus. No study participants discontinued burosomab. The mean age of patients was 2.9 years at study entry. At baseline, all study participants had radiographic evidence of rickets and had received oral phosphate and active vitamin D analogs for a mean (SD) duration of 16.9 (13.9) months. Discontinuation of oral phosphate and active vitamin D analogs occurred prior to study enrollment. At week 40, patients experienced an increased mean (SD) serum phosphorus levels from 2.5 (0.28) mg/dL at baseline to 3.5 (0.49) mg/dL. After 40 weeks of treatment, mean total RSS decreased from 2.9 to 1.2 and the mean (SE) RGI-C Global score was +2.3 (0.08). All 13 patients achieved a RGI-C global score  $\geq$  +2.0. The mean (SE) lower limb deformity as assessed by RGI-C, using standing long leg radiographs, was +1.3 (0.14).<sup>1,5</sup>

A randomized, double-blind, placebo-controlled study (NCT 02526160) in 134 adult XLH patients was completed. Burosomab was administered at a dose of 1 mg/kg every 4 weeks. At study entry, the patient age ranged from 16 to 66 years, with a mean of 40 years. The average age of diagnosis was 9 years and 81% of patients had received conventional therapy before the age of 18, for an average of approximately 12 years. 69% of patients had used phosphate and/or active vitamin D within 2 years of study baseline. At baseline, all patients had skeletal pain associated with XLH or osteomalacia. The baseline mean (SD) serum phosphorus concentration was below the lower limit of normal at 1.98 (0.31) mg/dL. Oral phosphate and active vitamin D analogs were not allowed during the study with one patient in the burosomab group discontinued treatment. Through week 24, a total of 94% of patients receiving burosomab achieved a serum phosphorus level above the lower limit of normal compared to 8% in the placebo group. Assessment of active fracture/pseudofractures at week 24 demonstrated a higher rate of complete healing in the group receiving burosomab compared to placebo. During the study, a total of 6 new fractures or pseudofractures appeared in 68 patients receiving burosomab, compared to 8 new abnormalities in 66 patients receiving placebo. The FDA conducted its own analysis in order to examine pain medication usage during burosomab treatment. The FDA determined that there is insufficient evidence to support that burosomab decreased use of pain medication during therapy. The FDA stated that it is possible that as longer term data is collected, a significant reduction in pain medication may become evident.<sup>1,5</sup>

A 48-week, open-label, single-arm study (NCT 02537431) was completed in 14 adult XLH patients to determine the effects of burosomab on improvement of osteomalacia as based on histologic and histomorphometric evaluation of iliac crest bone biopsies. Treatment was 1 mg/kg burosomab every four weeks. At study entry, the mean age of patients was 40 years (range 25 to 52 years) and 43% were male. Oral phosphate and active vitamin D analogs were not allowed during the study. After 48 weeks of treatment, healing of osteomalacia was observed in ten patients as demonstrated by decreases in Osteoid volume/Bone volume from a mean (SD) score of 26% (12.4) at baseline to 11% (6.5), a change of -57%. Osteoid thickness declined in eleven patients. Mineralization lag time) declined in 6 patients from a mean (SD) of 594 (675) days to 156 (77) days, a change of -74%.<sup>1,5</sup>

The pharmacokinetics, efficacy, and safety profile of burosomab was evaluated in a Phase 3 randomized, double blind, placebo controlled trial. The primary endpoint was the proportion of subjects achieving mean serum phosphate above 2.5 mg/dL at the dose interval mid-points of the dose interval between baseline and week 24. 94.1% of burosomab-treated subjects vs 7.6% of placebo-treated subjects achieved mean serum phosphorus  $>$  the lower limit of normal at mid-point of the dose interval, averaged across dose cycles ( $P < 0.0001$ ). At week 24, treatment was associated with healing of active fractures as well as pseudofractures in 44% of patients in the treatment group compared to 18% in the placebo group. The overall safety profile of patients on burosomab was similar to that of placebo.<sup>2</sup>

**Burosomab has been evaluated in two studies enrolling a total of 27 patients with tumor induced osteomalacia (TIO). A single-arm open-label study (NCT 02304237) enrolled 14 adult patients with a confirmed diagnosis of FGF23-related hypophosphatemia produced by an underlying tumor that was not amenable to surgical excision or could not be located. Patients received burosomab every 4 weeks at a weight based starting dose of 0.3 mg/kg that was titrated to achieve a fasting serum phosphorus level of 2.5 to 4.0 mg/dL. The mean dose was 0.83 mg/kg at Week 20, 0.87 mg/kg at Week 48, 0.77 mg/kg at Week 96 and 0.71 mg/kg at Week 144. Burosomab increased mean (SD) serum phosphorus levels from 1.60 (0.47) mg/dL at baseline to 2.64 (0.76) mg/dL averaged across the midpoint of dose intervals through Week 24 with 50% of patients (7/14) achieving a mean serum phosphorus level above the lower limit of normal (LLN) averaged across the midpoint of dose intervals through Week 24. Increase in the mean serum phosphorus concentrations was sustained near or above the LLN through Week 144. <sup>99m</sup>Technetium-labelled whole body bone scans were performed at baseline and subsequent timepoints during the study on all 14 patients. At baseline, all patients had areas of tracer uptake with a total of 249 bone abnormalities across 14 patients. The number of areas of tracer uptake decreased from Week 48 through Week 144, suggesting healing of the bone abnormalities.**

Formatted: Font: Bold

**A single-arm open-label study (NCT 02722798 assessed burosomab in 13 adult patients with a confirmed diagnosis of TIO received. Patients received burosomab every 4 weeks at a weight based starting dose of 0.3 mg/kg that was titrated to achieve a fasting serum phosphorus level of 2.5 to 4.0 mg/dL. The mean (SD) dose was 0.91 (0.59) mg/kg at Week 48, and 0.96 (0.70) mg/kg at Week 88. Burosomab increased**

Formatted: Font: Bold

mean (SD) serum phosphorus levels from 1.62 (0.49) mg/dL at baseline to 2.63 (0.87) mg/dL averaged across the midpoint of dose intervals through Week 24 with 69% of patients (9/13) achieving a mean serum phosphorus level above the LLN averaged across the midpoint on dose interval through Week 24. Mean serum phosphorus concentrations were sustained above LLN through Week 88.

#### U.S. FOOD AND DRUG ADMINISTRATION (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Crysvita is a fibroblast growth factor 23 (FGF23) blocking antibody indicated for the treatment of X-linked hypophosphatemia (XLH) in adult and pediatric patients 6 months of age and older. and for the treatment of FGF23-related hypophosphatemia in tumor-induced osteomalacia (TIO) associated with phosphaturic mesenchymal tumors that cannot be curatively resected or localized in adult and pediatric patients 2 years of age and older<sup>1</sup>

Formatted: Font: Bold

#### CENTERS FOR MEDICARE AND MEDICAID SERVICES (CMS)

Medicare does not have a National Coverage Determination (NCD) for CRYSVITA® (burosumab-twza). Local Coverage Determinations (LCDs)/Local Coverage Articles (LCAs) do not exist at this time.

In general, Medicare may cover outpatient (Part B) drugs that are "incident to" a physician's service provided that the drugs are not usually self-administered by the patients who take them. Refer to the Medicare Benefit Policy Manual, Chapter 15, §50 - Drugs and Biologicals. (Accessed March 20July 17, 2020)

Formatted: Font: Bold

Formatted: Font: Bold

#### REFERENCES

1. Crysvita [prescribing information]. Novato, CA: Ultragenyx Pharmaceutical; September 2019June 2020.
2. Osteocyte regulation of phosphate homeostasis and bone mineralization underlies the pathophysiology of the heritable disorders of rickets and osteomalacia. Feng JQ, Clinkenbeard EL, Yuan B, White KE, Drezner MK SOBone. 2013 Jun;54(2):213-21. Epub 2013 Feb 9.
3. Carpenter TO, Imel EA, Holm IA, et al. A clinician's guide to X-linked hypophosphatemia. J Bone Miner Res. 2011;26(7):1381-1388.
4. Carpenter TO, Whyte MP, Imel EA, et al. Burosumab therapy in children with X-linked hypophosphatemia. N Engl J Med 2018;378:1987-1998.
5. Food and Drug Administration/Center for Drug Evaluation and Research. Crysvita summary review. [https://www.accessdata.fda.gov/drugsatfda\\_docs/nda/2018/761068Orig1s000MultidisciplineR.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/nda/2018/761068Orig1s000MultidisciplineR.pdf). Accessed April 8, 2019.
6. Scheinman SJ, Carpenter T, Drezner MK. Hereditary hypophosphatemic rickets and tumor-induced osteomalacia. Hoppin AG, ed. UpToDate. Waltham, MA: UpToDate Inc. <https://www.uptodate.com>. Accessed March 18, 2020.
7. Imel EA, Glorieux FH, Whyte MP, et al. Burosumab versus conventional therapy in children with X-linked hypophosphataemia: a randomised, active-controlled, open-label, phase 3 trial. Lancet. 2019 Jun 15;393(10189):2416-2427.
8. Agus ZS. Oncogenic hypophosphatemic osteomalacia. Kidney Int. 1983;24(1):113-123. doi:10.1038/ki.1983.133
9. Folpe AL, Fanburg-Smith JC, Billings SD, et al. Most osteomalacia-associated mesenchymal tumors are a single histopathologic entity: an analysis of 32 cases and a comprehensive review of the literature. Am J Surg Pathol. 2004;28(1):1-30. doi:10.1097/00000478-200401000-00001

Shane E, Parisien M, Henderson JE, et al. Tumor-induced osteomalacia: clinical and basic studies. J Bone Miner Res. 1997;12(9):1502-1511. doi:10.1359/jbmr.1997.12.9.1502

#### POLICY HISTORY/REVISION INFORMATION

Date	Action/Description
TBD	<u>Updated Coverage rationale criteria with new indication. Added new ICD-10 codes with new indication. Updated Background information. Updated Clinical Evidence Section. Updated FDA Section. Updated CMS Statement section. Updated references.</u>

Formatted: Font: Bold

Formatted: Font: Bold