

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.



Virtual Upper Gastrointestinal Endoscopy (for Louisiana Only)

Policy Number: CS130LA.#K
Effective Date: February 1, 2024TBD

[Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Applicable Codes	1
Description of Services	2
Clinical Evidence	2
U.S. Food and Drug Administration	5
References	6
Policy History/Revision Information	6
Instructions for Use	7

Application

This Medical Policy only applies to the state of Louisiana.

Coverage Rationale

Virtual upper gastrointestinal (GI) endoscopy using **three-dimensional (3-D) computed tomography (CT)**, or 3-D magnetic resonance imaging (**MRI**) is unproven and not medically necessary for detecting and evaluating upper **gastrointestinal GI** lesions due to insufficient evidence of efficacy.

Formatted: Font: UHC Sans Medium

Formatted: Font: UHC Sans Medium

Formatted: Font: UHC Sans Medium

Formatted: Font: UHC Sans Medium

Formatted: Don't keep with next

Formatted: Font: UHC Sans Medium

Formatted: Font: UHC Sans Medium

Formatted: Font: UHC Sans Medium

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

CPT Code	Description
76497	Unlisted computed tomography procedure (e.g., diagnostic, interventional)
76498	Unlisted magnetic resonance procedure (e.g., diagnostic, interventional)

CPT® is a registered trademark of the American Medical Association

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Description of Services

Virtual upper gastrointestinal (GI) endoscopy is a noninvasive procedure that uses three-dimensional imaging and computed tomography (CT) to capture detailed pictures of the inside surfaces of organs (e.g., organs of the gastrointestinal (GI) tract). Magnetic resonance imaging (MRI) can also be used to perform virtual upper GI endoscopy. Virtual endoscopy is proposed as a means to determine the cause of symptoms, such as nausea, gastric reflux, abdominal pain, and unexplained weight loss; in identifying esophageal varices, inflammation, ulcers, precancerous conditions, and hernias; and in gastric cancer preoperative staging.

Individuals undergoing virtual upper gastrointestinal GI endoscopy usually do not need anesthesia or sedation. As this is an imaging procedure, physicians have the capability to modify the captured pictures by magnifying the images or altering the image angles. Disadvantages of virtual upper gastrointestinal GI endoscopy include the difficulty in showing fine detail compared to a standard endoscopy procedure; exposure to CT scan radiation, and the inability to perform a biopsy during the procedure. If a lesion is found, conventional upper GI endoscopy is necessary for ~~excision or~~ biopsy.

Clinical Evidence

The body of ~~evidence in the~~ published, peer reviewed, scientific literature ~~evidence~~ evaluating virtual upper gastrointestinal (GI) endoscopy using ~~three-dimensional (3-D)~~ computed tomography (CT) or 3-D magnetic resonance imaging (MRI) ~~is mainly~~ ~~constituted~~ ~~primarily consists~~ of observational studies and case reports. There ~~were~~ ~~are~~ no randomized controlled trials (RCTs) available to assess clinical utility. ~~Randomized controlled~~ ~~Studies are limited with small number of participants, lack of comparison, standard methods, and lack of studies~~ ~~designed to establish the clinical utility on long term patient and oncologic outcomes. RCTs~~ comparing ~~it~~ ~~virtual upper GI endoscopy~~ to conventional upper GI endoscopy are needed to determine ~~its~~ clinical value as well as safety and efficacy.

Formatted: Font: UHC Sans Medium

[Kumar et al. \(2023\) conducted a cross-sectional study to determine how effective multi-detector computed tomography \(MDCT\) virtual esophagography was at detecting and grading esophageal varices \(EV\) in individuals with chronic cirrhotic liver disease \(CLD\). The study included 62 individuals with CLD who underwent CT scan followed by endoscopic examination of the esophagus. CT scans were obtained after esophageal distention and processed using two-dimensional \(2-D\) and 3-D software that allowed processing of multiplanar reconstructions and volume renderings by using surface-shaded transparent and virtual endoscopy modes. EVs were graded at MDCT and at endoscopic examination using a standardized classification system. The study results revealed CT virtual esophagography had a sensitivity, specificity, positive predictive value, negative predictive value, and diagnostic accuracy of 86%, 90%, 98%, 56%, and 87%, respectively. Substantial agreement between endoscopic examination and CT virtual esophagography was determined to be statistically significant \(k = 0.616; p < 0.001\). However, agreement for the detection of grade I esophageal varices was poor \(k = 0.102\), possibly due to these lesions being less conspicuous on CT. The authors concluded the study results have the potential to change the way CLD is managed and encourage further research. A multicenter study with a large number of individuals is needed to support these findings. The authors noted several study limitations including small sample size and single center design. The study also included a small number of individuals with grade 0 and grade I EVs.](#)

In a 2022 retrospective study of individuals admitted to the emergency department, Kim et al. aimed to investigate the diagnostic value of ~~multi-detector computed tomography (MDCT)~~ for individuals with suspected upper gastrointestinal GI bleeding (UGIB). ~~In total, The study included 386 individuals were~~ ~~and~~ compared ~~between contrast~~ ~~enhanced~~ ~~abdominopelvic~~ ~~MCT~~ ~~MDCT~~ to endoscopy, ~~and the~~ ~~The~~ performance of MDCT in identifying the status, location of origin, and etiology of UGIB was analyzed. The outcomes measured were sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and accuracy. ~~The study results revealed~~ MDCT was able to accurately identify 32.9% [21.9–43.9, 95% confidence interval (CI)] for individuals with active bleeding, 27.4% (18.9–35.9, 95% CI) for individuals with recent bleeding, and 94.8% (91.8–97.8, 95% CI) for individuals without bleeding evidence (P < 0.001). MDCT showed an accuracy of 60.9%, 60.6%, and 50.9% in identifying bleeding in the esophagus, stomach, and duodenum ~~in that order, respectively~~ (P = 0.4028). ~~The~~ ~~Additionally, the~~ accuracy in differentiating ulcerative, cancerous, and variceal bleeding was 58.3%, 65.9%, and 56.6%, respectively (p = 0.6193). The authors concluded ~~that the~~ MDCT

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

has limited use as a supportive screening method to identify the presence of gastrointestinal bleeding UGIB. Limitations of the study noted by the authors included the retrospective nature of the research design. Unidentified confounders and selection bias may have been introduced. Non-blinded data interpretation carried out by different radiologists may have also generated variation in the reported findings. The study was conducted at a single-tertiary hospital. Multi-center studies are needed. False positive and false negative results could have been produced due to the widely diverse characteristics of UGIB.

Wani et al. (2021) in an observational study assessed/evaluated the diagnostic accuracy of MDCT using hydro and gaseous distension of stomach superseding endoscopic ultrasound in tumor (T) and nodal (N) staging of gastric cancer with an attempt to differentiate between early and advanced gastric carcinomas. A total of in this observational study, 160 individuals with endoscopically diagnosed and biopsy-proven gastric cancer were subjected to MDCT after adequate gaseous and hydro-distention of stomach. Multi-planar reformatted (MPR) and virtual gastroscopy (VG) images were also obtained. Gastric lesions were categorized into T1 to T4 stages with N staging from N0 to N3. Preoperative CT findings were correlated with histopathological findings. In the study, general results revealed the diagnostic accuracy of T staging was 82.5% (132/160) with an accuracy of 75% (120/160) for N staging. The diagnostic accuracy of CT for early gastric carcinoma in the study was 93.75%, with high specificity of 96%, but low sensitivity of 66.7%. The authors concluded that MDCT after gaseous and hydro-distension in MDCT of the stomach is an excellent method for near accurate preoperative T staging of gastric cancer. Nevertheless, CT has a limited role in the N staging of gastric cancer. This study also suggested that the use of virtual gastroscopy VG and MPR images helps in better detection of early gastric cancers/cancer. Larger more robust studies are needed to support the clinical utility of virtual gastroscopy VG in staging gastric cancer.

Almeida et al. (2018) evaluated the accuracy of multidetector computed tomography used in conjunction MDCT with virtual gastroscopy a stomach protocol in staging gastric cancer. The study included 14 participants who underwent computed tomography in a 16 channel scanner for preoperative staging of gastric adenocarcinoma between September 2015 in a 16-channel CT scanner. The stomach protocol included a period of fasting, followed by an intravenous antispasmodic agent, and December 2016. All images ingestion of two effervescent salt envelopes diluted in water just prior to CT. All participants were analyzed by the same radiologist then treated with extensive abdominal cancer imaging experience. The method sensitivity a total or partial gastrectomy. Biopsies were taken during surgery and submitted for histopathological analysis. Sensitivity, specificity, and accuracy were calculated by comparing it MDCT with the pathology result. All participants underwent partial or total gastrectomy. The mean age was 61.5 years, and 53.8% were male. The gastric lesions were classified as T1/T2 in 35.7% of the cases, as T3 in 28.5% of the cases, and as T4 in 35.7% of the cases. Eleven people/participants (68.7%) had suspicious (N₊positive) lymph nodes. The accuracy of the T1/T2, T3, T4, and lymph node staging tests was 85%, 78%, 90%, and 78%, respectively. The respective sensitivity and specificity values were 71% and 100% for T1/T2, 66% and 81% for T3, 100% and 90% for T4, and 88% and 60% for lymph nodes. The authors concluded that the multidetector computed tomography MDCT with a stomach protocol, used in conjunction with virtual gastroscopy, shows VG, showed good accuracy in the tumor and lymph node staging of gastric adenocarcinoma. This study is/was limited by a small sample size.

Kim et al. (2012) assessed the diagnostic accuracy of different reconstruction techniques using MDCT for gastric cancer detection compared with 2D-D axial CT. The authors performed CT examinations were performed in 104 consecutive individuals with gastric cancer, and the control group was composed of 35 individuals without gastric disease. All gastric cancer was pathologically proven by endoscopy and surgery. Among the 104 of those individuals with gastric cancer, 63 had early gastric cancer (EGC). Two radiologists, using a commercially available, 3-D workstation, retrospectively and independently interpreted the axial CT and three different reconstruction techniques, including. The techniques included multiplanar reformation (MPR), transparent imaging (TI), and virtual gastroscopy (VG). Overall, VG had significantly better performance than 2D-D axial CT. The sensitivity (p = 0.016), Sensitivity and specificity were as follows: 76.7% and 82.9% infor axial CT; 79.6% and 85.7% infor MPR; 91.3% and 80% infor TI; and 95.1% and 74.3% infor VG. For early gastric cancer, VG had significantly better performance than both 2D-D axial CT (P = 0.006) and MRP. The sensitivity (P = 0.038). Sensitivity and specificity were as follows: 62.9% and 82.9% infor axial CT; 67.7% and 85.7% infor MPR; 85.5% and 80% infor TI; and 91.9% and 74.3% infor VG. The There was substantial inter-observer agreement showed substantial agreement. The authors concluded that among the different reconstruction techniques, VG is a promising method for accurately detecting gastric cancer and is especially useful for early gastric cancer compared with 2D-D axial CT. Interpretation of these findings is limited due to the study's single institution, retrospective design, and relatively small patient sample size.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Okten et al. (2012) assessed the role of MDCT with ~~multiplanar (3-D) reconstruction (MPR)~~ and VG for the detection and differentiation of gastric subepithelial masses (~~SEM/SEMs~~) by comparison with endoscopic ultrasonography (EUS). Forty-one individuals with a suspected SEM were evaluated using EUS and MDCT. MDCT findings were analyzed based on the consensus of two radiologists blinded to the EUS findings. EUS and MDCT results were compared with histopathology for the pathologically proven lesions. For the non-pathologically proven lesions, MDCT results were compared with EUS. ~~Among~~**The study results revealed among the 41 participants/individuals, 34 SEM were detected using EUS. For the detection of SEM/SEMs with MDCT, a sensitivity of 85.3%, a specificity of 85.7%, a positive predictive value of 96.7%, and a negative predictive value of 54.5% were calculated. The overall accuracy of MDCT for detecting and classifying the SEM/SEMs was 85.3% and 78.8%, respectively. The authors concluded that MDCT with MPR and VG is a valuable method for evaluating SEMs. The authors stated that specific/Specific MDCT criteria for various SEMs may help make/in making an accurate diagnosis. These findings require confirmation in a larger/The authors noted several study limitations including small sample size. Additionally, MDCT was performed in individuals with a high probability of having a gastric SEM. This may have led to high accuracy rates for the detection of lesions.**

Moschetta et al. (2012) assessed the diagnostic accuracy of VG obtained by 320-row ~~computed tomography (CT)~~ examination in differentiating benign from malignant gastric ulcers (~~GUs~~). Forty-nine individuals with endoscopic and histological diagnoses of ~~GU~~**gastric ulcers** underwent CT examination. Based on morphological features, ~~GU~~**gastric ulcers** were subdivided into benign or malignant forms by two blinded radiologists. CT results were then compared with endoscopic and histological findings, having the latter as the reference standard. ~~Thirty-five~~**The study results revealed 35** out of 49 participants (71%) were affected by malignant ulcers, while in the remaining 14 cases, a diagnosis of benign ~~GU~~**gastric ulcers** was made. VG showed diagnostic accuracy, sensitivity, and specificity values of 94%, 91%, and 100%, respectively, in differentiating benign from malignant ~~gastric~~**gastric** ulcers. Almost perfect agreement between the two readers was found. The authors concluded that ~~the~~**CT VG improves the gastric ulcer identification of GUs** and allows the differentiation of benign from malignant forms. ~~The significance of this~~**authors noted several study is limited by a-limitations including small sample size. Additionally, only individuals with endoscopically diagnosed lesions were included in the study. Thus, results for the detection of gastric cancer may be biased. The value of VG in the screening of gastric cancer cannot be provided. Furthermore, a direct comparison with optical gastroscopy was not performed.**

~~In a prospective trial,~~ Ulla et al. (2010) evaluated the usefulness of ~~the~~**Pneumo-64-MDCT (PnCT64) technique** in the presurgical characterization of esophageal neoplasms in correlation with surgical findings ~~for~~. **The prospective study included 50 individuals with a diagnosis of an endoscopically confirmed esophageal neoplasm. A 14 French diagnosis. In order to optimize tumor visualization in the esophageal wall and in the gastroesophageal (GE) junction, individuals underwent insertion of a transoral Foley catheter was used trans-orally in all participants. Air was then instilled through the catheter to achieve esophageal distension. A 64-row MDCT scan was then performed, and the tumor was characterized according to scope, shape, and anatomic location using multiplanar ~~3D-3-D~~ reconstructions and virtual endoscopy. Wall infiltration and the presence of adenopathies were analyzed. ~~44/50 people,~~**Adequate GE distension was achieved in all individuals. The study results revealed that PnCT64 showed wall thickening was observed in 44/50 (84%) individuals and 34/50 regional adenopathies were found in 34/50(68%) individuals. In 29/50 individuals, the lesion was found in the lower third and the gastroesophageal/GE junction. The surgical correlation for wall infiltration was 85.7%. The investigators****Surgery was performed in 35/50 individuals. A comparison of PnCT64 with histopathological findings showed a surgical correlation for wall infiltration in 30/35 (85.7%) individuals. In 5/35 individuals who underwent surgery, PnCT64 did not show evidence of wall infiltration. Of these, 3 were T1 tumors, and the other 2 were T2 tumors.****Surgical correlation for the detection of adenopathies reached 75%. The authors** concluded that PnCT64 is useful and safe for identifying esophageal wall thickening and presurgical characterization. Optimal distension allows the definition of both upper and lower borders of the tumors located in the ~~gastroesophageal/GE~~**gastroesophageal/GE** junction, which is of utmost importance to determine the surgical approach. ~~These findings need confirmation~~**The authors noted study limitations including the lack of comparison between PnCT64 and other imaging modalities. There may have also been bias introduced in a larger investigation****the analysis, as the radiologists were not masked to the endoscopically confirmed esophageal neoplasm diagnosis.****

Chen et al. (2009) retrospectively compared CT VG to conventional optical ~~gastro-endoscopy~~**gastroendoscopy** when determining differences between benign and malignant gastric ulcers. ~~Gastric ulcers in~~**The study included 115 individuals (mean age, 64.7 years; range, 31-86 years; 61 men, 54 women)****with gastric ulcers who** were evaluated ~~by~~**using** endoscopy and VG. ~~At the histopathologic~~

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Histopathologic examination revealed 39 gastric ulcers were benign, while and 76 gastric ulcers were malignant. For the overall diagnosis of malignant gastric ulcers, VG and endoscopy had a sensitivity of 92.1% and 88.2% sensitivities, respectively, for the overall diagnosis of malignant gastric ulcers and specificities of . Specificity was 91.9% and 89.5%, respectively, for the overall diagnosis of malignant gastric ulcers. Endoscopy was more sensitive in depicting malignancy according to ulcer base (85.5% vs. 68.4%), (P = .034), and VG was more specific in depicting malignancy according to ulcer margin (78.4% vs. 63.2%), (P = .034). The authors concluded that VG and endoscopy were almost equally useful in distinguishing between malignant and benign gastric ulcers. These findings need The authors noted some study limitations including selection bias. There was also great variation noted in the elapsed time between the endoscopic and VG examinations. Additional prospective, RCTs comparing VG and optical endoscopy are needed to investigate the overall efficacy of VG.

Chen et al. (2007) prospectively evaluated the accuracy of MDCT for the preoperative staging of gastric cancer by using surgical and histopathologic results as reference standards. The study included 55 consecutive individuals who were diagnosed with gastric cancer. Prior to MDCT, individuals underwent endoscopic biopsy that provided histologic confirmation in of gastric cancer. MDCT included acquisition of VG images after air distention and contrast material-enhanced dynamic transverse and MPR images after water distention. All individuals underwent partial or complete gastrectomy. CT findings were compared with surgical and histopathologic results. The study results revealed detection rates of primary tumors with transverse images, MPRs, and combinations of MPR and virtual gastroscopy images were 91% (50 of 55), 96% (53 of 55), and 98% (54 of 55), respectively. The overall accuracy in assessment of tumor invasion into the gastric wall (T stage) was significantly better with MPR images (89%) (49 of 55) than with transverse images (73%) (40 of 55) (P < .01). Additionally, the overall accuracy for lymph node (N) staging was 78% (43 of 55) with MPR images and 71% (39 of 55) with transverse images. This difference was not significant (P = .103). The authors concluded MDCT with combined water and air distention can improve the accuracy of preoperative staging of gastric cancer. MPRs also yield significantly better overall accuracy than transverse images for tumor staging, but not for lymph node staging. The authors noted some study limitations including small sample size. Studies with a larger study number of individuals are needed.

Clinical Practice Guidelines

European Society of Gastrointestinal Endoscopy (ESGE)

The updated 2022 ESGE guideline for endoscopic submucosal dissection (ESD) for superficial gastrointestinal GI lesions recommends that the evaluation of superficial gastrointestinal GI lesions should be made by an experienced endoscopist using high-definition white-light and chromoendoscopy (virtual or dye-based), and validated classifications when available.

(Strong recommendation, high-quality evidence.) ESGE does not recommend routine performance of endoscopic ultrasonography (EUS), computed tomography (CT), magnetic resonance imaging (MRI), or positron emission tomography CT (PET-CT) before endoscopic resection (ER). (Strong recommendation, moderate quality evidence.)

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Imaging devices used to create virtual endoscopy images are classified under the following ~~product codes:~~ **Product Codes: JAK, LLZ (system, image processing, radiological); JAK (system, x ray, tomography, computed); and , or LNH (system, nuclear magnetic resonance imaging). Note that the devices listed under these codes are general imaging devices and may not be indicated explicitly for virtual upper gastrointestinal endoscopy. ~~To locate~~ Refer to the FDA's 510(k) Premarket Notification website for information regarding marketing clearance information for a specific device or manufacturer, search the following website by product and/or manufacturer name: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>. (Accessed July 14, 2023, 2024)**

~~Two software packages that can be used to convert two-dimensional helical computed tomography (CT) images into three-dimensional images are:~~
~~The Magic View package for use with the Somatom Plus 4 scanner [Siemens Medical Solutions, Erlangen, Germany, 510k (K964747) approval received on February 10, 1997] additional information at: (Accessed July 14, 2023), and;~~

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

~~The CT Colonography/Navigator 2 package (GE Medical Systems, Buc Cedex, France. 510k (K012313) approval received on August 7, 2001) additional information at: (Accessed July 14, 2023), and for use with the Hi-Speed Advantage CT Scanner (GE Medical Systems, Milwaukee, WI. 510k (K940606) approval received on August 23, 1994) additional information at: (Accessed July 14, 2023).~~

References

Almeida MFA, Verza L, Bitencourt AGV, et al. Computed tomography with a stomach protocol and virtual gastroscopy in the staging of gastric cancer: an initial experience. Radiol Bras. 2018 Jul-Aug;51(4):211-217.

Chen CY, Hsu JS, Wu DC, et al. Gastric cancer: preoperative local staging with 3D multi-detector row CT--correlation with surgical and histopathologic results. Radiology. 2007 Feb;242(2):472-82.

[Chen CY, Kuo YT, Lee CH, et al. Differentiation between malignant and benign gastric ulcers: CT virtual gastroscopy versus optical gastroendoscopy. Radiology. 2009 Aug;252\(2\):410-7.](#)

Kim D, Kim JH, Ko DR, et al. Usefulness of contrast-enhanced multi-detector computed tomography in identifying upper gastrointestinal bleeding: A retrospective study of patients admitted to the emergency department. PLoS One. 2022 Apr 7;17(4):e0266622.

Kim JH, Eun HW, Hong SS, et al. Gastric cancer detection using MDCT compared with 2D axial CT: diagnostic accuracy of three different reconstruction techniques. Abdom Imaging. 2012 Aug;37(4):541-8.

[Kumar R, Nandolia KK, Sharma P, et al. Computed tomography virtual oesophagography for the grading of oesophageal varices in cirrhotic liver disease patients with upper gastrointestinal endoscopic examination as the gold standard: a diagnostic validation study. Pol J Radiol. 2023 Apr 5;88:e187-e193.](#)

Moschetta M, Scardapane A, Telegrafo M, et al. Differential diagnosis between benign and malignant ulcers: 320-row CT virtual gastroscopy. Abdom Imaging. 2012 Dec;37(6):1066-73.

Okten RS, Kacar S, Kucukay F, et al. Gastric subepithelial masses: evaluation of multidetector CT (multiplanar reconstruction and virtual gastroscopy) versus endoscopic ultrasonography. Abdom Imaging. 2012 Aug;37(4):519-30.

Pimentel-Nunes P, Libânio D, Bastiaansen BAJ, et al. Endoscopic submucosal dissection for superficial gastrointestinal lesions: European Society of Gastrointestinal Endoscopy (ESGE) Guideline - Update 2022. Endoscopy. 2022 Jun;54(6):591-622. ~~Available at: Accessed July 17, 2023.~~

Ulla M, Cavadas D, Muñoz I, et al. Esophageal cancer: pneumo-64-MDCT. Abdom Imaging. 2010 Aug;35(4):383-9.

Wani AH, Parry AH, Feroz I, et al. Preoperative staging of gastric cancer using computed tomography and its correlation with histopathology with emphasis on multi-planar reformations and virtual gastroscopy. J Gastrointest Cancer. 2021 Jun;52(2):606-615.

Policy History/Revision Information

Date	Summary of Changes
TBD	Supporting Information <ul style="list-style-type: none">• Updated Description of Services, Clinical Evidence, FDA, and References sections to reflect the most current information• Archived previous policy version CS130LA.J

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual[®] criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

DRAFT