

Clinical Policy: Azacitidine (Vidaza)

Reference Number: LA.PHAR.387

Effective Date: 03.16.23

Last Review Date: 01.21.25~~05.06.24~~

Line of Business: Medicaid

[Coding Implications](#)[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

****Please note: This policy is for medical benefit****

Description

Azacitidine (Vidaza®) is a nucleoside metabolic inhibitor.

FDA Approved Indication(s)

Vidaza is indicated for the treatment of:

- Adult patients with the following French-American-British (FAB) myelodysplastic syndrome (MDS) subtypes: refractory anemia (RA) or refractory anemia with ringed sideroblasts (RARS) (if accompanied by neutropenia or thrombocytopenia or requiring transfusions), refractory anemia with excess blasts (RAEB), refractory anemia with excess blasts in transformation (RAEB-T), and chronic myelomonocytic leukemia (CMML).
- Pediatric patients aged 1 month and older with newly diagnosed juvenile myelomonocytic leukemia (JMML).

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that azacitidine, ~~and~~ Vidaza ~~is~~are medically necessary when the following criteria are met:

Formatted: Font color: Auto

I. Initial Approval Criteria**A. Myelodysplastic Syndromes** (must meet all):

1. Diagnosis of MDS, including JMML;
2. Request is for generic azacitidine or Vidaza;
3. Prescribed by or in consultation with an oncologist or hematologist;
4. One of the following (a or b):
 - a. Age ≥ 18 years;
 - b. Age ≥ 1 month, and request is for JMML;
5. For brand Vidaza requests, member must use generic azacitidine, unless contraindicated or clinically significant adverse effects are experienced;

5-6. Request meets one of the following (a, b, or c):*

- a. For MDS, dose does not exceed one of the following (i or ii):
 - i. Initial: 75 mg/m² per day for 7 days;
 - ii. Maintenance: 100 mg/m² per day for 7 days per 4-week cycle;
- b. For JMML, dose does not exceed one of the following administered daily for 7 days per 28-day cycle, for up to 6 cycles (i or ii):
 - i. Age 1 month to less than 1 year or weighing less than 10 kg: 2.5 mg/kg;
 - ii. Age 1 year and older and weighing 10 kg or greater: 75 mg/m²;
- c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

B. Acute Myeloid Leukemia (Vidaza off-label) (must meet all):

1. Diagnosis of AML;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age ≥ 18 years;
- ~~4. For Onureg requests, member meets all of the following (a, b, c, and d):~~
 - ~~a. Request is for maintenance therapy;~~
 - ~~b. Request is for Prescribed as a single agent therapy;~~
 - ~~c. Member achieved CR or CRi following intensive induction chemotherapy and is either not able or declines to complete intensive consolidation/curative therapy (see Appendix D);~~
 - ~~d. One of the following (i or ii):~~
 - ~~i. Medical justification supports inability to use SC/IV azacitidine (e.g., contraindication to excipients);~~
 - ~~ii. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (see Appendix E);~~
4. For Vidaza requests, prescribed in one of the following ways (a-f):
 - a. As a single agent;
 - b. In combination with Venclexta[®];
 - c. For FLT3-ITD (internal tandem duplication) mutation: In combination with Nexavar[®];
 - d. For IDH1 mutation: In combination with Tibsovo[®];
 - e. For IDH2 mutation: In combination with Idhifa[®];
 - f. For FLT3-ITD or TKD (tyrosine kinase domain) mutation in disease without IDH1 mutation: In combination with Xospata[®];
5. For brand product requests, member must use generic azacitidine, if available, unless contraindicated or clinically significant adverse effects are experienced;

5-6. Request meets one of the following (a, b, or c):*

- a. Onureg: Dose does not exceed 300 mg (1 tablet) per day for 14 days per 4-week cycle;
- ~~a-b~~. Vidaza: Dose does not exceed 100 mg/m² per day for 7 days per 4-week cycle;
- ~~b-c~~. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

Formatted: Page break before, Keep with next, Keep lines together

Formatted: Font: Italic

CLINICAL POLICY

Azacitidine



C. Myelofibrosis (off-label) (must meet all):

1. Diagnosis of advanced phase (i.e., accelerated- or blast-phase) ~~myelofibrosis (MF); myeloproliferative neoplasms;~~
2. Request is for generic azacitidine or Vidaza;
3. Prescribed by or in consultation with an oncologist or hematologist;
4. Age ≥ 18 years;
5. Prescribed as bridging therapy prior to transplant, unless member is not a candidate for transplant;
6. One of the following (a or b):
 - a. Prescribed as a single agent or in combination with Jakafi®, Inrebic®, Ojjaara®, or Vonjo® for palliation of splenomegaly or other disease-related symptoms;
 - b. Prescribed in combination with Venclexta;
7. For brand Vidaza requests, member must use generic azacitidine, unless contraindicated or clinically significant adverse effects are experienced;
- 5-8. Request meets one of the following (a or b):*
 - a. Dose does not exceed 100 mg/m² per day for 7 days per 4-week cycle;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

D. Peripheral T-Cell Lymphoma (off-label) (must meet all):

1. Diagnosis of one of the following peripheral T-cell lymphomas (a, b, or c):
 - a. Angioimmunoblastic T-cell lymphoma;
 - b. Nodal peripheral T-cell lymphoma with TFH phenotype;
 - c. Follicular T-cell lymphoma;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age ≥ 18 years;
4. Disease is relapsed or refractory;
5. Prescribed as a single agent for one of the following (a or b):
 - a. Initial palliative therapy;
 - b. Second-line or subsequent therapy;
- ~~For Onureg requests, one of the following (a or b):~~
 - ~~Medical justification supports inability to use SC/IV azacitidine (e.g., contraindication to excipients);~~
 - ~~Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (see Appendix E);~~
6. For brand product requests, member must use generic azacitidine, if available, unless contraindicated or clinically significant adverse effects are experienced;
7. Request meets one of the following (a, b, or c):*
 - ~~Onureg: Dose does not exceed 300 mg (1 tablet) per day for 14 days per 4-week cycle;~~
 - a. Vidaza: Dose does not exceed 100 mg/m² per day for 7 days per 4-week cycle;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

D-E. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53

Formatted: Font color: Auto

Formatted: Font color: Auto

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving Vidaza for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. For brand product requests, member must use generic azacitidine, if available, unless contraindicated or clinically significant adverse effects are experienced;
- 3.4. If request is for a dose increase, request meets one of the following (a, b, c, or d):*
 - a. Vidaza for MDS: New dose does not exceed 100 mg/m² per day for 7 days per 4-week cycle;
 - b. Vidaza for JMML: New dose does not exceed one of the following administered daily for 7 days per 28-day cycle, for up to 6 cycles (i or ii):
 - i. Age 1 month to less than 1 year or weighing less than 10 kg: 2.5 mg/kg;
 - ii. Age 1 year and older and weighing 10 kg or greater: 75 mg/m²;
 - c. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration: 12 months

Formatted: Font: 12 pt, Not Italic

Formatted: Indent: First line: 0"

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53

III. Diagnoses/Indications for which coverage is NOT authorized:

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – LA.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

AML: acute myelogenous leukemia

ANC: absolute neutrophil count

CLINICAL POLICY

Azacitidine



CMMoL/CMML: chronic myelomonocytic leukemia
 CR: complete response
 CRi: complete response with incomplete hematologic recovery
 FAB: French-American-British
 FDA: Food and Drug Administration
[ITD: internal tandem duplication](#)
 JMML: juvenile myelomonocytic leukemia
 MDS: myelodysplastic syndrome
[TKD: tyrosine kinase domain](#)

MF: myelofibrosis
 NCCN: National Comprehensive Cancer Network
 RA: refractory anemia
 RAEB: refractory anemia with excess blasts
 RAEB-T: refractory anemia with excess blasts in transformation
 RARS: refractory anemia with ringed sideroblasts

Appendix B: Therapeutic Alternatives
 Not applicable

Appendix C: Contraindications/Boxed Warnings:

- Contraindication(s): advanced malignant hepatic tumors (Vidaza only), hypersensitivity to azacitidine (or mannitol for Vidaza only)
- Boxed warning(s): none reported

Appendix D: General Information

The National Comprehensive Cancer Network (NCCN) AML treatment guidelines define morphologic CR in patients that are independent of transfusions as follows:

- Absolute neutrophil count (ANC) > 1,000/mcL (blasts < 5%)
- Platelets ≥ 100,000/mcL (blasts < 5%)

NCCN presents CRi (a variant of CR) for AML as follows based on clinical trial information:

- < 5% marrow blasts
- Either ANC < 1,000/mcL or platelets < 100,000/mcL
- Transfusion independence but with persistence of neutropenia (< 1,000/mcL) or thrombocytopenia (< 100,000/mcL)

Appendix E: States with Regulations against Redirections in Certain Oncology Settings

State	Step-Therapy Prohibited?	Notes
LA	Yes	For stage 4 advanced, metastatic cancer or associated conditions. Exception if "clinically equivalent therapy, contains identical active ingredient(s), and proven to have same efficacy."

V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Azacitidine (Onureg)	AML	300 mg PO QD on days 1 through 14 of each 28-day cycle	300 mg/day for 14 days/cycle

Formatted: Font: Italic

Formatted: Font: Bold, Not Italic

Formatted: Indent: First line: 0"

CLINICAL POLICY

Azacitidine



Drug Name	Indication	Dosing Regimen	Maximum Dose
Azacitidine (Vidaza)	MDS	75 mg/m ² SC or IV infusion QD for 7 days. Repeat cycle every 4 weeks. May increase to 100 mg/m ² (after 2 treatment cycles). Patients should be treated for a minimum of 4 to 6 cycles. Doses may be adjusted or delayed based on hematology lab values, renal function, or serum electrolytes. Continue treatment as long as the patient continues to benefit	100 mg/m ² /day for 7 days/cycle
	JMML	Age 1 month to less than 1 year or weighing less than 10 kg: 2.5 mg/kg Age 1 year and older and weighing 10 kg or greater: 75 mg/m ² Administer IV daily for 7 days in a 28-day cycle, for a minimum of 3 cycles and a maximum of 6 cycles	See dosing regimen

VI. Product Availability

Drug Name	Availability
Azacitidine (Vidaza)	Lyophilized powder in single dose vials/vial: 100 mg

VII. References

1. Onureg Prescribing Information. Summit, NJ: Celgene Corporation; October 2022. Available at: <https://onuregpro.com>. Accessed ~~June 30, 2023~~ August 8, 2024.
2. Vidaza Prescribing Information. Summit, NJ: Celgene Corporation; ~~May 2022~~ January 2024. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/050974s034lbl/2024/050794s036lbl.pdf. Accessed ~~June 30, 2023~~ August 8, 2024.
3. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed August ~~9, 2023~~ 8, 2024.
4. National Comprehensive Cancer Network. Myelodysplastic Syndromes Version ~~4.2023~~ 3.2024. Available at http://www.nccn.org/professionals/physician_gls/pdf/mds.pdf. Accessed ~~July 10, 2023~~ August 8, 2024.
5. National Comprehensive Cancer Network. Acute Myeloid Leukemia Version ~~4.2023~~ 3.2024. Available at http://www.nccn.org/professionals/physician_gls/pdf/aml.pdf. Accessed August ~~4, 2023~~ 8, 2024.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

Formatted: Keep with next

CLINICAL POLICY
Azacitidine



HCPCS Codes	Description
J9025	Injection, azacitidine, 1 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy	02.23	03.16.23
Updated criteria for other diagnoses/indications	06.25.23	10.24.23
Added Onureg to the policy and Onureg specific criteria		
Annual review: removed Onureg from policy, as drug is on PDL with non-preferred LDH Criteria. Added blurb "This policy is for medical benefit" references reviewed and updated	05.06.24	08.20.24
<u>Annual review: Removed any remaining policy mentions of Onureg, as this drug is part of the LDH PDL; revised policy/criteria section to also include generic azacitidine; for all indications where applicable, updated generic redirection to include Vidaza; for AML added requirements regarding usage of Vidaza (single agent and combination) per NCCN; updated off-label criteria for "myelofibrosis" to instead refer to "myeloproliferative neoplasms" and added specific requirements around recommended uses (bridging therapy prior to transplant and use as a single agent or in various combinations) per NCCN; added off-label criteria for peripheral T-cell lymphomas per NCCN; references reviewed and updated. Removal of Appendix E, as LDH previously advised it is not applicable to Medicaid.</u>	01.21.25	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

CLINICAL POLICY

Azacitidine



This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

| ©2025⁴ Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.