

Evolut Clinical Guideline 2006035 for Abdominal Aorta Computed Tomography Angiography with Lower Extremity Runoff

| | | |
|---|--|--|
| Guideline Number: Evolut_CG_ <u>2006035</u> | <u>Applicable Codes</u> | |
| <i>"Evolut" refers to Evolut Health LLC and Evolut Specialty Services, Inc.</i> © 2008 – 202 <u>6</u> 5 All rights Reserved | | |
| Original Date: July 2008 | Last Revised Date: July <u>June</u> 202 <u>5</u> 4 | Implementation Date: January 202 <u>6</u> 5 |

TABLE OF CONTENTS

| | |
|--|-------------------|
| STATEMENT | 3 |
| GENERAL INFORMATION | 3 |
| PURPOSE | 3 |
| INDICATIONS | 3 |
| PERIPHERAL VASCULAR DISEASE | 3 |
| PREOPERATIVE OR POSTOPERATIVE ASSESSMENT | 4 |
| FURTHER EVALUATION OF INDETERMINATE FINDINGS | <u>54</u> |
| IMAGING IN KNOWN GENETIC CONDITIONS | 5 |
| COMBINATION STUDIES FOR KNOWN GENETIC CONDITIONS | 5 |
| Abdominal Aorta CTA with Lower Extremity Runoff / Chest CTA | 5 |
| OTHER COMBINATIONS STUDIES WITH ABDOMINAL AORTA CTA WITH LOWER EXTREMITY RUNOFF | <u>65</u> |
| ABDOMINAL AORTA CTA WITH LOWER EXTREMITY RUNOFF AND CHEST CTA | <u>65</u> |
| CODING AND STANDARDS | <u>65</u> |
| CODES | <u>65</u> |
| APPLICABLE LINES OF BUSINESS | 6 |
| BACKGROUND | 6 |
| CONTRAINDICATIONS AND PREFERRED STUDIES | 6 |
| SUMMARY OF EVIDENCE | <u>96</u> |
| ANALYSIS OF EVIDENCE | <u>107</u> |
| POLICY HISTORY | <u>108</u> |
| LEGAL AND COMPLIANCE | <u>118</u> |
| GUIDELINE APPROVAL | <u>118</u> |
| Committee | <u>118</u> |

| | |
|-------------------------|--------------------|
| DISCLAIMER | <u>118</u> |
| REFERENCES | <u>1340</u> |

STATEMENT

General Information

- *It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.*
- *Where a specific clinical indication is not directly addressed in this guideline, medical necessity determination will be made based on widely accepted standard of care criteria. These criteria are supported by evidence-based or peer-reviewed sources such as medical literature, societal guidelines and state/national recommendations.*
- *The guideline criteria in the following sections were developed utilizing evidence-based and peer-reviewed resources from medical publications and societal organization guidelines as well as from widely accepted standard of care, best practice recommendations.*

Purpose

Computed tomography angiography (CTA) generates images of the blood vessels in the abdomen and lower extremities that can be evaluated for evidence of stenosis or occlusion with the use of ionizing radiation, which is a potential hazard in patients with impaired renal function.

INDICATIONS

Peripheral Vascular Disease ^(1–4)

NOTE: When vascular imaging of the aorta and both legs is required, i.e., CTA aortogram and runoff is desired (sometimes incorrectly requested as Abd/Pelvis CTA & Lower Extremity CTA Runoff), only **ONE** authorization request is required, using CPT Code 75635 (CTA Abdominal Aortogram with Lower Extremity Runoff). This study provides for imaging of the abdomen, pelvis, and both legs. A separate authorization for Lower Extremity CTA is **NOT** needed.

For evaluation of known or suspected lower extremity arterial-vascular disease:

- For known or suspected atherosclerotic peripheral arterial disease (such as claudication, or clinical concern for vascular causes of ulcers) when any **ONE** of the following non-invasive studies (pulse volume recording, ankle-brachial index, toe brachial index, segmental pressures, or doppler ultrasound) are abnormal or indeterminate: **OR**
 - Ankle-brachial index (ABI) (< 0.9 is the cutoff for diagnosis of peripheral arterial disease and >1.4 is considered inconclusive)
 - Toe brachial index (< 0.7 is the cutoff for diagnosis of peripheral arterial disease)
 - Segmental pressure test (a pressure gradient > 20 mmHg is considered abnormal)

- Doppler ultrasound
- Treadmill test
- 6-minute walking test
- For acute critical limb ischemia with any **ONE** of the below clinical signs of peripheral artery vascular disease: ~~(prior ultrasound is not needed; if done and negative, CTA should still be approved)~~
 - Ischemic rest pain
 - Tissue loss
 - Gangrene

NOTE: Prior ultrasound is **NOT** needed ^(1,2)

- After prior stenting or surgery (arterial and/or venous) with any **ONE** of the following:
 - Recurrent symptoms
 - Signs of recurrent disease on examination
 - Abnormal / indeterminate ~~prior~~ non-invasive testing or imaging (Such as, ankle brachial index (ABI), ultrasound)

PRE-OPERATIVE OR /POSTOPERATIVE ASSESSMENT PROCEDURE EVALUATION

When not otherwise specified in the guideline:

Preoperative Evaluation ⁽⁵⁾:

- Evaluation of interventional vascular procedures for luminal patency versus restenosis due to conditions such as atherosclerosis, thromboembolism, and intimal hyperplasia ⁽⁶⁾
- Imaging of the area requested is needed to develop a surgical plan

Post-Operative/Post-Procedural Evaluation ⁽⁵⁾

- Known or suspected complications
- A clinical reasoning is provided how imaging may change management ~~Evaluation of post-operative complications, e.g., pseudoaneurysms related to surgical bypass grafts, vascular stents, and stent-grafts~~ ⁽⁶⁾

NOTE: This section applies only within the first few months following surgery ~~Follow-up study may be needed to help evaluate a patient's progress after treatment, procedure, intervention, or surgery.~~

~~DOCUMENTATION REQUIRES A MEDICAL REASON THAT CLEARLY INDICATES WHY ADDITIONAL IMAGING IS NEEDED FOR THE TYPE AND AREA(S) REQUESTED.~~

FURTHER EVALUATION OF INDETERMINATE FINDINGS ~~ON PRIOR IMAGING~~

Unless follow-up is otherwise specified within the guideline:

- For initial evaluation of an inconclusive finding on a prior imaging report that requires further clarification
- One follow-up exam of a prior indeterminate MR/CT finding to ensure no suspicious interval change has occurred. (No further surveillance unless specified as highly suspicious or change was found on last follow-up exam)

IMAGING IN KNOWN GENETIC SYNDROMES AND RARE DISEASES/CONDITIONS

- Williams Syndrome ⁽⁶⁾:
 - Abnormal vascular exam ~~When there is concern for vascular disease or imaging findings (such as concern for including renal artery stenosis,) based on abnormal exam or imaging findings (such as~~ diminished pulses, bruits or signs of diffuse thoracic aortic stenosis)
- For other syndromes and rare diseases not otherwise addressed in the guideline, coverage is based on a case-by-case basis using societal guidance

Combination Studies for Known Genetic Conditions

NOTE: When medical necessity is met for an individual study AND conscious sedation is required (such as for young pediatric patients or patients with significant developmental delay), the entire combination is indicated)

Abdominal Aorta CTA with Lower Extremity Runoff / Chest CTA

- Williams Syndrome: ~~When there is concern for~~ Abnormal vascular ~~disease-exam or imaging findings (including such as concern for~~ renal artery stenosis,) ~~based on abnormal exam or imaging findings (such as~~ diminished pulses, bruits or signs of diffuse thoracic aortic stenosis ⁽⁶⁾

OTHER COMBINATIONS STUDIES WITH ABDOMINAL AORTA CTA WITH LOWER EXTREMITY RUNOFF

NOTE: When medical necessity is met for an individual study AND conscious sedation is required (such as for young pediatric patients or patients with significant developmental delay), the entire combination is indicated)

Abdominal Aorta CTA ~~Angiography~~ with Lower Extremity Runoff and Chest CTA

- To evaluate for an embolic source of lower extremity thromboembolic vascular disease ~~when other imaging such as echocardiography suggests a cardiac source of the embolism~~

NOTE: Echocardiogram is also indicated as the heart is the most commonly reported source of lower extremity emboli

CODING AND STANDARDS

Coding

CPT Codes

75635

Applicable Lines of Business

| | |
|---|--|
| ☒ | CHIP (Children’s Health Insurance Program) |
| ☒ | Commercial |
| ☒ | Exchange/Marketplace |
| ☒ | Medicaid |
| ☒ | Medicare Advantage |

BACKGROUND

Contraindications and Preferred Studies

- Contraindications and reasons why a CT/CTA cannot be performed may include: impaired renal function, significant allergy to IV contrast, pregnancy (depending on trimester)
- Contraindications and reasons why an MRI/MRA cannot be performed may include: impaired renal function, claustrophobia, non-MRI compatible devices (such as non-compatible defibrillator or pacemaker), metallic fragments in a high-risk location, patient exceeds weight limit/dimensions of MRI machine

~~SUSPECTED PERIPHERAL ARTERIAL DISEASE~~

~~CTA (OR MRA) IS AN EXCELLENT TOOL TO DIAGNOSE LOWER EXTREMITY PERIPHERAL ARTERIAL DISEASE (PAD). BENEFITS INCLUDE THE FAST-SCANNING TIME AND ACCURATE DETECTION OF OCCLUSIONS AND STENOSIS. ACCORDING TO THE SOCIETY FOR VASCULAR SURGERY GUIDELINES, “MEASUREMENT OF THE ANKLE-BRACHIAL INDEX (ABI) IS THE PRIMARY METHOD FOR ESTABLISHING THE DIAGNOSIS OF PAD. AN ABI OF ≤ 0.90 HAS BEEN DEMONSTRATED TO HAVE HIGH SENSITIVITY AND SPECIFICITY FOR THE IDENTIFICATION OF PAD COMPARED WITH THE GOLD STANDARD OF INVASIVE ARTERIOGRAPHY.” THE PRESENCE OF A NORMAL ABI AT REST AND FOLLOWING EXERCISE ALMOST EXCLUDES ATHEROSCLEROTIC DISEASE AS A CAUSE FOR LEG CLAUDICATION.~~

~~WHEN AN ABI IS > 1.40 (SUGGESTING NONCOMPRESSIBLE CALCIFIED VESSELS) AND~~

~~CLINICAL SUSPICION IS HIGH, OTHER TESTS SUCH AS TOE-BRACHIAL INDEX ≤ 8 , A RESTING TOE PRESSURE ≤ 40 MM HG, A SYSTOLIC PEAK POSTERIOR TIBIAL ARTERY FLOW VELOCITY ≤ 10 CM/S MAY BE USED. "IN SYMPTOMATIC PATIENTS IN WHOM REVASCULARIZATION TREATMENT IS BEING CONSIDERED, WE RECOMMEND ANATOMIC IMAGING STUDIES, SUCH AS ARTERIAL DUPLEX ULTRASOUND, CTA, MRA, AND CONTRAST ARTERIOGRAPHY." THIS LATER STATEMENT IS ACCOMPANIED BY A "B" (MODERATE) RATING FOR THE ACCOMPANYING EVIDENCE ("A" = HIGH, "C" = LOW) "IN PATIENTS WITH LIMITED RENAL FUNCTION OR PLANNED SURGICAL INTERVENTION, NONINVASIVE IMAGING TESTS (PARTICULARLY MRA AND CTA) MAY OBTAIN THE NEED FOR DIAGNOSTIC CATHETER ANGIOGRAPHY TO VISUALIZE THE LOCATION AND SEVERITY OF PERIPHERAL VASCULAR DISEASE."~~

~~FOLLOW-UP IMAGING POST VASCULAR SURGERY PROCEDURES HAVE NOT BEEN WELL RESEARCHED WITHOUT CLEAR SURVEILLANCE PROTOCOLS IN PLACE. CLINICAL EXAM, ABI AND EUS WITHIN THE FIRST MONTH OF ENDOVASCULAR THERAPY ARE GENERALLY RECOMMENDED TO ASSESS FOR RESIDUAL STENOSIS, AND AGAIN AT 6 AND 12 MONTHS, THEN ANNUALLY. MORE SOPHISTICATED IMAGING WITH CTA, MRA, OR INVASIVE CATHETER~~

ANGIOGRAPHY IS RESERVED FOR COMPLEX CASES. SUMMARY OF EVIDENCE

ACR Appropriateness Criteria® Lower Extremity Arterial Claudication-Imaging Assessment for Revascularization: 2022 Update ⁽¹⁾

Study Design: This document is an update on the ACR Appropriateness Criteria for imaging assessment in lower extremity arterial claudication, focusing on revascularization. It is based on evidence-based guidelines reviewed annually by a multidisciplinary expert panel.

Target Population: The target population includes patients with arterial claudication, a common manifestation of peripheral artery disease (PAD), which affects 3% to 7% of the general population and 20% of people over 70 years of age.

Key Factors: The document discusses the appropriate use of ultrasound, invasive arteriography, MR angiography, and CT angiography for diagnosing and treating claudication. Claudication prevalence ranges from 1.6% to almost 8%, depending on various factors. Noninvasive hemodynamic tests like the ankle-brachial index are essential for diagnosis. The document emphasizes the importance of imaging in diagnosing individual lesions and triaging patients for possible percutaneous or surgical intervention.

Society for Vascular Surgery practice guidelines for atherosclerotic occlusive disease of the lower extremities: Management of asymptomatic disease and claudication ⁽²⁾

Study Design: This document provides practice guidelines for managing atherosclerotic occlusive disease of the lower extremities, focusing on asymptomatic disease and claudication. The guidelines were developed by the Society for Vascular Surgery Lower Extremity Guidelines Writing Group.

Target Population: The target population includes patients with peripheral arterial disease (PAD), which affects an estimated 8 to 12 million Americans. The prevalence of PAD increases with age, smoking, diabetes, hypertension, and obesity.

Key Factors: The document highlights the association between PAD and risk factors such as age, smoking, diabetes, hypertension, and hypercholesterolemia. The ankle-brachial index (ABI) is recommended as the first-line noninvasive test to diagnose PAD. Routine screening for PAD in asymptomatic individuals is not recommended. Emphasis is placed on risk factor modification, medical therapies, and exercise programs to improve cardiovascular health and functional performance. Revascularization for intermittent claudication (IC) is appropriate for selected patients with disabling symptoms. The guidelines recommend multidisciplinary smoking cessation interventions, statin therapy, optimizing diabetes control, and antiplatelet therapy for patients with IC.

ANALYSIS OF EVIDENCE

Analysis ^(1,2):

In summary, both articles provide valuable insights into the use of CTA for imaging the abdominal aorta and lower extremities. They agree on the high accuracy and utility of CTA, its advantages over other imaging modalities, and its ability to detect nonvascular findings. However, they differ in their focus on technical considerations, clinical guidelines, and radiation exposure. By integrating the findings from both articles, healthcare providers can make more informed decisions about the use of CTA in the management of PAD.

Shared Findings

Both articles emphasize the importance of CTA (Computed Tomography Angiography) in the imaging and assessment of peripheral arterial disease (PAD), particularly for the abdominal aorta and lower extremities. They agree on several key points:

- Both articles highlight the high sensitivity and specificity of CTA for detecting stenoses greater than 50% in diameter. CTA is praised for its ability to provide volumetric images, which allow for extensive post-processing and the creation of arterial road maps.
- CTA is considered superior to catheter angiography for imaging the aorta and iliac arteries. It can readily detect stenosis caused by plaque or thrombus and is more clinically useful than duplex ultrasound (US) for assessing bypass grafts.
- Both articles note that CTA, like MRA (Magnetic Resonance Angiography), is a cross-sectional technique that can show nonvascular findings, such as aneurysms and cystic adventitial disease, which are not detected with catheter arteriography.

POLICY HISTORY

| Date | Summary |
|------------------|---|
| <u>July 2025</u> | <ul style="list-style-type: none"> ● <u>Added a Summary of Evidence and Analysis of Evidence</u> ● <u>Last revised date adjusted to July</u> ● <u>Removed “on prior imaging” under further evaluation to align with guidelines</u> |
| <u>June 2025</u> | <ul style="list-style-type: none"> ● <u>This guideline number changed from 035 to 2006</u> ● <u>Added in general information statement regarding guideline criteria development by reputable sources, standard of care, and best practices</u> ● <u>Updated language in the preoperative and postoperative section</u> ● <u>Segment added to the combinations studies about if the required use of conscious sedation is needed the entire combination is</u> |

| Date | Summary |
|-----------|---|
| | <p>indicated</p> <ul style="list-style-type: none"> ● Reduced background section |
| June 2024 | <ul style="list-style-type: none"> ● Updated references ● Added Combo studies, Genetic Syndromes and Rare Diseases, Contraindications and Preferred Studies sections ● Updated guideline name |
| May 2023 | <ul style="list-style-type: none"> ● General Information moved to beginning of guideline with added statement on clinical indications not addressed in this guideline ● Added statement regarding further evaluation of indeterminate findings on prior imaging |

LEGAL AND COMPLIANCE

Guideline Approval

Committee

Reviewed / Approved by Evolent Specialty [Services](#) Clinical Guideline Review Committee

Disclaimer

Evolent Clinical Guidelines do not constitute medical advice. Treating health care professionals are solely responsible for diagnosis, treatment, and medical advice. Evolent uses Clinical Guidelines in accordance with its contractual obligations to provide utilization management. Coverage for services varies for individual members according to the terms of their health care coverage or government program. Individual members' health care coverage may not utilize some Evolent Clinical Guidelines. [Evolent clinical guidelines contain guidance that requires prior authorization and service limitations.](#) A list of procedure codes, services or drugs may not be all inclusive and does not imply that a service or drug is a covered or non-covered service or drug. Evolent reserves the right to review and update this Clinical Guideline in its sole discretion. Notice of any changes shall be provided as required by applicable provider agreements and laws or regulations. Members should contact their Plan customer service representative for specific coverage information.

[Evolent Clinical Guidelines are comprehensive and inclusive of various procedural applications for each service type. Our guidelines may be used to supplement Medicare criteria when such criteria is not fully established. When Medicare criteria is determined to not be fully established, we only reference the relevant portion of the corresponding Evolent Clinical Guideline that is applicable to the specific service or item requested in order to determine medical necessity.](#)

REFERENCES

1. Azene EM, Steigner ML, Aghayev A, et al. ACR Appropriateness Criteria® Lower Extremity Arterial Claudication-Imaging Assessment for Revascularization: 2022 Update. *Journal of the American College of Radiology*. 2022;19(11):S364-S373. doi:10.1016/j.jacr.2022.09.002
2. Conte MS, Pomposelli FB, Clair DG, et al. Society for Vascular Surgery practice guidelines for atherosclerotic occlusive disease of the lower extremities: Management of asymptomatic disease and claudication. *J Vasc Surg*. 2015;61(3):2S-41S.e1. doi:10.1016/j.jvs.2014.12.009
3. Singh-Bhinder N, Kim DH, Holly BP, et al. ACR Appropriateness Criteria® Nonvariceal Upper Gastrointestinal Bleeding. *Journal of the American College of Radiology*. 2017;14(5):S177-S188. doi:10.1016/j.jacr.2017.02.038
4. Werncke T, Ringe KI, von Falck C, Kruschewski M, Wacker F, Meyer BC. Diagnostic Confidence of Run-Off CT-Angiography as the Primary Diagnostic Imaging Modality in Patients Presenting with Acute or Chronic Peripheral Arterial Disease. Zhang H, ed. *PLoS One*. 2015;10(4):e0119900. doi:10.1371/journal.pone.0119900
5. Browne WF, Sung J, Majdalany BS, et al. ACR Appropriateness Criteria® Sudden Onset of Cold, Painful Leg: 2023 Update. *Journal of the American College of Radiology*. 2023;20(11):S565-S573. doi:10.1016/j.jacr.2023.08.012
6. Morris CA. Williams Syndrome. *GeneReviews*®. Published online April 13, 2023. <https://www.ncbi.nlm.nih.gov/books/NBK1249/>