

National Imaging Associates, Inc.	
Clinical guidelines:	Original Date: August 2016
SHOULDER ARTHROSCOPY	
CPT Codes**:	Last Revised Date: May 2022 May
- Shoulder Rotator Cuff Repair: 23410, 23412, 23420,	<u>2023</u>
29827	
- Shoulder Labral Repair: 23450, 23455, 23460,	
23462, 23465, 23466, 29806, 29807	
- Frozen Shoulder Repair/Adhesive Capsulitis: 29825,	
23700	
- Shoulder Surgery Other: 23120, 23125, 23130,	
23405, 23415, 23430, 23700, 29805, 29819, 29820,	
29821, 29822, 29823, 29824, 29825, +29826, 29828	
**See UM Matrix for allowable billed groupings and	
additional covered codes	
Guideline Number: NIA_CG_318	Implementation Date: January
	20 <u>2423</u>

GENERAL INFORMATION

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

General Requirements for Elective Surgery of the Shoulder

Elective surgery of the shoulder may be considered if the following general criteria are met:

- There is clinical correlation of individual's subjective complaints with objective exam findings and/or imaging (when applicable)
- Individual has limited function (age-appropriate activities of daily living (ADLs), occupational, athletic)
- Individual is medically stable and optimized for surgery with no uncontrolled comorbidities (such as diabetes)

^{*} National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.

- Individual does not have an active local or systemic infection
- Individual does not have active, untreated drug dependency (including but not limited to narcotics, opioids, muscle relaxants) unless engaged in a treatment program
- A smoking cessation program is highly recommended and should be instituted preoperatively for all actively smoking patients^{1, 2}

Clinical notes should address:

- Symptom onset, duration, and severity
- Loss of function and/or limitations
- Type and duration of non-operative management modalities (where applicable)

Non-operative management, when required, will be specified within the clinical indications below and may include one or more of the following:

- Physical therapy or properly instructed home exercise program
- Rest or activity modification
- Application of heat or ice
- Minimum of 4 weeks of oral NSAIDs (if not medically contraindicated)
- Single injection of corticosteroid and local anesthetic into subacromial, <u>or</u>intraarticular space, or bicipital groove

INDICATIONS

DIAGNOSTIC SHOULDER ARTHROSCOPY

Diagnostic arthroscopy is considered medically necessary:

For the evaluation of pain prior to a painful total shoulder arthroplasty³⁻⁶

OR

When All of the following criteria have been met:

- Severe, disabling pain and/or a documented loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
- Individual demonstrates **any** of the following abnormal, shoulder physical examination findings, as compared to the non-involved side:
 - Functionally limited range of motion (active or passive)
 - Measurable loss in strength
 - Positive impingement signs
- Failure of non-surgical management for at least 12 weeks duration to include **TWO** of the following:
 - Rest or activity modifications/limitations

Page 2 of 32 Shoulder Arthroscopy



- o Ice/heat
- Use of a sling/immobilizer/brace
- Pharmacologic treatment: oral/topical NSAIDS, acetaminophen, analgesics, tramadol
- Physical therapy modalities
- Supervised home exercise program
- Individual has undergone an appropriate radiographic work-up that includes <u>both</u> routine x-rays and an MRI evaluation which are determined to be inconclusive for <u>internal derangement/pathology</u> <u>a specific diagnosis.</u>
- Other potential diagnostic conditions have been excluded, including, but not limited to, fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain and arthritis

NOTE: The following is not managed by NIA: Magellan:

• In-office diagnostic arthroscopy (e.g., Mi-Eye, VisionScope)⁷

ROTATOR CUFF REPAIR (RCR)

Surgical treatment of a rotator cuff tear (RCT) should only be performed when there is a clinical correlation of symptoms, clinical exam findings, imaging, and failed non-operative management (where required).⁸⁻¹⁰

NOTE: See section on subscapularis tears

Partial-Thickness Rotator Cuff Tear or Calcific Tendinitis

Surgical repair of **a partially torn rotator cuff** may be necessary when <u>ALL</u> of the following criteria are met:

- Reproducible rotator cuff pain patterns (lateral arm, deltoid pain rarely radiating past the elbow, night pain, or pain with overhead motions)
- Positive impingement signs and/or tests on exam (Hawkins, Neer, Jobe test or reproducible pain when arm is positioned overhead (above plane of shoulder) with relief of pain when arm is repositioned below the plane of the shoulder)¹¹
- Functional loss (age-appropriate activities of daily living (ADL), occupational, athletic)
- MRI or ultrasound^{12, 13}(if an MRI cannot be performed) that demonstrates a partial thickness tear (articular-sided, concealed, or bursal-sided) or evidence of calcific tendinitis
- Failure of at least 12 weeks of non-operative treatment, <u>including at least 6 weeks of</u> <u>physical therapy or a properly instructed home exercise program that includes exercises</u>



for scapular dyskinesis when present **AND** one of the following: including at least two of the following criteria:

o Rest or activity modification

o Minimum of 4 weeks of oral NSAIDs (if not medically contraindicated)

• Physical therapy or properly instructed home exercise program

Rest or activity modification

• Minimum of 4 weeks of oral NSAIDs (if not medically contraindicated) No cortisone injection within 12 weeks prior to surgery.¹⁴⁻²³

NOTE: The following is not managed by Magellan: NIA

• US-guided percutaneous debridement or tenotomy (e.g., Tenex, TenJet)

Small (< 1 cm), Full-Thickness Rotator Cuff Tear

Surgical repair of a **small full-thickness rotator cuff tear** may be necessary when <u>ALL</u> of the following criteria are met:

- Reproducible rotator cuff pain patterns (lateral arm, deltoid pain not radiating past the elbow, night pain, or pain with overhead motions)
- Positive impingement signs and/or tests on exam (Hawkins, Neer, Jobe test or reproducible pain when arm is positioned overhead (above plane of shoulder) with relief of pain when arm is repositioned below the plane of the shoulder)
- Functional loss (age-appropriate activities of daily living (ADLs), occupational, athletic)
- Rotator cuff weakness or severe pain with rotator cuff testing on physical exam
- MRI or Ultrasound^{12, 13} that demonstrates a small, full thickness tear (< 1 cm)
- Failure of at least 6 weeks of non-operative treatment*, including physical therapy or a properly instructed home exercise program (that includes exercises for scapular dyskinesis when present) **AND** one of the following:
 - Rest or activity modification
 - Minimum of 4 weeks of oral NSAIDs (if not medically contraindicated)
- No cortisone injection within 12 weeks prior to surgery.¹⁴⁻²³

***NOTE**: The requirement for conservative, non-operative treatment is waived in individuals less than age 55 with an acute traumatic tear (onset of shoulder pain attributed to a specific traumatic event with no prior history of significant shoulder pain). For ages > 55, non-operative treatment may be waived on a case-by-case basis.

Medium (1-3 cm) or Large (3-5 cm), Full-Thickness Rotator Cuff Tear

Page **4** of **32** Shoulder Arthroscopy



Surgical repair of a **medium or large full-thickness rotator cuff tear** may be necessary when the following criteria are met:

• Significant progression of a full-thickness tear on serial imaging performed at least 12 weeks apart (at least 50% increase in tear size)

OR

- When **ALL** of the following criteria are met:
 - Reproducible rotator cuff pain patterns (lateral arm, deltoid pain rarely not radiating past the elbow, night pain, or pain with overhead motions)
 - Positive impingement signs and/or tests on exam (Hawkins, Neer, Jobe, empty can or drop-arm test or reproducible pain when arm is positioned overhead (above plane of shoulder) with relief of pain when arm is repositioned below the plane of the shoulder
 - o Rotator cuff weakness or severe pain with rotator cuff testing on physical exam
 - Functional loss (age-appropriate activities of daily living (ADLs), occupational or athletic)
 - MRI or ultrasound^{12, 13} results support a medium (1-3 cm) or large (3-5 cm), fullthickness tear (tear must be a complete single tendon or greater)
 - MRI demonstrates no advanced fatty changes (Goutallier stage 0 (normal muscle), 1 (some fatty streaks), or 2 (less than 50% fatty degeneration or infiltration)²⁴⁻²⁶
 - Warner classification of atrophy "none" or "mild"^{27, 28}
 - No cortisone injection within 12 weeks prior to surgery.¹⁴⁻²³

Massive (> 5 cm and ≥ 2 tendons involved), Full-Thickness Rotator Cuff Tear

Surgical repair of **a massive torn rotator cuff** *WITH OR WITHOUT* a superior capsular reconstruction may be necessary when **ALL** of the following criteria are met²⁹⁻³²:

- MRI or ultrasound^{12, 13} demonstrates massive (> 5 cm), full-thickness tears (with intact or reparable subscapularis tendon for superior capsular reconstruction)
- MRI demonstrates no advanced fatty changes (Goutallier stage 0 (normal muscle), 1 (some fatty streaks), or 2 (less than 50% fatty degeneration or infiltration)²⁴⁻²⁶
- Warner classification of atrophy "none" or "mild"^{27, 28}
- No x-ray evidence of chronic subacromial articulation of the humeral head, defined as an acromiohumeral space less than 5 mm (Hamada grade 2)^{31, 33, 34}
- No advanced or severe arthritis (severe narrowing of glenohumeral space or bone-onbone articulation, large osteophytes, subchondral sclerosis, or cysts, etc.)
- No cortisone injection within 12 weeks prior to surgery.¹⁴⁻²³



NOTE: AAOS consensus guidelines state that partial repair and superior capsular reconstruction, can improve patient reported outcomes.²⁹

Subscapularis Tears

Surgical repair of a subscapularis rotator cuff tear may be necessary when the following criteria are met:³⁵⁻⁴²

- History of an acute injury or chronic complaints of anterior shoulder pain, weakness, or functional impairment
- Positive physical examination findings of subscapularis deficiency lift-off, bear-hug, belly press test, etc.
- MRI demonstrates a significant partial thickness tear (at least 50% of tendon), fullthickness tear, or any tear associated with subluxation of the biceps tendon
- No cortisone injection within 12 weeks prior to surgery.¹⁴⁻²³

AN ISOLATED SUPERIOR CAPSULAR RECONSTRUCTION MAY BE NECESSARY WHEN ALL OF THE FOLLOWING CRITERIA ARE MET⁴³⁻⁵⁰:

- MRI or ultrasound^{12, 13} demonstrates massive (> 5 cm), full-thickness tears with an intact or reparable subscapularis tendon
- No x-ray evidence of chronic subacromial articulation of the humeral head, defined as an acromiohumeral space less than 5 mm (Hamada grade 2)^{31, 33, 34}
- No advanced or severe arthritis (severe narrowing of glenohumeral space or bone-onbone articulation, large osteophytes, subchondral sclerosis, or cysts, etc.)

NOTE: A CONCOMITANT ROTATOR CUFF REPAIR IS NOT ALLOWABLE WITH ADVANCED GOUTALLIER OR WARNER MUSCLE ATROPHY CHANGES AS NOTED IN THE PREVIOUS SECTION.

Rotator Cuff Repair (RCR) Revision

Surgical revision within 1 year of a previously repaired small, medium, large or massive torn rotator cuff will be reviewed on a case-by-case basis and must include an MRI (with or without arthrogram) or CT arthrogram that demonstrate failure of healing (Sugaya type 4-5, see background section) or recurrent tear > 12 weeks after index surgery.⁵¹

All RCR revision cases greater than 1 year following an initial repair must again meet indications as specified by tear size listed in Background section.

Contraindications (applies to all rotator cuff repair):

- Active infection (local or remote)
- Treatment of asymptomatic, full thickness rotator cuff tear

Page 6 of 32 Shoulder Arthroscopy



- Active systemic bacteremia
- Deltoid or rotator cuff paralysis
- Advanced or severe arthritis (severe narrowing of glenohumeral space or bone-on-bone articulation, large osteophytes, subchondral sclerosis, or cysts, etc.)
- No cortisone injection within 12 weeks prior to surgery.¹⁴⁻²³

LABRAL REPAIRS

Repair of Superior Labral Anterior-Posterior (SLAP) Tear

Surgical indications should be focused on clinical symptoms and failure to respond to nonoperative treatments, rather than imaging (due to a higher percentage of tears being missed on images **AND** significant over-diagnosing of tears based on imaging-alone).

Repair (not debridement of a SLAP lesion) may be necessary when **ALL** of the following criteria are met:

- History compatible with tear (acute onset in thrower or overhead athlete, fall, traction injury, shear injury (MVA), lifting injury
- Pain localized to the glenohumeral joint (often only associated with certain reaching or lifting activities and at night) or painful catching/popping/locking sensations
- Inability to perform desired tasks without pain (age-appropriate ADLs, sports, occupation)
- Physical examination demonstrates findings of a SLAP tear (active compression test (O'Brien test), compression rotation test, clunk, or crank test, etc.)
- Age < 40; requests for SLAP repair in an individual age > 40 will be reviewed on a caseby-case basis⁵²
- MRI demonstrating superior labral tear
- Type 2 or 4 SLAP tear (not type 1 or 3)
 - I Labral and biceps fraying, anchor intact
 - o II Labral fraying with detached biceps tendon anchor
 - III Bucket handle tear, intact biceps tendon anchor (biceps separates from bucket handle tear)
 - IV Bucket handle tear with detached biceps tendon anchor (remains attached to bucket handle tear)
- Failure of at least 12 weeks of non-operative treatment, including activity modification/avoidance of painful activities **AND** <u>at least-**ONE**</u> of the following:
 - o Minimum of 4 weeks of oral NSAIDs (if not medically contraindicated)
 - Physical therapy or a properly instructed home exercise program

Contraindications:

• ANY evidence of degenerative disease upon imaging

Page **7** of **32** Shoulder Arthroscopy



- Smoker and age > 40
- Diabetics with poor control HgBA1c > 7
- MRI findings not attributable to normal common variants (for example, labral overhang)

NOTE: In cases where a true SLAP tear exists, but the individual has one or more contraindications or findings at the time of surgery that indicates a repair is not feasible, a SLAP debridement (limited, extensive debridement), biceps tenotomy or tenodesis may be an alternative.⁵²⁻⁵⁴ Even with repairable type II SLAP tears, biceps tenodesis is a viable alternative to repair.⁵⁵⁻⁵⁷ See Tenotomy and Tenodesis Indications.

Anterior-Inferior Labral Tear (Bankart lesion)58

Bankart repair of **an acute labral tear** may be necessary when **ALL** of the following criteria are met:

- History of an acute event of instability (subluxation or dislocation) or acute onset of pain following activity
- Acute labral tear on MRI or CT imaging
- Age < 30
- Range of motion is not limited by stiffness upon physical exam (not required if there has been a recent episode of instability)
- Clinical exam findings demonstrate positive apprehension test, positive relocation test, positive labral grind test, or objective laxity with pain.

Bankart repair for **recurrent (two or more episodes of subluxation or dislocation) associated with a labral tear** may be necessary when **ALL** of the following criteria are met⁵⁹:

- Recurrent instability (subluxation or dislocation)
- <u>MRI e</u>Evidence of a labral tear with or without bony Bankart fracture of the glenoid upon imaging
- Range of motion is not limited by stiffness upon physical exam (not required if there has been a recent episode of instability)
- Clinical exam findings demonstrate positive apprehension test, positive relocation test, positive labral grind test, or objective laxity with pain.

Contraindications:

Pain only (no documented recurrent instability events) in individuals over 40
X-ray, MRI, or CT documentation of significant degenerative arthritis of the glenohumeral joint

 Radiographic findings of an engaging Hill Sachs humeral head defect or glenoid bone loss (if surgery only includes Bankart repair).; (see below indications for Latarjet and/or remplissage procedure) Latarjet or Remplissage procedures should be considered for anterior dislocations of the shoulder when there is evidence of an engaging ("off-



track")* Hill-Sachs lesion of the humerus, or if there is greater than 20% glenoid bone loss by x-ray, CT, or MRI⁶⁰⁻⁷⁹

- Pain only (no documented recurrent instability events) in individuals over 40
- X-ray, MRI, or CT documentation of significant degenerative arthritis of the glenohumeral joint

* The glenoid track, a zone of dynamic contact during arm elevation, is a unique biomechanical model that uses both glenoid and humeral head bone loss to predict subsequent risk of humeral head engagement and possible dislocation. An *engaging* Hill-Sachs bony defect, or "off-track" lesion, is one in which the width of the bony defect is greater than the width of the glenoid track. Off-track engagement occurs when the medial margin of the Hill-Sachs defect engages the glenoid track. If there is bony loss of the glenoid as well, the glenoid track will proportionately be less, causing greater risk of engagement. A *nonengaging*, or "on-track" Hill-Sachs lesion is one in which the width of the bony defect is less than the width of the glenoid track. Using preoperative CT or MR imaging, the glenoid track can identify individuals who are more likely to fail only a primary capsuloligamentous Bankart repair. Glenoid track evaluation shows that restoring the track (glenoid) to its normal width should be the first priority in restoring shoulder stability.⁸⁰⁻⁸⁷

Posterior Labral Tear

Surgical repair of a posterior labral tear may be necessary when **ALL** of the following criteria are met:

- Symptoms of pain **OR** painful catching/popping OR instability
- MRI findings of posterior labral tear
- Exam findings demonstrate positive load-shift test, jerk test, glenohumeral grind test, or objective laxity with pain or profound weakness⁸⁸
- Failure of at least 12 weeks of non-operative treatment (unless presenting as a traumatic tear in a competitive athlete at any level) that includes any **two** of the following:
 - Physical therapy or a properly instructed home exercise program
 - Rest or activity modification
 - Minimum of 4 weeks of oral NSAIDs (if not medically contraindicated)
- Age < 40
- No radiographic evidence of degenerative disease (e.g., posterior glenoid cartilage loss, subchondral glenoid cysts, mucoid degeneration of labrum, narrowing of joint space with posterior humeral head subluxation on axillary x-ray or axial MRI images)

<u>Combined Labral Tears (e.g., Anterior / Posterior, SLAP / Anterior, SLAP / Posterior, SLAP /</u> <u>Ant. / Post.)⁸⁹</u>



Surgical repair of an **acute combination tear** may be necessary when **ALL** of the following criteria are met:

- History of an acute event of instability (subluxation or dislocation)
- Acute labral tear on MRI/CT imaging with/without bony Bankart fracture not > 25% of glenoid width upon imaging
- Age < 30
- Range of motion not limited by stiffness upon physical exam
- Clinical exam findings demonstrate positive apprehension test and positive relocation test, **OR** positive labral grind test **OR** objective laxity with pain
- Minimal to no evidence of degenerative changes on imaging

•

Surgical repair of **recurrent combination tear** may be necessary when **ALL** of the following criteria are met:

- Recurrent instability (subluxation or dislocation) with at least 2 instability events
- Labral tear on MRI or CT, with/without bony Bankart fracture not > 25% of glenoid width upon imaging
- Range of motion not limited by stiffness upon physical exam
- Clinical exam findings demonstrate positive apprehension test and positive relocation test, or positive labral grind test, or objective laxity with pain
- Minimal to no evidence of degenerative changes on imaging

Open or Arthroscopic Capsulorrhaphy for Multidirectional Instability of the Shoulder (MDI)

Surgical repair for MDI may be necessary when **ALL** of the following criteria are met:

- Individual has pain and limited function (age-appropriate ADLs, occupation, or sports)
- Individual has recurrent instability due to hyperlaxity or mobility and no traumatic dislocation
- Physical exam supports repeatable increased glenohumeral joint translation (greater than 1 cm of movement during the sulcus test)
- Imaging (x-ray and MRI) rules out fracture and/or other soft-tissue injury
- Failure of at least 6 months of formal physical therapy and activity modification

Adhesive Capsulitis (Lysis of Adhesions, Capsulotomy/Capsular Release or Manipulation under Anesthesia)

Surgery for adhesive capsulitis may be necessary when **ALL** of the following criteria are met⁹⁰⁻⁹⁴:

• Individual has pain, loss of motion, and limited function (age-appropriate ADLs, occupation, or sports)





- Physical exam demonstrates loss of motion of at least 50% in 2 planes, <u>as compared to</u> <u>the contralateral shoulder. {use contralateral shoulder for comparison}</u>
- A chest X-ray has been obtained in the past 12 months and co-morbidities (such as diabetes, thyroid disease, lung disease, etc.), and other causes of loss of shoulder motion have been ruled out
- Failure of at least 12 weeks of non-operative treatment that includes physical therapy or a properly instructed home exercise program and documentation of **any one** of the following:
 - Minimum of 4 weeks of oral or topical NSAIDs (if not medically contraindicated)
 - Rest or activity modification
 - Heat/Ice
 - Corticosteroid injection

Distal Clavicle Excision (DCE)

Distal Clavicle Excision may be necessary when ALL of the following criteria are met⁹⁵:

- Positive clinical exam findings as evidenced by pain upon palpation over AC joint and/or pain with cross-body adduction test
- Positive findings on x-Ray **OR** MRI⁹⁶:
 - Radiographic (x-ray) demonstrates narrowed joint space, distal clavicle or medial acromial sclerosis, and/or osteophytes or cystic degeneration of distal clavicle or medial acromion correlating with the clinical findings, patient symptoms and diagnosis; OR MRI findings with edema in the distal clavicle and/or inflammatory change within the joint space correlating with the clinical findings, patient symptoms and diagnosis
- Failure of at least 12 weeks of non-operative treatment that includes **at least two** of the following:
 - Minimum of 4 weeks of oral or topical NSAIDs (if not medically contraindicated)
 - Rest or activity modification
 - AC joint corticosteroid injection (if DCE is to be performed as a standalone procedure, AC injection must be performed*)
 - o Physical therapy or a properly instructed home exercise program

***NOTE:** If DCE is to be performed *in isolation* of other shoulder procedures, an AC joint injection is required for diagnostic purposes and documentation should support pain relief from injection. If no response to injection, this is a strong negative predictor to surgical outcome for isolated DCE.



Long Head Biceps (LHB) Tenotomy/Tenodesis

The indications and outcomes for tenodesis and tenotomy are the same^{35, 97, 98} with the exception that tenodesis is typically better for more active, muscular individuals that are performing higher-demand activities for work or sport. Tenotomy is often preferred in individuals that smoke (this is a relative indication of tenotomy over tenodesis) due to healing problems in tenodesis.

Tenotomy or tenodesis may be necessary when the following criteria are met^{35, 52, 99, 100}:

• Any of the following:

- o When performed in conjunction with a total shoulder arthroplasty
- \circ $\;$ When performed in conjunction with a subscapularis tendon repair
- Age > 50 with SLAP tear
- Smoker with SLAP labral tear (regardless of age, more significant with increasing age)
- Failed SLAP repair
- \circ $\,$ SLAP tear in diabetic or individual with loss of motion or predisposition to stiff shoulder $\,$
- LHB hypertrophy/tearing/subluxation in association with RCR

OR

• Diagnosis of chronic LHB groove pain from tenosynovitis*

AND

- Failure of at least 12 weeks of non-operative treatment to include **TWO** of the following:
 - Minimum of 4 weeks of oral or topical NSAIDs (if not medically contraindicated)
 - Rest or activity modification
 - Bicipital groove corticosteroid injection
 - Physical therapy or a properly instructed home exercise program

NOTE: The following is not managed by NIAMagellan:

• US-guided percutaneous debridement or tenotomy (e.g., Tenex, TenJet)

Loose Body Removal

Loose body removal may be medically necessary when the following criteria are met:

- Documentation of pain, mechanical symptoms (catching or locking), stiffness, loss of motion, feelings of instability or loss of function
- X-ray, CT, or MRI documentation of a loose body

Page **12** of **32** Shoulder Arthroscopy



Synovectomy

Synovectomy as an isolated procedure is usually reserved for primary synovial disease or in cases where secondary hypertrophic synovitis is documented during arthroscopy (these include adhesive capsulitis, osteoarthritis, chronic rotator cuff tear). These should be evident on arthroscopic photographs taken at surgery but may be missed on preoperative images.¹⁰¹

Subacromial Decompression (SAD)

Subacromial decompression may be necessary **in conjunction with** other shoulder procedures (listed below) if there is radiographic (x-ray) evidence of mechanical outlet impingement as evidenced by a Bigliani type 3 morphology. Subacromial decompression should not be performed in isolation.^{102, 103}

- Rotator cuff repair
- Labral repair
- Capsulorrhaphy
- Loose body removal
- Synovectomy
- Debridement
- Distal clavicle excision
- Lysis of adhesions
- Biceps tenodesis/tenotomy

Contraindications:

- Type 1 or Type 2 or a thinned acromion. Subacromial bursectomy may be a reasonable option.
- If individual has received an injection in the subacromial space and there is failure to adequately respond—significant relief (> 50%) for minimum of 1 week—to injection in the subacromial space (pain should respond temporarily if impingement)
- Prior subacromial decompression with either a Type 1 or a thinned acromion or no evidence of overhang on x-ray (unnecessary revision can thin the acromion and lead to deltoid avulsion and/or acromial fracture)
- Open SAD procedures should rarely be performed given the increased morbidity due to deltoid disruption.

BACKGROUND:

This guideline addresses the following elective, non-emergent, arthroscopic shoulder repair procedures:

Page **13** of **32** Shoulder Arthroscopy



- Rotator Cuff Repair
- Labral Repairs
- Lysis of Adhesions (Capsulotomy)
- Distal Clavicle Excision (DCE)
- Long Head Biceps (LHB) Tenotomy or Tenodesis
- Loose body removal
- Synovectomy
- Subacromial Decompression (SAD)

Arthroscopy introduces a fiber-optic camera into the shoulder joint through a small incision for diagnostic visualization purposes. Other instruments may then be introduced to remove, repair, or reconstruct joint pathology.

Surgical indications are based on relevant subjective clinical symptoms, objective physical exam & radiologic findings, and response to previous non-operative treatments when medically appropriate.

Open, non-arthroplasty shoulder repair surgeries are performed as dictated by the type and severity of injury and/or disease.

Rotator Cuff Repair (RCR)

Traditional open rotator cuff repair (RCR) with deltoid take-down should be rare given increased morbidity when compared to arthroscopic or mini-open surgery.

Goutallier Classification of Fatty Infiltration of Rotator Cuff Musculature^{26, 104-107}

Grade 0 – Normal Grade 1 – Mild - muscle contains some fatty streaks Grade 2 – Moderate – more muscle than fat Grade 3 – Severe – equal amounts of fat and muscle Grade 4 – More fat than muscle

Hamada Classification of Rotator Cuff Arthropathy^{33, 34, 108}

Acromiohumeral Interval (AHI)

Grade 1 – AHI over 6 mm Grade 2 – AHI < 5mm Grade 3 – Acetabulization Grade 4 – Acetabulization and narrowed GH joint Grade 5 - Acetabulization with humeral head collapse

Revision Rotator Cuff Repair

Page **14** of **32** Shoulder Arthroscopy



The Sugaya classification for evaluation in revision rotator cuff repair is as follows:

Sugaya Classification

Type I - Sufficient thickness, homogeneous tendon (low signal on T2 images) Type II - Sufficient thickness, partial high-intensity from within the tendon Type III - Insufficient thickness without discontinuity Type IV - Minor discontinuity on more than one slice, suggesting a small tear Type V - Major discontinuity suggesting a moderate or large tear

Labral Repairs

There is a tendency to misinterpret normal degenerative labral changes and variations as "tears" which may lead to over-utilization of surgery if decisions are made upon imaging reports alone. In addition, the anterior-superior labrum (from 12 to 3 o'clock for a right shoulder) has many normal variations that can be misinterpreted as a tear, including sublabral hole/foramen, Buford complex, and a labral overhang with an intact biceps anchor. In general, true labral tears lead to pain, catching, popping, functional limitations (including age-appropriate activities of daily living (ADLs), occupational and athletic), micro-instability, and gross instability. Labral repairs are most-frequently associated with a specific traumatic event.

Anterior-Inferior Labral-Tear (Bankart lesion)

A Bankart tear of the glenoid labrum is located at the 3-6 o'clock position of a right shoulder. It is typically caused by a traumatic instability event (dislocation or subluxation). It can involve the labrum, the capsular ligaments (IGHL [inferior glenohumeral ligamentous complex]) and/or the bone (bony Bankart fracture). If symptomatic, Bankart lesions typically require surgical repair as individuals less than 30 have a high recurrence rate of instability. If there has been significant bone loss of the anterior glenoid, further stabilization might be required by transferring the coracoid process and attached conjoined tendon (Latarjet Procedure) or using a bone graft to the anterior glenoid. Engaging or "off-track" defects of the humeral head (Hill-Sachs lesion) may require the use of portions of the rotator cuff (Remplissage Procedure) to fill the bony defect, in order to further stabilize the shoulder.

Posterior Labral Tear

Similar to Bankart tears, posterior labral tears are often associated with a paralabral cyst that grows large enough to compress the suprascapular nerve (isolated to infraspinatus). Posterior labral tears are frequently associated with contact sports or a history of a traumatic fall/posterior loading of the joint. They are often observed in athletes performing repetitive posterior loading of the joint (offensive linemen in football, weightlifting, push-ups, and bench press). These tears are more likely to result in pain and weakness rather than recurrent dislocations/instability. Posterior labral changes are often misinterpreted on MRI as a "tear" in age > 40 years old, when changes due to early glenohumeral degeneration begin to appear.

Page **15** of **32** Shoulder Arthroscopy



Combined Labral Tears (e.g., Anterior / Posterior, SLAP / Anterior, SLAP / Posterior, SLAP / Ant. / Post.)

Combined tears that require repair are almost always associated with significant recurrent instability. Often tears begin within one area and overtime the failure to repair the original injury causes the tear to extend.

Adhesive Capsulitis (Lysis of Adhesions; Capsulotomy/Capsular Release)

Adhesive capsulitis is a thickening and tightening of the soft tissue capsule that surrounds the glenohumeral joint. Adhesive capsulitis usually begins with the gradual onset of pain and limitation of shoulder motion, with a progression to interference of activities of daily living. Primary adhesive capsulitis is the subject of much debate as the specific causes of this condition are not fully understood. Individuals with uncontrolled diabetes have a significantly higher risk of developing adhesive capsulitis than the general population. Secondary (acquired) adhesive capsulitis develops from a known cause, such as stiffness following a shoulder injury, surgery, or a prolonged period of immobilization. Adhesive capsulitis may last from one to three years, despite active treatment, and is more common in women.

Distal Clavicle Excision (DCE)

The AC joint (acromioclavicular joint) can develop degenerative disease in those over 30 years of age, those with a history of a prior grade I or II AC sprain/separation, those with a history of heavy lifting (labor occupation or strength training), or those with evidence of remote trauma. It can occur in isolated form in younger individuals (distal clavicle osteolysis) but is more commonly observed concomitantly with rotator cuff disease in those over age 40 years of age.

Long Head Biceps (LHB) Tenotomy/Tenodesis

Pain in the area of the long head of the bicep tendon is common, especially in overhead sports and in the presence of rotator cuff tears (especially subscapularis). It can be an isolated source of pain in chronic tenosynovitis, SLAP tears, or small tears of the biceps sling, resulting in dynamic or static subluxation or dislocation of the tendon. LHB problems are frequently missed on MRI (especially using contrast which can mask the pathology). The choice of tenodesis versus tenotomy is controversial. Typically, tenodesis is better for more active, muscular individuals performing higher demand activity (labor, sports). Tenotomy is generally a better option for older, less active individuals with poor muscle definition, as it generally leaves the individual with a "Popeye" deformity and the possibility of biceps cramping or anterior shoulder pain with activity. The choice of tenotomy vs. tenodesis is generally left up to the surgeon/patient.

Loose Body Removal

Page **16** of **32** Shoulder Arthroscopy



Although not as common as in the knee, a loose body in the shoulder may require arthroscopic removal if symptoms such as pain, catching or locking are present. Because of the non-weightbearing status of the shoulder and the axillary fold where a loose body might be positioned, not every loose body diagnosed by imaging requires removal.

Synovectomy

Synovitis is common in many shoulder conditions and typically resolves when the primary pathology is treated. Most commonly, this includes loose bodies, inflammatory arthritis or degenerative arthritis, labral tears, and adhesive capsulitis. Primary synovial diseases include pigmented villonodular synovitis, synovial chondromatosis, rheumatoid arthritis, other inflammatory arthritides, traumatic synovial hypertrophy or metaplasia.

Subacromial Decompression (SAD)

There are 3 types of acromion anatomy according to Bigliani classification: type 1, flat (20%), type 2, curved (40%) and type 3, hooked, (40%). Acromioplasty involves removing bone from the undersurface of the acromion to change a type 3 (hooked) acromion to a type 1 (flat) acromion. Although debated for decades, current evidence concludes that there is no role for isolated acromioplasty (subacromial decompression), which prompted conversion of CPT code 29826 (acromioplasty, subacromial decompression) from an index, primary, "stand-alone" code to an "add-on" code only.¹⁰⁹

Date	Summary
2023	Added the requirement of 6 weeks of physical therapy for partial
	rotator cuff repairs
	 Added the requirement for no significant muscle atrophy or fatty
	infiltration for medium or large rotator cuff repairs
	 <u>Clarification of the indications for Latarjet or Remplissage</u>
	procedures
	 Added requirement for 50% decreased ROM in 2 planes, as
	compared to the opposite shoulder, for frozen shoulder surgery
	 Added requirement for a chest X ray in the past 12 months for
	frozen shoulder surgery
May 2022	Updated background and references
	Further defined the glenoid track verbiage for "off-track" and "on-
	track" Hill Sachs lesions (bony defects of the humeral head)
	 Clarified individual is medically stable and optimized for surgery
	Revised Partial-Thickness Rotator Cuff Tear or Calcific Tendinitis to
	<u>"include two of the following criteria"</u>
	Revised criteria for Latarjet or Remplissage to "Recurrent anterior
	dislocations"

POLICY HISTORY SUMMARIES

Page **17** of **32** Shoulder Arthroscopy



	Non-operative treatment for small RCT revised to ONE of the
	following (previously "at least one")
	 Revised 3 months to 12 weeks throughout
	 Replace "patient" with "individual" where appropriate
	Added:
	Evaluation of pain prior total shoulder arthroplasty as indication for
	a diagnostic arthroscopy
	Cortisone injection within 12 weeks of a rotator cuff repair or
	revision as a contraindication
	 Added more specific indications for repair of a subscapularis
	rotator cuff tear
	 Physical examination findings requirement for SLAP tears
	Criteria for loose body removal
	 "performed in conjunction with a subscapularis tendon repair" to
	criteria for Long Head Biceps Tenotomy/Tenodesis
	Deleted:
	 Requirement for a cortisone injection for calcific tendinopathy
	 Deleted cortisone injections from lists of treatment modalities
	 IA joint corticosteroid injection from non-operative treatments for
	LHB Tenotomy/Tenodesis
	 Rotator cuff repair surgical management statement
June 2021	Added heat and ice as acceptable non-operative treatment
	modalities
	 Added "subluxation" to instability language for Bankart and
	remplissage and Latarjet surgery
	Deleted X-ray/MRI language as procedures to r/o co-morbidities
	for frozen shoulder surgery
	Deleted the requirement for ROM testing for stiffness if there has
	been a recent episode of instability
January 2021	Revised section for massive rotator cuff tears
	 Added section for Superior Capsular Reconstruction, including
	references
	Added classification tables for Goutallier and Hamada grading
November 2020	 Added CPT code 29825 to Shoulder Surgery Other
October 2020	
	 Added statement pertaining to in-office arthroscopy: these
	procedures are considered investigational and are not managed by
	Magellan.



	 Added statement pertaining to US-guided percutaneous
	tenotomy/debridement: these procedures are considered to be
	investigational and are not managed by Magellan.
	Deleted requirement for non-operative treatment for massive
	rotator cuff tears
	Added indications for Latarjet and Remplissage procedures for
	shoulder instability.
	Added indication for biceps tenotomy/tenodesis when performed
	in conjunction with a TSA.
	Updated references
July 2020	Removed CPT code S2300
October 2019	Updated references
	Added US (ultrasound) as acceptable diagnostic test for rotator
	cuff tears
	Addition Superior Capsular Reconstruction to rotator cuff repair
	section
	Added manipulation under anesthesia (MUA) to adhesive capsulitis
	section
November 2018	Added new section: 'Diagnostic Shoulder Arthroscopy'
	Added and update references

REFERENCES:

1. Bishop JY, Santiago-Torres JE, Rimmke N, Flanigan DC. Smoking Predisposes to Rotator Cuff Pathology and Shoulder Dysfunction: A Systematic Review. *Arthroscopy*. Aug 2015;31(8):1598-605. doi:10.1016/j.arthro.2015.01.026

2. Santiago-Torres J, Flanigan DC, Butler RB, Bishop JY. The effect of smoking on rotator cuff and glenoid labrum surgery: a systematic review. *Am J Sports Med*. Mar 2015;43(3):745-51. doi:10.1177/0363546514533776

3. Akgün D, Maziak N, Plachel F, et al. Diagnostic Arthroscopy for Detection of Periprosthetic Infection in Painful Shoulder Arthroplasty. *Arthroscopy*. Sep 2019;35(9):2571-2577. doi:10.1016/j.arthro.2019.03.058

4. Guild T, Kuhn G, Rivers M, Cheski R, Trenhaile S, Izquierdo R. The Role of Arthroscopy in Painful Shoulder Arthroplasty: Is Revision Always Necessary? *Arthroscopy*. Jun 2020;36(6):1508-1514. doi:10.1016/j.arthro.2020.01.045

 5. Khan M, Ayeni OR. Editorial Commentary: Does the Scope Have a Role in Painful Shoulder Arthroplasty? *Arthroscopy*. Jun 2020;36(6):1515-1516. doi:10.1016/j.arthro.2020.02.031
 6. Parker DB, Smith AC, Fleckenstein CM, Hasan SS. Arthroscopic Evaluation and Treatment of Complications That Arise Following Prosthetic Shoulder Arthroplasty. *JBJS Rev*. Aug 2020;8(8):e2000020-8. doi:10.2106/jbjs.Rvw.20.00020

7. Zhang K, Crum RJ, Samuelsson K, Cadet E, Ayeni OR, de Sa D. In-Office Needle Arthroscopy: A Systematic Review of Indications and Clinical Utility. *Arthroscopy*. Sep 2019;35(9):2709-2721. doi:10.1016/j.arthro.2019.03.045

8. Harris JD, Pedroza A, Jones GL. Predictors of pain and function in patients with symptomatic, atraumatic full-thickness rotator cuff tears: a time-zero analysis of a prospective patient cohort enrolled in a structured physical therapy program. *Am J Sports Med*. Feb 2012;40(2):359-66. doi:10.1177/0363546511426003

9. Okoroha KR, Mehran N, Duncan J, et al. Characterization of Rotator Cuff Tears: Ultrasound Versus Magnetic Resonance Imaging. *Orthopedics*. Jan 1 2017;40(1):e124-e130. doi:10.3928/01477447-20161013-04

10. Barreto RPG, Braman JP, Ludewig PM, Ribeiro LP, Camargo PR. Bilateral magnetic resonance imaging findings in individuals with unilateral shoulder pain. *J Shoulder Elbow Surg*. Sep 2019;28(9):1699-1706. doi:10.1016/j.jse.2019.04.001

11. Hegedus EJ, Goode AP, Cook CE, et al. Which physical examination tests provide clinicians with the most value when examining the shoulder? Update of a systematic review with metaanalysis of individual tests. *Br J Sports Med*. Nov 2012;46(14):964-78. doi:10.1136/bjsports-2012-091066

12. Apostolopoulos AP, Angelis S, Yallapragada RK, et al. The Sensitivity of Magnetic Resonance Imaging and Ultrasonography in Detecting Rotator Cuff Tears. *Cureus*. May 1 2019;11(5):e4581. doi:10.7759/cureus.4581

13. Kruse KK, Dilisio MF, Wang WL, Schmidt CC. Do we really need to order magnetic resonance imaging? Shoulder surgeon ultrasound practice patterns and beliefs. *JSES Open Access*. Jul 2019;3(2):93-98. doi:10.1016/j.jses.2019.01.004

14. Agarwalla A, Puzzitiello RN, Mascarenhas R, Sumner S, Romeo AA, Forsythe B. Preoperative Injections May Be an latrogenic Cause of Reoperation After Arthroscopic Rotator Cuff Repair. *Arthroscopy*. Feb 2019;35(2):325-331. doi:10.1016/j.arthro.2018.08.042

15. Cimino AM, Veazey GC, McMurtrie JT, et al. Corticosteroid Injections May Increase Retear and Revision Rates of Rotator Cuff Repair: A Systematic Review. *Arthroscopy*. Aug 2020;36(8):2334-2341. doi:10.1016/j.arthro.2020.04.044

16. Lubowitz JH, Brand JC, Rossi MJ. Preoperative Shoulder Corticosteroid Injection Is Associated With Revision After Primary Rotator Cuff Repair. *Arthroscopy*. Mar 2019;35(3):693-694. doi:10.1016/j.arthro.2018.12.025

17. von Knoch M, Baums MH, Lehmann W, Frosch S. Risk Analysis of Perioperative Injections in Arthroscopic Reconstruction of the Rotator Cuff of the Shoulder - A Systematic Review. *Z Orthop Unfall*. Mar 29 2021;Risikoanalyse perioperativer Infiltrationen bei arthroskopischer Rotatorenmanschettennaht der Schulter – ein systematischer Review. doi:10.1055/a-1394-6469

18. Weber AE, Trasolini NA, Mayer EN, et al. Injections Prior to Rotator Cuff Repair Are Associated With Increased Rotator Cuff Revision Rates. *Arthroscopy*. Mar 2019;35(3):717-724. doi:10.1016/j.arthro.2018.10.116

19. Forsythe B, Agarwalla A, Puzzitiello RN, Sumner S, Romeo AA, Mascarenhas R. The Timing of Injections Prior to Arthroscopic Rotator Cuff Repair Impacts the Risk of Surgical Site Infection. *J Bone Joint Surg Am*. Apr 17 2019;101(8):682-687. doi:10.2106/jbjs.18.00631

20. Saffarini M. Editorial Commentary: "Doctor, Are You Sure the Steroid Injection Won't Harm My Shoulder?" Perhaps We Should Stop Injecting Corticosteroids and Just Repair Those Rotator Cuffs. *Arthroscopy*. Jan 2019;35(1):51-53. doi:10.1016/j.arthro.2018.10.009

21. Traven SA, Brinton D, Simpson KN, et al. Preoperative Shoulder Injections Are Associated With Increased Risk of Revision Rotator Cuff Repair. *Arthroscopy*. Mar 2019;35(3):706-713. doi:10.1016/j.arthro.2018.10.107

22. Woolf SK. Editorial Commentary: Steroid Injections Prior to Arthroscopic Rotator Cuff Repair-Is It Time to Rethink a Conservative Treatment Paradigm? *Arthroscopy*. Mar 2019;35(3):714-716. doi:10.1016/j.arthro.2018.12.017

23. Werner BC, Cancienne JM, Burrus MT, Griffin JW, Gwathmey FW, Brockmeier SF. The timing of elective shoulder surgery after shoulder injection affects postoperative infection risk in Medicare patients. *J Shoulder Elbow Surg*. Mar 2016;25(3):390-7. doi:10.1016/j.jse.2015.08.039

24. Chung SW, Kim JY, Kim MH, Kim SH, Oh JH. Arthroscopic repair of massive rotator cuff tears: outcome and analysis of factors associated with healing failure or poor postoperative function. *Am J Sports Med.* Jul 2013;41(7):1674-83. doi:10.1177/0363546513485719

25. Khair MM, Lehman J, Tsouris N, Gulotta LV. A Systematic Review of Preoperative Fatty Infiltration and Rotator Cuff Outcomes. *Hss j*. Jul 2016;12(2):170-6. doi:10.1007/s11420-015-9465-5

26. Somerson JS, Hsu JE, Gorbaty JD, Gee AO. Classifications in Brief: Goutallier Classification of Fatty Infiltration of the Rotator Cuff Musculature. *Clin Orthop Relat Res*. May 2016;474(5):1328-32. doi:10.1007/s11999-015-4630-1

27. Naimark M, Trinh T, Robbins C, et al. Effect of Muscle Quality on Operative and Nonoperative Treatment of Rotator Cuff Tears. *Orthop J Sports Med*. Aug

2019;7(8):2325967119863010. doi:10.1177/2325967119863010

28. Warner JJ, Higgins L, Parsons IMt, Dowdy P. Diagnosis and treatment of anterosuperior rotator cuff tears. *J Shoulder Elbow Surg*. Jan-Feb 2001;10(1):37-46. doi:10.1067/mse.2001.112022

29. Weber S, Chahal J. Management of Rotator Cuff Injuries. *J Am Acad Orthop Surg*. Mar 1 2020;28(5):e193-e201. doi:10.5435/jaaos-d-19-00463

30. Shon MS, Koh KH, Lim TK, Kim WJ, Kim KC, Yoo JC. Arthroscopic Partial Repair of Irreparable Rotator Cuff Tears: Preoperative Factors Associated With Outcome Deterioration Over 2 Years. *Am J Sports Med.* Aug 2015;43(8):1965-75. doi:10.1177/0363546515585122

31. Denard PJ, Brady PC, Adams CR, Tokish JM, Burkhart SS. Preliminary Results of Arthroscopic Superior Capsule Reconstruction with Dermal Allograft. *Arthroscopy*. Jan 2018;34(1):93-99. doi:10.1016/j.arthro.2017.08.265

32. Henry P, Wasserstein D, Park S, et al. Arthroscopic Repair for Chronic Massive Rotator Cuff Tears: A Systematic Review. *Arthroscopy*. Dec 2015;31(12):2472-80.

doi:10.1016/j.arthro.2015.06.038

33. Brolin TJ, Updegrove GF, Horneff JG. Classifications in Brief: Hamada Classification of Massive Rotator Cuff Tears. *Clin Orthop Relat Res*. Nov 2017;475(11):2819-2823. doi:10.1007/s11999-017-5340-7

34. Hamada K, Yamanaka K, Uchiyama Y, Mikasa T, Mikasa M. A radiographic classification of massive rotator cuff tear arthritis. *Clin Orthop Relat Res*. Sep 2011;469(9):2452-60. doi:10.1007/s11999-011-1896-9

35. Chae SH, Jung TW, Lee SH, et al. Hidden Long Head of the Biceps Tendon Instability and Concealed Intratendinous Subscapularis Tears. *Orthop J Sports Med*. Jan

2020;8(1):2325967119898123. doi:10.1177/2325967119898123

36. Chauvet T, Haritinian E, Baudin F, Collotte P, Nové-Josserand L. The Invisible MGHL Test: Diagnostic Value and Benefits for the Repair of Retracted Subscapularis Tears. *Am J Sports Med*. Jul 2020;48(9):2144-2150. doi:10.1177/0363546520929695

37. Hsu KL, Chuang HC, Chang HM, et al. Association Between Medial Displacement of the Middle Glenohumeral Ligament and Subscapularis Tear Severity. *Orthop J Sports Med*. Apr 2022;10(4):23259671221083593. doi:10.1177/23259671221083593

38. Katthagen JC, Vap AR, Tahal DS, Horan MP, Millett PJ. Arthroscopic Repair of Isolated Partial- and Full-Thickness Upper Third Subscapularis Tendon Tears: Minimum 2-Year Outcomes

After Single-Anchor Repair and Biceps Tenodesis. *Arthroscopy*. Jul 2017;33(7):1286-1293. doi:10.1016/j.arthro.2017.01.027

39. Kim SJ, Jung M, Lee JH, Park JH, Chun YM. Arthroscopic repair of a significant (>50%) partialthickness subscapularis tear concomitant with a full-thickness supraspinatus tear: technical considerations for subscapularis repair (transtendon technique versus tear completion). *J Shoulder Elbow Surg*. Jun 2015;24(6):875-81. doi:10.1016/j.jse.2014.10.009

40. Lee TQ. Editorial Commentary: Precise Repair of Partial Subscapularis Tendon Tears Is Essential. *Arthroscopy*. May 2019;35(5):1314-1315. doi:10.1016/j.arthro.2019.02.022

41. Nové-Josserand L, Saffarini M, Hannink G, Carrillon Y. Influence of pre-operative tear size and tendon retraction on repair outcomes for isolated subscapularis tears. *Int Orthop*. Dec 2016;40(12):2559-2566. doi:10.1007/s00264-016-3299-8

42. Saltzman BM, Collins MJ, Leroux T, et al. Arthroscopic Repair of Isolated Subscapularis Tears: A Systematic Review of Technique-Specific Outcomes. *Arthroscopy*. Apr 2017;33(4):849-860. doi:10.1016/j.arthro.2016.10.020

43. Burkhart SS, Pranckun JJ, Hartzler RU. Superior Capsular Reconstruction for the Operatively Irreparable Rotator Cuff Tear: Clinical Outcomes Are Maintained 2 Years After Surgery. *Arthroscopy*. Feb 2020;36(2):373-380. doi:10.1016/j.arthro.2019.08.035

44. Frank RM, Cvetanovich G, Savin D, Romeo AA. Superior Capsular Reconstruction: Indications, Techniques, and Clinical Outcomes. *JBJS Rev*. Jul 2018;6(7):e10. doi:10.2106/jbjs.Rvw.17.00141

45. Galvin JW, Kenney R, Curry EJ, et al. Superior Capsular Reconstruction for Massive Rotator Cuff Tears: A Critical Analysis Review. *JBJS Rev*. Jun 2019;7(6):e1.

doi:10.2106/jbjs.Rvw.18.00072

46. Hartzler RU, Burkhart SS. Superior Capsular Reconstruction. *Orthopedics*. Oct 1 2017;40(5):271-280. doi:10.3928/01477447-20170920-02

47. Mihata T, Lee TQ, Hasegawa A, et al. Five-Year Follow-up of Arthroscopic Superior Capsule Reconstruction for Irreparable Rotator Cuff Tears. *J Bone Joint Surg Am*. Nov 6 2019;101(21):1921-1930. doi:10.2106/jbjs.19.00135

48. Smith TJ, Gowd AK, Kunkel J, Kaplin L, Waterman BR. Superior Capsular Reconstruction Provides Sufficient Biomechanical Outcomes for Massive, Irreparable Rotator Cuff Tears: A Systematic Review. *Arthroscopy*. Jan 2021;37(1):402-410. doi:10.1016/j.arthro.2020.09.007 49. Tokish JM, Makovicka JL. The Superior Capsular Reconstruction: Lessons Learned and Future Directions. *J Am Acad Orthop Surg*. Jul 1 2020;28(13):528-537. doi:10.5435/jaaos-d-19-00057

50. Ohta S, Komai O, Onochi Y. Outcomes of superior capsule reconstruction for massive rotator cuff tears and risk factors for postoperative retear. *Arch Orthop Trauma Surg*. Oct 2020;140(10):1319-1325. doi:10.1007/s00402-019-03316-2

51. Strauss EJ, McCormack RA, Onyekwelu I, Rokito AS. Management of failed arthroscopic rotator cuff repair. *J Am Acad Orthop Surg*. May 2012;20(5):301-9. doi:10.5435/jaaos-20-05-301

52. Erickson J, Lavery K, Monica J, Gatt C, Dhawan A. Surgical treatment of symptomatic superior labrum anterior-posterior tears in patients older than 40 years: a systematic review. *Am J Sports Med*. May 2015;43(5):1274-82. doi:10.1177/0363546514536874

53. McCormick F, Bhatia S, Chalmers P, Gupta A, Verma N, Romeo AA. The management of type II superior labral anterior to posterior injuries. *Orthop Clin North Am*. Jan 2014;45(1):121-8. doi:10.1016/j.ocl.2013.08.008

54. Belk JW, Kraeutler MJ, Houck DA, Chrisman AN, Scillia AJ, McCarty EC. Biceps tenodesis versus tenotomy: a systematic review and meta-analysis of level I randomized controlled trials. *J Shoulder Elbow Surg*. May 2021;30(5):951-960. doi:10.1016/j.jse.2020.11.012

55. Denard PJ, Lädermann A, Parsley BK, Burkhart SS. Arthroscopic biceps tenodesis compared with repair of isolated type II SLAP lesions in patients older than 35 years. *Orthopedics*. Mar 2014;37(3):e292-7. doi:10.3928/01477447-20140225-63

56. Dunne KF, Knesek M, Tjong VK, et al. Arthroscopic treatment of type II superior labral anterior to posterior (SLAP) lesions in a younger population: minimum 2-year outcomes are similar between SLAP repair and biceps tenodesis. *Knee Surg Sports Traumatol Arthrosc.* Jan 2021;29(1):257-265. doi:10.1007/s00167-020-05971-0

57. Ek ET, Shi LL, Tompson JD, Freehill MT, Warner JJ. Surgical treatment of isolated type II superior labrum anterior-posterior (SLAP) lesions: repair versus biceps tenodesis. *J Shoulder Elbow Surg*. Jul 2014;23(7):1059-65. doi:10.1016/j.jse.2013.09.030

58. Harris JD, Romeo AA. Arthroscopic management of the contact athlete with instability. *Clin Sports Med*. Oct 2013;32(4):709-30. doi:10.1016/j.csm.2013.07.007

59. Harris JD, Gupta AK, Mall NA, et al. Long-term outcomes after Bankart shoulder stabilization. *Arthroscopy*. May 2013;29(5):920-33. doi:10.1016/j.arthro.2012.11.010 60. Alkaduhimi H, Verweij LPE, Willigenburg NW, van Deurzen DFP, van den Bekerom MPJ. Remplissage With Bankart Repair in Anterior Shoulder Instability: A Systematic Review of the Clinical and Cadaveric Literature. *Arthroscopy*. Apr 2019;35(4):1257-1266. doi:10.1016/j.arthro.2018.10.117

61. Bhatia S, Frank RM, Ghodadra NS, et al. The outcomes and surgical techniques of the latarjet procedure. *Arthroscopy*. Feb 2014;30(2):227-35. doi:10.1016/j.arthro.2013.10.013 62. Bishop JY, Hidden KA, Jones GL, Hettrich CM, Wolf BR. Factors Influencing Surgeon's Choice of Procedure for Anterior Shoulder Instability: A Multicenter Prospective Cohort Study. *Arthroscopy*. Jul 2019;35(7):2014-2025. doi:10.1016/j.arthro.2019.02.035

63. Di Giacomo G, Itoi E, Burkhart SS. Evolving concept of bipolar bone loss and the Hill-Sachs lesion: from "engaging/non-engaging" lesion to "on-track/off-track" lesion. *Arthroscopy*. Jan 2014;30(1):90-8. doi:10.1016/j.arthro.2013.10.004

64. Dyrna FGE, Ludwig M, Imhoff AB, Martetschläger F. Off-track Hill-Sachs lesions predispose to recurrence after nonoperative management of first-time anterior shoulder dislocations. *Knee Surg Sports Traumatol Arthrosc.* Jul 2021;29(7):2289-2296. doi:10.1007/s00167-020-06176-1 65. Hurley ET, Jamal MS, Ali ZS, Montgomery C, Pauzenberger L, Mullett H. Long-term outcomes of the Latarjet procedure for anterior shoulder instability: a systematic review of

studies at 10-year follow-up. *J Shoulder Elbow Surg*. Feb 2019;28(2):e33-e39. doi:10.1016/j.jse.2018.08.028

66. Hurley ET, Lim Fat D, Farrington SK, Mullett H. Open Versus Arthroscopic Latarjet Procedure for Anterior Shoulder Instability: A Systematic Review and Meta-analysis. *Am J Sports Med*. Apr 2019;47(5):1248-1253. doi:10.1177/0363546518759540

67. Maio M, Sarmento M, Moura N, Cartucho A. How to measure a Hill-Sachs lesion: a systematic review. *EFORT Open Rev.* Apr 2019;4(4):151-157. doi:10.1302/2058-5241.4.180031
68. Milano G, Grasso A, Russo A, et al. Analysis of risk factors for glenoid bone defect in anterior shoulder instability. *Am J Sports Med.* Sep 2011;39(9):1870-6. doi:10.1177/0363546511411699

69. Moroder P, Schulz E, Wierer G, et al. Neer Award 2019: Latarjet procedure vs. iliac crest bone graft transfer for treatment of anterior shoulder instability with glenoid bone loss: a prospective randomized trial. *J Shoulder Elbow Surg*. Jul 2019;28(7):1298-1307. doi:10.1016/j.jse.2019.03.035

70. Rashid MS, Crichton J, Butt U, Akimau PI, Charalambous CP. Arthroscopic "Remplissage" for shoulder instability: a systematic review. *Knee Surg Sports Traumatol Arthrosc*. Feb 2016;24(2):578-84. doi:10.1007/s00167-014-2881-0

71. Werthel JD, Sabatier V, Schoch B, et al. Outcomes of the Latarjet Procedure for the Treatment of Chronic Anterior Shoulder Instability: Patients With Prior Arthroscopic Bankart Repair Versus Primary Cases. *Am J Sports Med*. Jan 2020;48(1):27-32. doi:10.1177/0363546519888909

Yang JS, Mazzocca AD, Cote MP, Edgar CM, Arciero RA. Recurrent Anterior Shoulder Instability With Combined Bone Loss: Treatment and Results With the Modified Latarjet Procedure. *Am J Sports Med*. Apr 2016;44(4):922-32. doi:10.1177/0363546515623929
Calvo E, Delgado C. Management of off-track Hill-Sachs lesions in anterior glenohumeral instability. *J Exp Orthop*. Mar 21 2023;10(1):30. doi:10.1186/s40634-023-00588-x

74. Provencher MT, Midtgaard KS, Owens BD, Tokish JM. Diagnosis and Management of Traumatic Anterior Shoulder Instability. *J Am Acad Orthop Surg*. Jan 15 2021;29(2):e51-e61. doi:10.5435/jaaos-d-20-00202

75. Calvo C, Calvo J, Rojas D, Valencia M, Calvo E. Clinical Relevance of Persistent Off-Track Hill-Sachs Lesion After Arthroscopic Latarjet Procedure. *Am J Sports Med*. Jul 2021;49(8):2006-2012. doi:10.1177/03635465211019979

76. Lee YJ, Kim C, Kim SJ, Yoon TH, Cho JY, Chun YM. Does an "Off-Track" Hill-Sachs Lesion That Is Misclassified as "Non-Engaging" Affect Outcomes From Bankart Repair Alone Compared With Bankart Repair Combined With Remplissage? *Arthroscopy*. Feb 2021;37(2):450-456. doi:10.1016/j.arthro.2020.09.037

77. Sheean AJ. Editorial Commentary: Remplissage Is Not Needed When Performing the Latarjet Procedure in the Setting of Off-Track Hill-Sachs Lesions: One of the Classics Continues to Get Better With Age (and Some Help From New Data). *Arthroscopy*. Aug 2021;37(8):2462-2464. doi:10.1016/j.arthro.2021.05.049

78. Brandariz RN, Gorodischer TD, Pasqualini I, Rossi LA, Tanoira I, Ranalletta M. The Latarjet Procedure Without Remplissage Is Effective to Restore Stability in Athletes With Glenoid Bone Defects Greater Than 25% and Off-Track Hill-Sachs Lesions. *Arthroscopy*. Aug 2021;37(8):2455-2461. doi:10.1016/j.arthro.2021.03.048

79. Verweij LPE, van Iersel TP, van Deurzen DFP, van den Bekerom MPJ, Floor S. "Nearly offtrack lesions" or a short distance from the medial edge of the Hill-Sachs lesion to the medial edge of the glenoid track does not seem to be accurate in predicting recurrence after an arthroscopic Bankart repair in a military population: a case-control study. *J Shoulder Elbow Surg*. Apr 2023;32(4):e145-e152. doi:10.1016/j.jse.2022.10.003

80. Arner JW, Peebles LA, Bradley JP, Provencher MT. Anterior Shoulder Instability Management: Indications, Techniques, and Outcomes. *Arthroscopy*. Nov 2020;36(11):2791-2793. doi:10.1016/j.arthro.2020.09.024

81. Lubowitz JH, Brand JC, Rossi MJ. Proper Evaluation of Bone Loss Determines Shoulder Instability Indications and Outcomes. *Arthroscopy*. Mar 2021;37(3):785-786. doi:10.1016/j.arthro.2021.01.004

82. Pécora JOR, Neves Junior AT, Roesler CRM, et al. Glenoid track evaluation by a validated finite-element shoulder numerical model. *Orthop Traumatol Surg Res.* Jun 2020;106(4):735-742. doi:10.1016/j.otsr.2020.03.004

83. Trivedi S, Pomerantz ML, Gross D, Golijanan P, Provencher MT. Shoulder instability in the setting of bipolar (glenoid and humeral head) bone loss: the glenoid track concept. *Clin Orthop Relat Res*. Aug 2014;472(8):2352-62. doi:10.1007/s11999-014-3589-7

84. Vopat ML, Hermanns CA, Midtgaard KS, et al. Imaging Modalities for the Glenoid Track in Recurrent Shoulder Instability: A Systematic Review. *Orthop J Sports Med*. Jun 2021;9(6):23259671211006750. doi:10.1177/23259671211006750

85. Barber FA, Howard MS. Editorial Commentary: Glenoid Track Instability Management Score or Instability Severity Index Score-Will This Decrease Latarjet Abuse? *Arthroscopy*. Jan 2020;36(1):68-70. doi:10.1016/j.arthro.2019.09.002

86. Di Giacomo G, Peebles LA, Pugliese M, et al. Glenoid Track Instability Management Score: Radiographic Modification of the Instability Severity Index Score. *Arthroscopy*. Jan 2020;36(1):56-67. doi:10.1016/j.arthro.2019.07.020

87. Younan Y, Wong PK, Karas S, et al. The glenoid track: a review of the clinical relevance, method of calculation and current evidence behind this method. *Skeletal Radiol*. Dec 2017;46(12):1625-1634. doi:10.1007/s00256-017-2687-5

88. Kibler WB, Sciascia A. Current Practice for the Diagnosis of a SLAP Lesion: Systematic Review and Physician Survey. *Arthroscopy*. Dec 2015;31(12):2456-69. doi:10.1016/j.arthro.2015.06.033

89. Forsythe B, Frank RM, Ahmed M, et al. Identification and treatment of existing copathology in anterior shoulder instability repair. *Arthroscopy*. Jan 2015;31(1):154-66. doi:10.1016/j.arthro.2014.06.014

90. Kraal T, Beimers L, The B, Sierevelt I, van den Bekerom M, Eygendaal D. Manipulation under anaesthesia for frozen shoulders: outdated technique or well-established quick fix? *EFORT Open Rev.* Mar 2019;4(3):98-109. doi:10.1302/2058-5241.4.180044

91. Rangan A, Hanchard N, McDaid C. What is the most effective treatment for frozen shoulder? *Bmj*. Aug 23 2016;354:i4162. doi:10.1136/bmj.i4162

92. Kim DH, Song KS, Min BW, Bae KC, Lim YJ, Cho CH. Early Clinical Outcomes of Manipulation under Anesthesia for Refractory Adhesive Capsulitis: Comparison with Arthroscopic Capsular Release. *Clin Orthop Surg*. Jun 2020;12(2):217-223. doi:10.4055/cios19027

93. Pandey V, Chidambaram R, Modi A, et al. Trends in Practice Among Shoulder Specialists in the Management of Frozen Shoulder: A Consensus Survey. *Orthop J Sports Med*. Oct 2022;10(10):23259671221118834. doi:10.1177/23259671221118834

94. Pandey V, Madi S. Clinical Guidelines in the Management of Frozen Shoulder: An Update! *Indian J Orthop*. Apr 2021;55(2):299-309. doi:10.1007/s43465-021-00351-3

95. Pensak M, Grumet RC, Slabaugh MA, Bach BR, Jr. Open versus arthroscopic distal clavicle resection. *Arthroscopy*. May 2010;26(5):697-704. doi:10.1016/j.arthro.2009.12.007

96. Singh B, Gulihar A, Bilagi P, et al. Magnetic resonance imaging scans are not a reliable tool for predicting symptomatic acromioclavicular arthritis. *Shoulder Elbow*. Oct 2018;10(4):250-254. doi:10.1177/1758573217724080

97. Aflatooni JO, Meeks BD, Froehle AW, Bonner KF. Biceps tenotomy versus tenodesis: patient-reported outcomes and satisfaction. *J Orthop Surg Res*. Feb 18 2020;15(1):56. doi:10.1186/s13018-020-1581-3

98. MacDonald P, Verhulst F, McRae S, et al. Biceps Tenodesis Versus Tenotomy in the Treatment of Lesions of the Long Head of the Biceps Tendon in Patients Undergoing Arthroscopic Shoulder Surgery: A Prospective Double-Blinded Randomized Controlled Trial. *Am J Sports Med*. May 2020;48(6):1439-1449. doi:10.1177/0363546520912212

99. Chalmers PN, Verma NN. Proximal Biceps in Overhead Athletes. *Clin Sports Med*. Jan 2016;35(1):163-79. doi:10.1016/j.csm.2015.08.009

100. Creech MJ, Yeung M, Denkers M, Simunovic N, Athwal GS, Ayeni OR. Surgical indications for long head biceps tenodesis: a systematic review. *Knee Surg Sports Traumatol Arthrosc*. Jul 2016;24(7):2156-66. doi:10.1007/s00167-014-3383-9

101. Kanbe K, Chiba J, Inoue Y, Taguchi M, Iwamatsu A. Analysis of clinical factors related to the efficacy of shoulder arthroscopic synovectomy plus capsular release in patients with rheumatoid arthritis. *Eur J Orthop Surg Traumatol*. Apr 2015;25(3):451-5. doi:10.1007/s00590-014-1570-5

102. Familiari F, Gonzalez-Zapata A, Iannò B, Galasso O, Gasparini G, McFarland EG. Is acromioplasty necessary in the setting of full-thickness rotator cuff tears? A systematic review. *J Orthop Traumatol*. Sep 2015;16(3):167-74. doi:10.1007/s10195-015-0353-z

103. Frank JM, Chahal J, Frank RM, Cole BJ, Verma NN, Romeo AA. The role of acromioplasty for rotator cuff problems. *Orthop Clin North Am*. Apr 2014;45(2):219-24. doi:10.1016/j.ocl.2013.12.003

104. Goutallier D, Postel JM, Bernageau J, Lavau L, Voisin MC. Fatty muscle degeneration in cuff ruptures. Pre- and postoperative evaluation by CT scan. *Clin Orthop Relat Res*. Jul 1994;(304):78-83.

105. Goutallier D, Postel JM, Gleyze P, Leguilloux P, Van Driessche S. Influence of cuff muscle fatty degeneration on anatomic and functional outcomes after simple suture of full-thickness tears. *J Shoulder Elbow Surg*. Nov-Dec 2003;12(6):550-4. doi:10.1016/s1058-2746(03)00211-8 106. Lippe J, Spang JT, Leger RR, Arciero RA, Mazzocca AD, Shea KP. Inter-rater agreement of the Goutallier, Patte, and Warner classification scores using preoperative magnetic resonance imaging in patients with rotator cuff tears. *Arthroscopy*. Feb 2012;28(2):154-9. doi:10.1016/j.arthro.2011.07.016

107. Schiefer M, Mendonça R, Magnanini MM, et al. Intraobserver and interobserver agreement of Goutallier classification applied to magnetic resonance images. *J Shoulder Elbow Surg*. Aug 2015;24(8):1314-21. doi:10.1016/j.jse.2015.02.011

108. Hamada K, Fukuda H, Mikasa M, Kobayashi Y. Roentgenographic findings in massive rotator cuff tears. A long-term observation. *Clin Orthop Relat Res.* May 1990;(254):92-6. 109. Chahal J, Mall N, MacDonald PB, et al. The role of subacromial decompression in patients undergoing arthroscopic repair of full-thickness tears of the rotator cuff: a systematic review and meta-analysis. *Arthroscopy*. May 2012;28(5):720-7. doi:10.1016/j.arthro.2011.11.022

ADDITIONAL RESOURCES

1. Harris JD, Frank JM, Jordan MA, et al. Return to sport following shoulder surgery in the elite pitcher: a systematic review. *Sports Health*. Jul 2013;5(4):367–76. doi:10.1177/1941738113482673

<u>1</u>2. Melis B, DeFranco MJ, Chuinard C, Walch G. Natural history of fatty infiltration and atrophy of the supraspinatus muscle in rotator cuff tears. *Clin Orthop Relat Res*. Jun 2010;468(6):1498-505. doi:10.1007/s11999-009-1207-x

<u>2</u>3. Paavola M, Malmivaara A, Taimela S, et al. Subacromial decompression versus diagnostic arthroscopy for shoulder impingement: <u>randomised</u><u>randomized</u>, placebo surgery controlled clinical trial. *BMJ*. Jul 19 2018;362:k2860. doi:10.1136/bmj.k2860

4. Saltzman BM, Jain A, Campbell KA, et al. Does the Use of Platelet-Rich Plasma at the Time of Surgery Improve Clinical Outcomes in Arthroscopic Rotator Cuff Repair When Compared With Control Cohorts? A Systematic Review of Meta-analyses. Arthroscopy. May 2016;32(5):906-18. doi:10.1016/j.arthro.2015.10.007

<mark>5. Strauss EJ, Barker JU, McGill K, Verma NN. The evaluation and management of failed distal clavicle excision. *Sports Med Arthrosc Rev*. Sep 2010;18(3):213–9. dei:10.1007/JSA.0b012e2181e802de</mark>

<mark>doi:10.1097/JSA.0b013e3181e892da</mark>

 Werner BC, Brockmeier SF, Miller MD. Etiology, Diagnosis, and Management of Failed SLAP Repair. J Am Acad Orthop Surg. Sep 2014;22(9):554-65. doi:10.5435/jaaos-22-09-554

<u>3</u>7. Yang JS, Mehran N, Mazzocca AD, Pearl ML, Chen VW, Arciero RA. Remplissage Versus Modified Latarjet for Off-Track Hill-Sachs Lesions With Subcritical Glenoid Bone Loss. *Am J Sports Med*. Jul 2018;46(8):1885-1891. doi:10.1177/0363546518767850



Reviewed / Approved by NIA Clinical Guideline Committee

Page **29** of **32** Shoulder Arthroscopy



POLICY HISTORY

<u>Date</u>	Summary
<u>May 2023</u>	Added the requirement of 6 weeks of physical therapy for partial
	rotator cuff repairs
	Added the requirement for no significant muscle atrophy or fatty
	infiltration for medium or large rotator cuff repairs
	Clarification of the indications for Latarjet or Remplissage
	<u>procedures</u>
	Added requirement for 50% decreased ROM in 2 planes, as
	compared to the opposite shoulder, for frozen shoulder surgery
	• Added requirement for a chest X-ray in the past 12 months for
	frozen shoulder surgery
May 2022	Updated background and references
	• Further defined the glenoid track verbiage for "off-track" and "on-
	track" Hill-Sachs lesions (bony defects of the humeral head)
	• Clarified individual is medically stable and <i>optimized for surgery</i>
	Revised Partial-Thickness Rotator Cuff Tear or Calcific Tendinitis to
	"include <i>two</i> of the following criteria"
	Revised criteria for Latarjet or Remplissage to "Recurrent anterior
	dislocations"
	Non-operative treatment for small RCT revised to ONE of the
	following (previously "at least one")
	Revised 3 months to 12 weeks throughout
	Replace "patient" with "individual" where appropriate
	Added:
	• Evaluation of pain prior total shoulder arthroplasty as indication for
	a diagnostic arthroscopy
	Cortisone injection within 12 weeks of a rotator cuff repair or
	revision as a contraindication
	Added more specific indications for repair of a subscapularis
	rotator cuff tear
	Physical examination findings requirement for SLAP tears
	Criteria for loose body removal
	• "performed in conjunction with a subscapularis tendon repair" to
	criteria for Long Head Biceps Tenotomy/Tenodesis
	Deleted:
	Requirement for a cortisone injection for calcific tendinopathy
	Deleted cortisone injections from lists of treatment modalities
	IA joint corticosteroid injection from non-operative treatments for
	LHB Tenotomy/Tenodesis



|--|

Page **31** of **32** Shoulder Arthroscopy



Reviewed / Approved by NIA Clinical Guideline Committee

GENERAL INFORMATION

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

Disclaimer: Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates ("Magellan"). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.

Disclaimer: National Imaging Associates, Inc. (NIA) authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Evolent Health LLC subsidiaries including, but not limited to, National Imaging Associates ("NIA"). The policies constitute only the reimbursement and coverage guidelines of NIA. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. NIA reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.

Page **32** of **32** Shoulder Arthroscopy

