

## AmeriHealth Caritas Louisiana

|   |  |
|---|--|
| <b>National Imaging Associates, Inc.*</b>     |  |
| <b>Clinical guidelines</b><br><b>NECK CTA</b> | <b>Original Date: September 1997</b>     |
| <b>CPT Codes: 70498</b>                       | <b>Last Revised Date: April 2021</b>     |
| <b>Guideline Number: NIA_CG_012-1</b>         | <b>Implementation Date: January 2022</b> |

### INDICATIONS FOR NECK CTA

Patients with claustrophobia, limited ability to cooperate or an implanted device may be better suited for CTA, whereas those with extensive calcification, renal disease or iodine contrast allergy should have MRA (Adla, 2015).

#### For evaluation of known or suspected extracranial vascular disease

##### Cerebrovascular Disease

- Recent ischemic stroke or transient ischemic attack ([Robertson, 2020; SalmelaACR, 2017; 2019](#); Sanelli, 2014).
- Known or suspected vertebrobasilar insufficiency (VBI) in patients with symptoms such as dizziness, vertigo, headaches, diplopia, blindness, vomiting, ataxia, weakness in both sides of the body, or abnormal speech (Yang, 2005; Lima-Neto, 2017; Searls, 2012)**
- Asymptomatic patients with an abnormal ultrasound of the neck or carotid duplex imaging (e.g., carotid stenosis  $\geq 70\%$ , technically limited study, aberrant direction of flow in the carotid or vertebral arteries) (Brott, 2011; DaCosta, 2019; Marquardt, 2010).
- Symptomatic patients with an abnormal ultrasound of the neck or carotid duplex imaging (e.g., carotid stenosis  $\geq 50\%$ , technically limited study, aberrant direction of flow in the carotid or vertebral arteries) ([BAAN, 2010; Brott, 2011; Chaturvedi, 2005](#); Rerkasem, 2011).

##### Aneurysm screening

- Screening for aneurysm in Loeys-Dietz syndrome\*\*, fibromuscular dysplasia or spontaneous coronary arteries dissection (SCAD) (Hayes, 2018; Hitchcock, 2014; Macaya, 2019; MacCarrick, 2014)**

\* National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.

1— Neck CTA

[Copyright](#) © 2019-2021 National Imaging Associates, Inc., All Rights Reserved

~~\*\*For Loeyzs-Dietz imaging should be repeated at least every two years Screening for aneurysm in fibromuscular dysplasia or spontaneous coronary arteries dissection (SCAD) (Hayes, 2018; Hitchcock, 2014; Macaya, 2019).~~

#### Tumor/pulsatile mass

- Pulsatile mass on exam (Aulino, 2019)

#### Known

- ~~Carotid body tumors, or other masses such as a paraganglioma, arteriovenous fistula pseudoaneurysm, atypical lymphovascular malformation (Nguyen, 2011).~~  
**Note:** Ultrasound (US) may be used to identify a mass overlying or next to an artery in initial work up of a pulsatile mass.

#### Other extracranial vascular disease

- Takayasu arteritis based on findings in other blood vessels on previous imaging (Zhu, 2012)
- 
- Giant cell arteritis with suspected extracranial involvement (Abdel Razek, 2014; Halbach, 2018; Khan, 2015; Koster, 2018; Khan, 2015; Halbach, 2018) (Abdel Razek, 2014; Koster, 2018)
- Subclavian steal syndrome when ultrasound is positive or indeterminate ~~or OR~~ for planning interventions (Potter, 2014)
- Suspected carotid or vertebral artery dissection; due to trauma or spontaneous due to weakness of vessel wall ~~leading to dissection~~ (Franz, 2012; Shakir, 2016)
- Horner's syndrome (miosis, ptosis, and anhidrosis) (Kim, 2012).
- For evaluation of pulsatile tinnitus (subjective or objective) for vascular etiology (Pegge, 2017)
- Known extracranial vascular disease that needs follow-up or further evaluation

#### Pre-operative/procedural evaluation

- Pre-operative evaluation for a planned surgery or procedure if the imaging provides diagnostic information that is not available on prior studies (provider should be referred to the health plan for nondiagnostic surgical planning studies).
- 

#### Post-operative/procedural evaluation (e.g. carotid endarterectomy):

- A follow-up study may be needed to help evaluate a patient's progress after treatment, procedure, intervention, or surgery. Documentation requires a medical reason that clearly indicates why additional imaging is needed for the type and area(s) requested.

## **INDICATIONS FOR COMBINATION STUDIES**

### **Neck CTA/Brain CTA**

- Recent ischemic stroke or transient ischemic attack (TIA) ([Robertson, 2020; Salmela, 2017](#)[ACR, 2017; 2019](#); Wintermark, 2013)
- Known or suspected vertebrobasilar insufficiency (VBI) in patients with symptoms such as dizziness, vertigo, headaches, diplopia, blindness, vomiting, ataxia, and weakness in both sides of the body, or abnormal speech (Lima-Neto 2017; Searls, 2012)
- Suspected carotid or vertebral artery dissection; due to trauma or spontaneous due to weakness of vessel wall [leading to dissection](#) (Franz, 2012; Shakir, 2016)
- Asymptomatic patients with an abnormal ultrasound of the neck or carotid duplex imaging (e.g., carotid stenosis  $\geq 70\%$ , technically limited study, aberrant direction of flow in the carotid or vertebral arteries) and patient is surgery or angioplasty candidate (Brott, 2011; DaCosta, 2019; Marquardt, 2010)
- Symptomatic patients with an abnormal ultrasound of the neck or carotid duplex imaging (e.g., carotid stenosis  $\geq 50\%$ , technically limited study, aberrant direction of flow in the carotid or vertebral arteries) and patient is surgery or angioplasty candidate ([AAN, 2010](#); Brott, 2011; [Chaturvedi, 2005](#); Rerkasem, 2011)
- Pulsatile tinnitus (subjective or objective) for vascular etiology (Pegge, 2017)

---



---

## BACKGROUND

For vascular disease, MRA and CTA are generally comparable. [There is no](#) current literature [comparing](#) [compares](#) the efficacy of contrast enhanced CT to CTA or MRI and MRA for evaluation of pulsatile neck mass, so any are approvable (Guneyli, 2014). CTA may be complementary to CT in the following settings: evaluation of a pulsatile neck mass to assess vascular detail when needed; assessment of relevant vascular anatomy for pre-procedural evaluation; vascular supply to tumors and vessel encasement and narrowing by tumors; extent of disease in vasculitis; and to help determine the nature and extent of congenital or acquired vascular anomalies.

**CTA and dissection-** Craniocervical dissections can be spontaneous or traumatic. Patients with blunt head or neck trauma who meet Denver Screening criteria should be assessed for cerebrovascular injury (although about 20% will not meet criteria). The criteria include: focal or lateralizing neurological deficits (not explained by head CT), infarct on head CT, face, basilar skull, or cervical spine fractures, cervical hematomas that are not expanding, glasgow coma score less than 8 without CT findings, massive epistaxis, cervical bruit or thrill (Franz, 2012; Liang, 2013; Mundinger, 2013; Simon, 2019). Spontaneous dissection presents with headache, neck pain with neurological signs or symptoms. There is often minor trauma or precipitating factor (e.g., exercise, neck manipulation). Dissection is thought to occur due to weakness of the vessel wall, and there may be an underlying connective tissue disorder. Dissection of the extracranial vessels can extend intracranially and/or lead to thrombus, which

can migrate into the intracranial circulation causing ischemia. Therefore, MRA of the head and neck is warranted (Nash, 2019; Shakir, 2016).

**CTA and recent stroke or transient ischemic attack (TIA)**-A stroke or central nervous system infarction is defined as “brain, spinal cord, or retinal cell death attributable to ischemia, based on neuropathological, neuroimaging, and/or clinical evidence of permanent injury. ... Ischemic stroke specifically refers to central nervous system infarction accompanied by overt symptoms, whereas silent infarction causes no known symptoms” (Sacco, 2013).<sup>12</sup> If imaging or pathology is not available, a clinical stroke is diagnosed by symptoms persisting for more than 24 hours. Ischemic stroke can be further classified by the type and location of ischemia and the presumed etiology of the brain injury. These include large-artery atherosclerotic occlusion (extracranial or intracranial), cardiac embolism, small-vessel disease and less commonly dissection, hypercoagulable states, sickle cell disease and undetermined causes (Kernan, 2014). TIAs in contrast, “are a brief episode of neurological dysfunction caused by focal brain or retinal ischemia, with clinical symptoms typically lasting less than one hour, and without evidence of acute infarction on imaging” (Easton, 2009).<sup>13</sup> On average, the annual risk of future ischemic stroke after a TIA or initial ischemic stroke is 3–4%, with an incidence as high as 11% over the next 7 days and 24–29% over the following 5 years. This has significantly decreased in the last half century due to advances in secondary prevention (Hong, 2011).

When revascularization therapy is not indicated or available in patients with an ischemic stroke or TIA, the focus of the work-up is on secondary prevention. This includes noninvasive vascular imaging to identify the underlying etiology, assess immediate complications and risk of future stroke. The majority of stroke evaluations take place in the inpatient setting. Admitting TIA patients is reasonable if they present within 72 hours and have an ABCD(2) score  $\geq 3$ , indicating high risk of early recurrence, or the evaluation cannot be rapidly completed on an outpatient basis (Easton, 2009). Minimally, both stroke and TIA should have an evaluation for high-risk modifiable factors, such as carotid stenosis atrial fibrillation, as the cause of ischemic symptoms (Kernan, 2014). Diagnostic recommendations include neuroimaging evaluation as soon as possible, preferably with magnetic resonance imaging, including DWI; noninvasive imaging of the extracranial vessels should be performed, and noninvasive imaging of intracranial vessels is reasonable (Wintermark, 2013).

Patients with a history of stroke and recent work up with new signs or symptoms indicating progression or complications of the initial CVA should have repeat brain imaging as an initial study. Patients with remote or silent strokes discovered on imaging should be evaluated for high-risk modifiable risk factors based on the location and type of the presumed etiology of the brain injury (Easton, 2009; Hong, 2011; Kernan, 2014; Sacco, 2013; Wintermark, 2013).

#### POLICY HISTORY

| Date           | Summary   |
|----------------|---|
| April-May 2021 | <u>Updated references</u><br><u>Added:</u><br>• <u>Loeys-Dietz syndrome to aneurysm screening section</u> |

|                 |  |
|-----------------|--|
|                 | <ul style="list-style-type: none"> <li>● <u>Known or suspected vertebrobasilar insufficiency (VBI) in patients with symptoms such as dizziness, vertigo, headaches, diplopia, blindness, vomiting, ataxia and weakness in both sides of the body, or abnormal speech – which was before only in the combo section</u></li> <li>● <u>Pulsatile mass on exam</u></li> <li>● <u>For evaluation of pulsatile tinnitus (subjective or objective) for vascular etiology - which was before only in the combo section</u></li> <li>● <u>Pre-operative evaluation for a planned surgery or procedure if the imaging provides diagnostic information that is not available on prior studies (provider should be referred to the health plan for nondiagnostic surgical planning studies).</u></li> <li>● <u>Clarified:</u></li> <li>● <u>Giant cell arteritis with suspected extracranial involvement</u></li> <li>● <u>Known carotid body tumors, or other masses such as a paraganglioma, arteriovenous fistula pseudoaneurysm, atypical lymphovascular malformation</u></li> </ul> |
| <u>May 2020</u> | <p><u>Clarified:</u></p> <ul style="list-style-type: none"> <li>● <u>Patients with claustrophobia, limited ability to cooperate or an implanted device may be better suited for CTA, whereas those with extensive calcification, renal disease or iodine contrast allergy should have MRA</u></li> <li>● <u>Recent ischemic stroke or transient ischemic attack (also in combo section)</u></li> <li>● <u>Pulsatile mass on exam after ultrasound (US)</u></li> <li>● <u>Takayasu arteritis based on findings in other blood vessels on previous imaging</u></li> <li>● <u>Giant cell arteritis</u></li> <li>● <u>Known or suspected vertebrobasilar insufficiency (VBI) in patients with symptoms such as dizziness, vertigo, headaches, diplopia, blindness, vomiting, ataxia and weakness in both sides of the body, or abnormal speech</u></li> <li>● <u>Suspected carotid or vertebral artery dissection; due to trauma or spontaneous due to weakness of vessel wall leading to dissection (combo section)</u></li> </ul> <p><u>Added:</u></p>                         |

|                   |   |
|-------------------|---|
|                   | <ul style="list-style-type: none"> <li>Known extracranial vascular disease that needs follow-up or further evaluation</li> <li>Spontaneous coronary arteries dissection (SCAD) in screening for aneurysm</li> <li>Suspected carotid or vertebral artery dissection; due to trauma or spontaneous due to weakness of vessel wall leading to dissection</li> <li>Horner's syndrome (miosis, ptosis, and anhidrosis)</li> <li>Known extracranial vascular disease that needs follow-up or further evaluation</li> </ul> <p>-</p> <p><u>Deleted:</u></p> <ul style="list-style-type: none"> <li>Ehlers-Danlos syndrome and neurofibromatosis in screening for aneurysm</li> </ul> |
| <u>April 2019</u> | <ul style="list-style-type: none"> <li>Added initial statement describing the use of CTA versus MRA</li> <li>Suspected or known disease: Added "Giant cell arteritis" and "Subclavian steal syndrome when ultrasound is positive or indeterminate or for planning interventions"</li> <li>"Known or suspected tumor/pulsatile mass": Added 'pulsatile';</li> <li>Neck CTA/Brain CTA: Added Denver screening criteria to assess for cerebrovascular injury</li> <li>Added background information describing CTA and MRA as complimentary information to CT or MRI</li> </ul>   |

#### April 2019

- Added initial statement describing the use of CTA versus MRA
- Suspected or known disease: Added "Giant cell arteritis" and "Subclavian steal syndrome when ultrasound is positive or indeterminate or for planning interventions"
- "Known or suspected tumor/pulsatile mass": Added 'pulsatile';
- Neck CTA/Brain CTA: Added Denver screening criteria to assess for cerebrovascular injury
- Added background information describing CTA and MRA as complimentary information to CT or MRI

#### May 2020

##### Clarified:

- Patients with claustrophobia, limited ability to cooperate or an implanted device may be better suited for CTA, whereas those with extensive calcification, renal disease or iodine contrast allergy should have MRA
- Recent ischemic stroke or transient ischemic attack (also in combo section)
- Pulsatile mass on exam after ultrasound (US)
- Takayasu arteritis based on findings in other blood vessels on previous imaging
- Giant cell arteritis

#### 6— Neck CTA

- Known or suspected vertebrobasilar insufficiency (VBI) in patients with symptoms such as dizziness, vertigo, headaches, diplopia, blindness, vomiting, ataxia and weakness in both sides of the body, or abnormal speech
- Suspected carotid or vertebral artery dissection; due to trauma or spontaneous due to weakness of vessel wall leading to dissection (combo section)

**Added:**

- Known extracranial vascular disease that needs follow up or further evaluation
- Spontaneous coronary arteries dissection (SCAD) in screening for aneurysm
- Suspected carotid or vertebral artery dissection; due to trauma or spontaneous due to weakness of vessel wall leading to dissection
- Horner's syndrome (miosis, ptosis, and anhidrosis)
- Known extracranial vascular disease that needs follow up or further evaluation

-

**Deleted:**

- Ehlers-Danlos syndrome and neurofibromatosis in screening for aneurysm



## REFERENCES

Abdel Razek AA, Alvarez H, Bagg S, et al. Imaging spectrum of CNS vasculitis. *Radiographics*. 2014 Jul-Aug; 34(4):873-94.

Adla T, Adlova R. Multimodality imaging of carotid stenosis. *Int J Angiol*. 2015 Sep; 24(3):179-84.

[Aulino JM, Kirsch CFE, Burns J, et al. ACR Appropriateness Criteria® Neck Mass-Adenopathy. Journal of the American College of Radiology. 2019;16\(5\):S150-S160.](#)  
[doi:10.1016/j.jacr.2019.02.025](#)

[American Academy of Neurology \(AAN\). Carotid Endarterectomy An evidence-based review. 2010.](#)

[American College of Radiology \(ACR\). ACR Appropriateness Criteria® Cerebrovascular Disease. 2017.](#)

[American College of Radiology \(ACR\). ACR Appropriateness Criteria® Cerebrovascular Disease-Child. 2019.](#)

Brott TG, Halperin JL, Abbara S, et al. ASA / ACCF / AHA / AANN / AANS / ACR / ASNR / CNS / SAIP / SCAI / SIR / SNIS / SVM / SVS guideline on the management of patients with extracranial carotid and vertebral artery disease: Executive summary. *Circulation*. 2011; 124:489-532.  
<http://www.ncbi.nlm.nih.gov/pubmed/21282493>.

[Chaturvedi S, Bruno A, Feasby T, et al. Carotid endarterectomy--an evidence-based review: Report of the therapeutics and technology assessment subcommittee of the American Academy of Neurology. Neurology. 2005;65\(6\):794-801.](#)  
[doi:10.1212/01.wnl.0000176036.07558.82](#)

DaCosta M, Surowiec SM. Carotid Endarterectomy. *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2019.

Easton JD, Saver JL, Albers GW, et al. Definition and evaluation of transient ischemic attack. *Stroke*. 2009; 40(6):2276.

Franz RW, Willette PA, Wood MJ, et al. A systematic review and meta-analysis of diagnostic screening criteria for blunt cerebrovascular injuries. *J Am Coll Surg*. March 2012; 214(3):313-327.

Guneyli S, Ceylan N, Bayraktaroglu S, et al. Imaging findings of vascular lesions in the head and neck. *Diagn Interv Radiol.* 2014 Sep-Oct; 20(5):432-7.

**Halbach C, McClelland CM, Chen J, et al. Use of noninvasive imaging in giant cell arteritis. Asia Pac J Ophthalmol (Phila). 2018;7(4):260-264. doi:10.22608/APO.2018133**

Hayes SN, Kim ESH, Saw J, et al. Spontaneous coronary artery dissection: current state of the science: A scientific statement from the American Heart Association. *Circulation.* 2018 May 8; 137(19):e523-e557. doi:10.1161/CIR.0000000000000564. Epub 2018 Feb 22.

Hitchcock E, Gibson WT. A review of the genetics of intracranial berry aneurysms and implications for genetic counseling. *J Genet Couns.* 2017;26(1):21–31. doi:10.1007/s10897-016-0029-8.

Hong KS, Yegiaian S, Lee M, et al. Declining stroke and vascular event recurrence rates in secondary prevention trials over the past 50 years and consequences for current trial design. *Circulation.* 2011 May 17;123(19):2111-9. Epub 2011 May 2. doi: 10.1161/CIRCULATIONAHA.109.934786.

Jadhav AP, Jovin TG. Vascular Imaging of the Head and Neck. *Semin Neurol.* 2012; 32(04):401-410. doi: 10.1055/s-0032-1331811.

Kernan WN, Ovbiagele B, Black HR, et al. Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke.* 2014 Jul; 45(7):2160-236. Epub 2014 May. doi: 10.1161/STR.0000000000000024.

**Khan A, Dasgupta B. Imaging in giant cell arteritis. Curr Rheumatol Rep. 2015;17(8):52. doi:10.1007/s11926-015-0527-y**

Kim JD, Hashemi N, Gelman R, et al. Neuroimaging in ophthalmology. *Saudi J Ophthalmol.* 2012; 26(4):401–407. doi:10.1016/j.sjopt.2012.07.001.

Koster MJ, Matteson EL, Warrington KJ. Large-vessel giant cell arteritis: Diagnosis, monitoring, and management. *Rheumatol.* 2018; 57(suppl2):ii32-42.

Liang T, Tso DK, Chiu RY, et al. Imaging of blunt vascular neck injuries: A review of screening and imaging modalities. *AJR Am J Roentgenol.* 2013 Oct; 201(4):884-92.

Lima-Neto AC, Bittar R, Gattas GS, et al. Pathophysiology and diagnosis of vertebrobasilar insufficiency: A review of the literature. *Int Arch Otorhinolaryngol.* 2017 July; 21(3):302-07.

Macaya F, Moreu M, Ruiz-Pizarro V, et al. Screening of extra-coronary arteriopathy with magnetic resonance angiography in patients with spontaneous coronary artery dissection: A single-centre experience. *Cardiovasc Diagn Ther.* 2019 Jun; 9(3):229-238. doi: 10.21037/cdt.2019.04.09.

[MacCarrick G, Black JH, Bowdin S, et al. Loeys–Dietz syndrome: A primer for diagnosis and management. \*Genet Med.\* 2014;16\(8\):576-587. doi:10.1038/gim.2014.11](#)

Marquardt L, Geraghty OC, Mehta Z, et al. Low risk of ipsilateral stroke in patients with asymptomatic carotid stenosis on best medical treatment: A prospective, population-based study. *Stroke.* 2010; 41(1):e11.

Mundinger GS, Dorafshar AH, Gilson MM, et al. Blunt-mechanism facial fracture patterns associated with internal carotid artery injuries: recommendations for additional screening criteria based on analysis of 4,398 patients. *J Oral Maxillofac Surg.* December 2013; 71(12):2092-2100.

Nash M, Rafay MF. Craniocervical arterial dissection in children: Pathophysiology and management. *Pediatr Neurol.* 2019 Jun; 95:9-18. Epub 2019 Feb 2. doi:10.1016/j.pediatrneurol.2019.01.020.

Nguyen RP, Shah LM, Quigley EP, et al. Carotid body detection on CT angiography. *AJNR Am J Neuroradiol.* 2011 June; 32(6):1096-99.

Pegge SAH, Steens SCA, Kunst HPM, et al. Pulsatile tinnitus: Differential diagnosis and work-up. *Curr Radiol Rep.* 2017; 5(1):5.

Potter BJ, Pinto DS. Subclavian steal syndrome. *Circulation.* 2014 Jun 3; 129(22):2320-23.

Rerkasem K, Rothwell PM. Carotid endarterectomy for symptomatic carotid stenosis. *Cochrane Database Syst Rev.* 2011.

[Robertson RL, Palasis S, Rivkin, MJ, et al. ACR Appropriateness Criteria® Cerebrovascular Disease-Child. \*J Am Coll Radiol.\* 2020;17\(5S\):S36-S54. doi:10.1016/j.jacr.2020.01.036](#)

Sacco RL, Kasner SE, Broderick JP, et al. An updated definition of stroke for the 21st century: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke.* 2013; 44:2064–2089.

[Salmela MB, Mortazavi S, Jagadeesan, BD, et al. ACR Appropriateness Criteria® Cerebrovascular Disease. \*J Am Coll Radiol.\* 2017;14\(5S\):S34-S61. doi:10.1016/j.jacr.2017.01.051](#)

Sanelli PC, Sykesa JB, Ford AL, et al. Imaging and treatment of patients with acute stroke: An evidence-based review. *AJNR Am J Neuroradiol.* 2014; 35:1045-1051.

Searls DE, Pazdera L, Korbel E, et al. Symptoms and signs of posterior circulation ischemia in the new England medical center posterior circulation registry. *Arch Neurol.* 2012; 69(3):346.


Shakir HJ, Davies JM, Shallwani H, et al. Carotid and vertebral dissection imaging. *Curr Pain Headache Rep.* 2016 Dec; 20(12):68. Review.

Simon LV, [Nassar, AK](#), Mohseni M. Vertebral Artery Injury. *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2021. <https://www.ncbi.nlm.nih.gov/books/NBK470363/>.

Wintermark M, Sanelli PC, Albers GW, et al. Imaging recommendations for acute stroke and transient ischemic attack patients: A joint statement by the American Society of Neuroradiology, the American College of Radiology, and the Society of NeuroInterventional Surgery. *AJNR Am J Neuroradiol.* 2013 Nov-Dec; 34(11):E117-27. Epub 2013 Aug 1. doi: 10.3174/ajnr.A3690.

[Yang CW, Carr JC, Futterer SF, et al. Contrast-enhanced MR angiography of the carotid and vertebrobasilar circulations. AJNR Am J Neuroradiol. 2005;26\(8\):2095-2101.](#)

Zhu FP, Luo S, Wang ZJ, et al. Takayasu arteritis: Imaging spectrum at multidetector CT angiography. *Br J Radiol.* 2012 Dec; 85(1020):e1282-1292.

Reviewed / Approved by  M. Atif Khalid, M.D., Medical Director, Radiology  
[Reviewed / Approved by](#) [ee](#)

## **GENERAL INFORMATION**

**It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.**

**Disclaimer:** Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates ("Magellan"). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.