

AmeriHealth Caritas Louisiana

National Imaging Associates, Inc.*	
Clinical guidelines THORACIC SPINE CT	Original Date: September 1997
CPT Codes: 72128, 72129, 72130	Last Revised Date: April-March 2021
Guideline Number: NIA_CG_043	Implementation Date: January 2023 2

INDICATIONS FOR THORACIC SPINE CT

If there is a combination request for an overlapping body part, either requested at the same time or sequentially (within the past 3 months) the results of the prior study should be:

- Inconclusive or show a need for additional or follow up imaging evaluation OR
- The office notes should clearly document an indication why overlapping imaging is needed and how it will change management for the patient (the entire spinal cord and/or autonomic postganglionic chain must be assessed)

(*Unless approvable in the combination section as noted in the guidelines)

(Combination requests at end of the document)

For evaluation of neurologic deficits when Thoracic Spine MRI is contraindicated or inappropriate¹⁻³ (~~Acharya, 2019; ACR, 2013; NASS, 2010~~)

- With any of the following new neurological deficits documented on physical exam
 - Extremity muscular weakness (and not likely caused by plexopathy, or peripheral neuropathy)^{4, 5}
 - Pathologic (e.g., Babinski, ~~Lhermitte's sign~~, Chaddock Sign,⁷) or abnormal reflexes⁶ (~~Teoli, 2021~~)
 - Absent/decreased sensory changes along a particular thoracic dermatome (nerve distribution): pin prick, touch, vibration, proprioception, or temperature
 - Upper or lower extremity increase muscle tone/spasticity and likely localized to the thoracic spinal cord
 - New onset bowel or bladder dysfunction (e.g., retention or incontinence) - not related to an inherent bowel or bladder process
 - Gait abnormalities (see [Table 1](#) for more details)
- Suspected cord compression with any neurological deficits as listed above

* National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.

1— Thoracic Spine CT

© 2019-202~~1~~ National Imaging Associates, Inc., All Rights Reserved

- Toe walking in a child when associated with upper motor neuron signs including hyperreflexia, spasticity; or orthopedic deformity with concern for spinal cord pathology (e.g., pes cavus, clawed toes, leg or foot length deformity (excluding tight heel cords))

For evaluation of back pain with any of the following when Thoracic Spine MRI is contraindicated⁷⁻¹⁰

~~(AANSCNS, 2014; Allegri, 2016; Jarvik, 2015; Last, 2009)~~

- With new or worsening objective neurologic deficits on exam, as above
- Failure of conservative treatment* for at least six (6) weeks within the last six (6) months¹¹
~~(ACR, 2013; Eubanks, 2010)~~
- With progression or worsening of symptoms during the course of conservative treatment*
- With an abnormal electromyography (EMG) or nerve conduction study (if performed) indicating a thoracic radiculopathy. (EMG is not recommended to determine the cause of axial lumbar, thoracic, or cervical spine pain~~(NASS, 2013)~~)¹²
- Isolated back-thoracic pain in pediatric population¹³ ~~-(ACR, 2016)~~— conservative care not required if red flags present ~~(see combination request below cervical and lumbar spine may also be indicated)~~
 - Red flags that prompt imaging should include the presence of: age 5 or younger, constant pain, pain lasting >4 weeks, abnormal neurologic examination, early morning stiffness and/or gelling; night pain that prevents or disrupts sleep; radicular pain; fever; weight loss; malaise; postural changes (e.g., kyphosis or scoliosis); and limp (or refusal to walk in a younger child <5yo) AND initial radiographs have been performed^{14, 15} ~~(Bernstein, 2007; Feldman, 2006)~~
- ~~Back pain associated with suspected inflammation, infection, or malignancy~~

As part of initial pre-operative/post-operative/procedural evaluation (“CT best examination to assess for hardware complication, extent of fusion”^{16, 17} ~~-(ACR, 2015; Rao, 2018)~~ and MRI for cord, nerve root compression, disc pathology, or post-op infection)

If ordered by Neurosurgeon or orthopedic surgeon for purposes of surgical planning. A contraindication to MRI is not required

- For preoperative evaluation/planning
- CT discogram
- CSF leak highly suspected and supported by patient history and/or physical exam findings (leak (known or suspected spontaneous (idiopathic) intracranial hypotension (SIH), post lumbar puncture headache, post spinal surgery headache, orthostatic headache, rhinorrhea or otorrhea, or cerebrospinal-venous fistula -preferred exam CT myelogram)¹⁸ ~~(Starling, 2013)~~
- Prior to spinal cord stimulator to exclude canal stenosis if no prior imaging of the thoracic spine has been done recently and MRI is contraindicated

- A follow-up study may be needed to help evaluate a patient's progress after treatment, procedure, intervention, or surgery in the last 6 months. Documentation requires a medical reason that clearly indicates why additional imaging is needed for the type and area(s) requested (routine surveillance post-op not indicated without symptoms)
- ~~Changing neurologic status post-operatively~~
- Surgical infection as evidenced by signs/symptoms, laboratory, or prior imaging findings
- New or changing neurological deficits or symptoms post-operatively ~~Residual or new neurological deficits or symptoms~~^{16, 19} ~~(Rao, 2018)~~ see neurological deficit section above
- When combo requests are submitted (i.e., MRI and CT of the spine), the office notes should clearly document the need for both studies to be done simultaneously, i.e., the need for both soft tissue and bony anatomy is required²⁰ ~~(Fisher, 2013)~~
 - Combination requests where both thoracic spine CT and MRI thoracic spine are both approvable (not an all-inclusive list):
 - OPLL (Ossification of posterior longitudinal ligament)
 - Most common in cervical spine (rare but more severe in thoracic spine)²¹ ~~(Choi, 2011)~~
 - Pathologic or complex fractures
 - Malignant process of spine with both bony and soft tissue involvement
 - Clearly documented indication for bony and soft tissue abnormality where assessment will change management for the patient

For evaluation of suspected myelopathy when Thoracic Spine MRI is contraindicated²²⁻²⁶
~~(ACR, 2015; Behrbalk, 2013; Davies, 2018; Vilaca, 2016; Waly, 2017)~~

- Does NOT require conservative care
- Progressive symptoms including unsteadiness; broad-based gait; increased muscle tone; pins and needles sensation; weakness and wasting of the lower limbs; and diminished sensation to light touch, temperature, proprioception, and vibration; limb hyperreflexia and pathologic reflexes; bowel and bladder dysfunction in more severe cases.
- ~~Progressive symptoms including hand clumsiness, worsening handwriting, difficulty with grasping and holding objects, diffuse numbness in the hands, pins and needles sensation, increasing difficulty with balance and ambulation~~
- Any of the neurological deficits as noted above

For evaluation of trauma or acute injury²⁷
~~(ACR, 2018)~~

- Presents with any of the following neurological deficits as above
- With progression or worsening of symptoms during the course of conservative treatment*
- History of underlying spinal abnormalities (i.e., ankylosing spondylitis, diffuse idiopathic skeletal hyperostosis) ~~(B)~~, both MRI and CT are would be approvable²⁸⁻³⁰ ~~(ACR, 2021; Koivikko, 2008; Taljanovic, 2009)~~
- When the patient is clinically unevaluable or there are preliminary imaging findings (x-ray or CT) needing further evaluation

("MRI and CT provide complementary information. When indicated It is appropriate to perform both examinations")²⁷ ~~(ACR, 2018)~~.

For evaluation of known fracture or known/new compression fractures with worsening back pain^{27, 31}

~~(ACR, 2018)~~

- To assess union of a fracture when physical examination, plain radiographs, or prior imaging suggest delayed or non-healing
- To determine the position of fracture fragments
- With history of malignancy (if MRI is contraindicated or cannot be performed)
- With an associated new focal neurologic deficit as above³² ~~(Alexandru, 2012)~~
- Prior to a planned surgery/intervention or if the results of the CT will change management

CT myelogram ~~is indicated when signs and symptoms are incongruent with MRI findings or:~~ When MRI cannot be performed/contraindicated/surgeon preference³³⁻³⁷

~~(Grams, 2010; Morita, 2011; Naganawa, 2011; NASS, 2012; Ozdoba, 2011)~~

- When signs and symptoms are inconsistent or not explained by the MRI findings
- Demonstration of the site of a CSF leak (known or suspected spontaneous (idiopathic) intracranial hypotension (SIH), post lumbar puncture headache, post spinal surgery headache, orthostatic headache, rhinorrhea or otorrhea, or cerebrospinal-venous fistula)
- Surgical planning, especially regarding to the nerve roots or evaluation of dural sac

For evaluation of tumor, cancer, or metastasis with any of the following:

(MRI is usually the preferred study- CT may be needed to further characterize solitary indeterminate lesions seen on MRI)³⁸

~~(Kim, 2012)~~

- **Primary tumor**

- Initial staging or re-staging of a known primary spinal tumor³⁹ ~~(NCCN 2021)~~
- Known spinal tumor with new signs or symptoms (e.g., new or increasing nontraumatic pain, physical, laboratory, and/or imaging findings)
- With an associated new focal neurologic deficit as above³² ~~(Alexandru, 2012)~~

- **Metastatic tumor**

- With evidence of metastasis on bone scan needing further clarification OR inconclusive findings on a prior imaging exam
- ~~Known malignancy with new signs or symptoms (e.g., new or increasing nontraumatic pain, physical, laboratory, and/or imaging findings) in a tumor that tends to metastasize to the spine~~
- With an associated new focal neurologic deficit³² ~~(Alexandru, 2012)~~

- Known malignancy with new signs or symptoms (e.g., new or increasing nontraumatic pain, radiculopathy or neck pain that occurs at night and wakes the patient from sleep with known active cancer, physical, laboratory, and/or imaging findings) in a tumor that tends to metastasize to the spine^{40, 41} ~~(ACR, 2018; Ziu, 2019)~~
- **For evaluation of inconclusive/indeterminate finding on prior imaging that requires further clarification**
 - One follow-up exam to ensure no suspicious change has occurred in prior imaging finding. No further surveillance unless specified as highly suspicious or change was found on last follow-up exam. When MRI cannot be performed or is contraindicated or CT is preferred to characterize the finding⁴¹ ~~(ACR, 2018)~~

Indication for combination studies for the initial pre-therapy staging of cancer, OR active monitoring for recurrence as clinically indicated, OR evaluation of suspected metastases

- ≤ 5 concurrent studies to include CT or MRI of any of the following areas as appropriate depending on the cancer: Neck, Abdomen, Pelvis, Chest, Brain, Cervical Spine, Thoracic Spine, or Lumbar Spine

For evaluation of known or suspected infection/-abscess when Thoracic MRI is contraindicated^{42, 43}

~~(ACR, 2018; Lerner, 2018)~~

- As evidenced by signs and/or symptoms, laboratory (i.e., abnormal white blood cell count, ESR and/or CRP) or prior imaging findings⁴⁴ ~~(Bond, 2016)~~
- Follow-up imaging of infection
 - With worsening symptoms/laboratory values (i.e., white blood cell count, ESR/CRP) or radiographic findings⁴⁵ ~~(Berbari, 2015)~~

For evaluation of known or suspected inflammatory disease when MRI is contraindicated or cannot be performed²⁸

~~(ACR, 2021)~~

- ~~For known or suspected~~ Ankylosing Spondylitis/Spondyloarthropathies with non-diagnostic or indeterminate x-ray and appropriate rheumatology workup

For evaluation of spine abnormalities related to immune system suppression, e.g., HIV, chemotherapy, leukemia, or lymphoma when Thoracic MRI is contraindicated⁴²

~~(ACR, 2018)~~

- As evidenced by signs/symptoms, laboratory, or prior imaging findings

Other Indications for a Thoracic Spine CT when MRI is contraindicated or cannot be performed

(Note- See [combination requests](#), below, for initial advanced imaging assessment and pre-operatively)

- Tethered cord, or spinal dysraphism (known or suspected) based on preliminary imaging, neurological exam, and/or high-risk cutaneous stigmata⁴⁶⁻⁴⁸ (~~AANS, 2019; Duz, 2008; Milhorat, 2009~~)
- Known Arnold-Chiari syndrome (For [initial imaging](#) see combination below)
 - Known Chiari I malformation without syrinx or hydrocephalus, follow-up imaging after initial diagnosis with new or changing signs/symptoms or exam findings consistent with spinal cord pathology⁴⁹ (~~Hitson, 2015~~)
 - Known Chiari II (Arnold-Chiari syndrome), III, or IV malformation
- Syrinx or syringomyelia (known or suspected)
 - With neurologic findings and/or predisposing conditions (e.g., Chiari malformation, prior trauma, neoplasm, arachnoiditis, severe spondylosis (~~Timpone, 2015~~))⁵⁰;
 - To further characterize a suspicious abnormality seen on prior imaging
 - Known syrinx with new/worsening symptoms
- [Toe walking in a child with signs/symptoms of myelopathy localized to the Thoracic Spine](#)
- [Suspected neuroinflammatory Conditions/Diseases \(e.g., sarcoidosis, Behcet's\)](#)
 - [After detailed neurological exam and basic testing completed](#)
- ~~Toe walking in a child when associated with upper motor neuron signs, including hyperreflexia, spasticity; or orthopedic deformity with concern for spinal cord pathology (e.g., pes cavus, clawed toes, leg or foot length deformity (excluding tight heel cords))~~

COMBINATION STUDIES WITH THORACIC SPINE CT WHEN MRI IS CONTRAINDICATED OR CANNOT BE PERFORMED OR SURGEON PREFERENCE

[Cervical and Thoracic CT](#)

- [Initial evaluation of known syrinx or syringomyelia](#)
 - [With neurologic findings and/or predisposing conditions \(e.g., Chiari malformation, prior trauma, neoplasm, arachnoiditis, severe spondylosis⁵⁰\)](#)
 - [To further characterize a suspicious abnormality seen on prior imaging](#)
 - [Known syrinx with new/worsening symptom](#)

~~Indications for combination studies^{51, 52}: (ACR, 2017, 2019) — For approved indications as noted below and being performed in a child under 8 years of age who will need anesthesia for the procedure~~

[Any combination of Cervical and/or Thoracic and/or Lumbar CTs](#)

Note: [These body regions might be evaluated separately or in combination as documented in the clinical notes by physical examination findings \(e.g., localization to a particular segment of the spinal cord\), patient history, and other available information, including prior imaging.](#)

[Exception- Indications for combination studies^{51, 52}: Are approved indications as noted below and being performed in children who will need anesthesia for the procedure](#)

- Any combination of these studies for:
 - Survey/complete initial assessment **of** infant/child with congenital scoliosis or juvenile idiopathic scoliosis under the age of 10⁵³⁻⁵⁵ (e.g., congenital scoliosis, idiopathic scoliosis, scoliosis with vertebral anomalies)
 - In the presence of neurological deficit, progressive spinal deformity, or for preoperative planning⁵⁶
 - Back pain with known vertebral anomalies (hemivertebrae, hypoplasia, agenesis, butterfly, segmentation defect, bars, or congenital wedging) in a child on preliminary imaging
 - Scoliosis with any of the following⁵⁷:
 - Progressive spinal deformity;
 - Neurologic deficit (new or unexplained);
 - Early onset;
 - Atypical curve (e.g., short segment, >30° kyphosis, left thoracic curve, associated organ anomalies);
 - Pre-operative planning; OR
 - When office notes clearly document how imaging will change management
- Arnold-Chiari malformations^{58, 59}
 - Arnold-Chiari I
 - For evaluation of spinal abnormalities associated with initial diagnosis of Arnold-Chiari Malformation. (C/T/L spine due to association with tethered cord and syringomyelia), and initial imaging has not been completed^{47, 53}
 - Arnold-Chiari II-IV - For initial evaluation and follow-up as appropriate
 - Usually associated with open and closed spinal dysraphism, particularly meningocele
- Tethered cord, or spinal dysraphism (known or suspected) based on preliminary imaging, neurological exam, and/or high-risk cutaneous stigmata,⁴⁶⁻⁴⁸ ~~40-42~~ when anesthesia required for imaging⁶⁰ ~~54~~ (e.g., meningocele, lipomenocele, diastematomyelia, fatty/thickened filum terminale, and other spinal cord malformations)
- Oncological Applications (e.g., primary nervous system, metastatic)
 - Drop metastasis from brain or spine (imaging also includes brain; CT spine imaging in this scenario is usually CT myelogram)- See **Background Overview**
 - Suspected leptomeningeal carcinomatosis (LC)⁶¹- See **Overview**
 - Any combination of these for spinal survey in patient with metastases
 - Tumor evaluation and monitoring in neurocutaneous syndromes—See **Background**
- CSF leak highly suspected and supported by patient history and/or physical exam findings (leak (known or suspected spontaneous (idiopathic) intracranial hypotension (SIH), post lumbar puncture headache, post spinal surgery headache, orthostatic headache, rhinorrhea or otorrhea, or cerebrospinal-venous fistula -preferred exam CT myelogram))¹⁸
- CT myelogram when meets above guidelines and MRI is contraindicated or for surgical planning
- Post-procedure (discogram) CT

Any combination of Cervical and/or Thoracic and/or Lumbar CTs

- ~~Any combination of these studies for:~~
 - ~~Scoliosis survey in infant/child with congenital scoliosis or juvenile idiopathic scoliosis under the age of 10⁴⁸⁻⁵⁰ (ACR, 2018; SRS, 2019; Strahle, 2015)~~
 - ~~In the presence of neurological deficit, progressive spinal deformity, or for preoperative planning⁵¹ (Trenga, 2016)~~
 - ~~Back pain and vertebral anomalies (hemivertebrae, hypoplasia, agenesis, butterfly, segmentation defect, bars, or congenital wedging) in a child on preliminary imaging~~
 - ~~Scoliosis with any of the following⁵² (Ozturk, 2010):~~
 - ~~Progressive spinal deformity;~~
 - ~~Neurologic deficit;~~
 - ~~Early onset;~~
 - ~~Atypical curve (e.g., short segment, >30° kyphosis, left thoracic curve, associated organ anomalies);~~
 - ~~Pre-operative planning; OR~~
 - ~~When office notes clearly document how imaging will change management~~
- ~~Arnold-Chiari I^{53,54} (Radic, 2018; Strahle, 2011)~~
 - ~~For evaluation of spinal abnormalities associated with initial diagnosis of Arnold-Chiari Malformation. (C/T/L spine due to association with tethered cord and syringomyelia), and initial imaging has not been completed^{42,49} (Milhorat, 2009; Strahle, 2015)~~
- ~~Arnold-Chiari II-IV~~
 - ~~For initial evaluation and follow-up as appropriate~~
- ~~Tethered cord, or spinal dysraphism (known or suspected) based on preliminary imaging, neurological exam, and/or high-risk cutaneous stigmata,⁴¹⁻⁴³ (AANS, 2019; Duz, 2008; Milhorat, 2009), when anesthesia required for imaging⁵⁵ (Hertzler, 2010)~~
- ~~Toe walking in a child when associated with upper motor neuron signs including hyperreflexia, spasticity; or orthopedic deformity with concern for spinal cord pathology (e.g., pes cavus, clawed toes, leg or foot length deformity (excluding tight heel cords))~~
- ~~Back pain in a child with any of the following red flags (conservative care not required when red flags present):~~
 - ~~Red flags that prompt imaging should include the presence of: age 5 or younger, constant pain, pain lasting >4 weeks, abnormal neurologic examination, early morning stiffness and/or gelling; night pain that prevents or disrupts sleep; radicular pain; fever; weight loss; malaise; postural changes (e.g., kyphosis or scoliosis); and limp (or refusal to walk in a younger child <5yo), AND initial radiographs have been performed^{41,42} (Bernstein, 2007; Feldman, 2006)~~
- ~~Drop metastasis from brain or spine (imaging also includes brain; CT spine imaging in this scenario is usually CT myelogram)~~
 - ~~Suspected leptomeningeal carcinomatosis (LC)⁵⁶ (Shah, 2011)~~
 - ~~Any combination of these for spinal survey in patient with metastases.~~

- ~~Tumor evaluation and monitoring in neurocutaneous syndromes—See Background~~
- ~~CSF leak highly suspected and supported by patient history and/or physical exam findings (leak (known or suspected spontaneous (idiopathic) intracranial hypotension (SIH), post lumbar puncture headache, post spinal surgery headache, orthostatic headache, rhinorrhea or otorrhea, or cerebrospinal venous fistula—preferred exam CT myelogram)⁴⁵ (Starling, 2013)~~
- ~~CT myelogram when meets above guidelines and MRI is contraindicated or for surgical planning~~
- ~~Post procedure (discogram) CT~~

BACKGROUND

Computed tomography is used for the evaluation, assessment of severity, and follow-up of diseases of the spine. Its use in the thoracic spine is limited, however, due to the lack of epidural fat in this part of the body. CT myelography improves the contrast severity of CT, but it is also invasive. CT may be used for conditions, e.g., degenerative changes, infection, and immune suppression, when magnetic resonance imaging (MRI) is contraindicated. It may also be used in the evaluation of tumors, cancer, or metastasis in the thoracic spine, and it may be used for preoperative and post-surgical evaluations. CT obtains images from different angles and uses computer processing to show a cross-section of body tissues and organs. CT is fast and is often performed in acute settings. It provides good visualization of cortical bone.

OVERVIEW

Ankylosing Spondylitis/Spondyloarthropathies can cause back or sacroiliac pain of insidious onset (usually > 3 month), associated with morning stiffness not relieved with rest (usually age at onset <40). They are associated with any of the following⁶²⁻⁶⁵ ~~(Akgul, 2011; Bennett, 2010; Østergaard, 2012; Sieper, 2014):~~

- Sedimentation rate and/or C-reactive protein (not an essential criteria)
- HLA B27 (not an essential criteria)
- Non-diagnostic or indeterminate x-ray
- Personal or family history of ~~sacroilitis~~**sacroiliitis**, peripheral inflammatory arthritis, and/or inflammatory bowel disease.

***Conservative Therapy** ~~÷~~ — (Spine) should include a multimodality approach consisting of a **combination of active and inactive components**. Inactive components, such as rest, ice, heat, modified activities, medical devices, acupuncture and/or stimulators, medications, injections (epidural, facet, bursal, and/or joint, not including trigger point), and diathermy can be utilized. Active modalities may consist of physical therapy, a physician-supervised home exercise program**, regular osteopathic manipulative medicine treatments (OMT), and/or chiropractic care when considered safe and appropriate.

****Home Exercise Program - (HEP)/Therapy** – the following elements are required to meet guidelines for completion of conservative therapy^{10, 17} ~~(ACR, 2015; Last, 2009)~~:

- Information provided on exercise prescription/plan AND
- Follow-up with member with documentation provided regarding lack of improvement (failed) after completion of HEP (after suitable 6-week period), or inability to complete HEP due to physical reason- i.e., increased pain, inability to physically perform exercises. (Patient inconvenience or noncompliance without explanation does not constitute “inability to complete” HEP).
- Dates and duration of failed PT, physician-supervised HEP, or chiropractic treatment should be documented in the original office notes or an addendum to the notes.

Table 1: Gait and spine imaging⁶⁶⁻⁷¹

Gait	Characteristic	Work up/Imaging
Hemiparetic	Spastic unilateral, circumduction	Brain and/or, Cervical spine imaging based on associated symptoms
Diplegic	Spastic bilateral, circumduction	Brain, Cervical and Thoracic Spine imaging
Myelopathic	Wide based, stiff, unsteady	Cervical and/or Thoracic spine MRI based on associated symptoms
Ataxic	Broad based, clumsy, staggering, lack of coordination, usually also with limb ataxia	Brain imaging
Apraxic	Magnetic, shuffling, difficulty initiating	Brain imaging
Parkinsonian	Stooped, small steps, rigid, turning en bloc, decreased arm swing	Brain Imaging
Choreiform	Irregular, jerky, involuntary movements	Medication review, consider brain imaging as per movement disorder Brain MR guidelines
Sensory ataxic	Cautious, stomping, worsening without visual input (ie + Romberg)	EMG, blood work, consider spinal (cervical or thoracic cord imaging) imaging based on EMG
Neurogenic	Steppage, dragging of toes	EMG foot drop Lumbar spine MRI <ul style="list-style-type: none"> • <u>Pelvis MR appropriate evidence of plexopathy-EMG initial testing;</u> • <u>BUT if there is a foot drop, lumbar spine MRI is appropriate without EMG</u> • <u>Pelvis MR if there is evidence of plexopathy</u>

Vestibular	Insecure, veer to one side, worse when eyes closed, vertigo	Consider Brain/IAC MRI as per GL
------------	---	----------------------------------

([#]References: Chhetri, 2014; Clinch, 2021; Gait, 2021; Haynes, 2018; Marshall, 2012; Pirker, 2017)

Myelopathy – Symptom severity varies, and a high index of suspicion is essential for making the proper diagnosis in early cases. Symptoms of pain and radiculopathy may not be present. The natural history of myelopathy is characterized by neurological deterioration. The most frequently encountered symptom is gait abnormality (86%), followed by increased muscular reflexes (79.1%), pathological reflexes (65.1%), paresthesia of upper limb (69.8%), and pain (67.4%) (Vitzthum, 2007).⁷²

CT and Infection of the spine – Infection of the spine is not easy to differentiate from other spinal disorders, e.g., degenerative disease, spinal neoplasms, and non-infective inflammatory lesions. Infections may affect different parts of the spine, e.g., vertebrae, intervertebral discs, and paraspinal tissues. Imaging is important to obtain early diagnosis and treatment to avoid permanent neurology deficits. When MRI is contraindicated, CT may be used to evaluate infections of the spine.

CT and Degenerative Disc Disease – Degenerative disc disease is very common, and CT may be indicated when MRI is contraindicated and when chronic degenerative changes are accompanied by conditions, e.g., new neurological deficits; onset of joint tenderness of a localized area of the spine; new abnormal nerve conduction studies; exacerbation of chronic back pain unresponsive to conservative treatment; and unsuccessful physical therapy/home exercise program.

Infection, Abscess, or Inflammatory disease

- Most common site is the lumbar spine (58%), followed by the thoracic spine (30%) and the cervical spine (11%) (Graeber, 2019)
- High risk populations (indwelling hardware, history of endocarditis, IVDA, recent procedures) with appropriate signs/symptoms

CT Myelogram

Myelography is the instillation of intrathecal contrast media under fluoroscopy. Patients are then imaged with CT to evaluate for spinal canal pathology. Although this technique has diminished greatly due to the advent of MRI and its non-invasiveness and superior soft-tissue contrast, myelography is still a useful technique for conventional indications, such as spinal stenosis, when MRI is contraindicated or nondiagnostic, brachial plexus injury in neonates, radiation therapy treatment planning, and cerebrospinal fluid (CSF) leak (Pomerantz, 2016).⁷³

Cauda Equina Syndrome

- Symptoms include severe back pain or sciatica along with one or more of the following:

- Saddle anesthesia - loss of sensation restricted to the area of the buttocks, perineum, and inner surfaces of the thighs (areas that would sit on a saddle)
- Recent bladder/bowel dysfunction
- Achilles reflex absent on both sides
- Sexual dysfunction that can come on suddenly
- Absent anal reflex and bulbocavernosus reflex

Back Pain with Cancer History — Radiographic (x-ray) examination should be performed in cases of back pain when a patient has a cancer history, but without known active cancer or a tumor that tends to metastasize to the spine. This can make a diagnosis in many cases. This may occasionally allow for selection of bone scan in lieu of MRI in some cases. When radiographs do not answer the clinical question, then MRI may be appropriate after a consideration of conservative care.

“Neoplasms causing VCF (vertebral compression fractures) include: primary bone neoplasms, such as hemangioma or giant cell tumors, and tumor-like conditions causing bony and cellular remodeling, such as aneurysmal bone cysts, or Paget’s disease (osteitis deformans); infiltrative neoplasms, including and not limited to, multiple myeloma and lymphoma, and metastatic neoplasms (ACR, 2018).”³¹

Most common spine metastasis involving primary metastasis originate from the following tumors in descending order: breast (21%), lung (19%), prostate (7.5%), renal (5%), gastrointestinal (4.5%), and thyroid (2.5%). While all tumors can seed to the spine, the cancers mentioned above metastasize to the spinal column early in the disease process (Ziu, 2019).⁴⁰

Drop Metastases⁷⁴ —

Drop metastases are intradural extramedullary spinal metastases that arise from intracranial lesions. Common examples of intracranial neoplasms that result in drop metastases include pineal tumors, ependymomas, medulloblastomas, germinomas, primitive neuroectodermal tumors (PNET), glioblastomas multiform, anaplastic astrocytomas, oligodendrogliomas and less commonly choroid plexus neoplasms and teratomas.

Leptomeningeal Carcinomatosis⁷⁵ —

Leptomeningeal carcinomatosis is a complication of cancer in which cancerous cells spread to the membranes (meninges) that covers the brain and spinal cord. The most common solid tumors that involve the leptomeninges are breast, lung, and melanoma, gastrointestinal, and primary central nervous system tumors.

Table 2: MRI and Cutaneous Stigmata⁷⁶ (Dias, 2015)

Risk Stratification for Various Cutaneous Markers

High Risk	Intermediate Risk	Low Risk
<ul style="list-style-type: none"> • Hypertrichosis • Infantile hemangioma • Anorectal meningocele • DST • Subcutaneous lipoma • Caudal appendage • Segmental hemangiomas in association with LUMBAR[‡] syndrome 	<ul style="list-style-type: none"> • Capillary malformations (also referred to as NFS or salmon patch when pink and poorly defined or PWS when darker red and well-defined) 	<ul style="list-style-type: none"> • Coccygeal dimple • Light hair • Isolated café au lait spots • Mongolian spots • Hypo- and hypermelanotic macules or papules • Deviated or forked gluteal cleft • Nonmidline lesions
[‡] LUMBAR, lower body hemangioma and other cutaneous defects, urogenital abnormalities, ulcerations, myelopathy, bony defects, anorectal malformations, arterial anomalies, and renal anomalies.		

POLICY HISTORY

Date	Summary
<u>March 2022</u>	<p><u>Added</u></p> <ul style="list-style-type: none"> • <u>Combination request for overlapping body part statement</u> • <u>Clarified muscle weakness not related to plexopathy or peripheral neuropathy</u> • <u>Clarified bowel and bladder dysfunction – not related to an inherent bowel or bladder problem</u> • <u>Added subsection for cervical and thoracic spine section for syrinx and syringomyelia</u> • <u>Descriptions for tethered cord</u> • <u>Clarified CT myelogram section</u> • <u>Background section of Drop Metastases</u> • <u>Background section of Leptomeningeal Carcinomatosis</u> • <u>Clarified toe walking in pediatric patient with myelopathy for thoracic spine</u> <p><u>Removed</u></p> <ul style="list-style-type: none"> • <u>Removed from combination section syrinx and syringomyelia and added subsection for cervical and thoracic spine section</u> • <u>Removed pediatric back pain from the total spine combination section</u>
April 2021	<ul style="list-style-type: none"> • Added/modified <ul style="list-style-type: none"> ○ Modified section on neurological deficits ○ Back pain in a child added/modified red flags ○ Gait table in background

	<ul style="list-style-type: none"> ○ Post-surgical modified/clarified surgical criteria for combination exams and surgeon preference for exam type ○ Removed myelopathy combination studies ○ Updated/added MS Criteria <ul style="list-style-type: none"> ▪ Combination section for initial imaging and follow up ▪ Added pediatric MS ○ Modified known tumor imaging into primary and metastatic disease ○ Added toe walking for pediatric patients ○ Modified Combination exam wording
May 2020	<ul style="list-style-type: none"> ● For evaluation of neurologic deficits when new deficits are present ● Removed pars defect section ● Added ankylosing spondylitis for evaluation of trauma/acute injury ● Modified Initial imaging of new or increasing non-traumatic back pain or radiculopathy or back pain that occurs at night and wakes the patient from sleep with known active cancer and a tumor that tends to metastasize to the spine ● Added Imaging of Ossification of the Posterior Longitudinal Ligament (OPPL) ● Added Osteopathic Manipulative medicine to conservative care therapy
June 2019	<ul style="list-style-type: none"> ● Added: <ul style="list-style-type: none"> ○ new or worsening objective neuro deficits for chronic and acute back pain; CSF leak ○ last 6 months for allowable post op f/u period and removed EMG comment ○ red flags specifically for peds back pain and pain related to malignancy, infection, inflammation ○ new sections: pars defect; compression fractures; congenital abnormalities including section on scoliosis and vertebral anomalies in children w/back pain; ○ For combination studies cervical/thoracic/lumbar added drop metastasis, tumor evaluation for neurocutaneous syndromes, and abnormalities associated w/Arnold Chiari, as well as separate indication for tethered cord or spinal dysraphism ○ Spinal cord stimulator

	<ul style="list-style-type: none">○ New or increasing back pain in cancer patients with high suspicion of mets
--	--

REFERENCES

- Acharya AB, Fowler JB. Chaddock Reflex. [Updated 2019 Dec 15]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan. <https://www.ncbi.nlm.nih.gov/books/NBK519555/>.
- Akgul O, Ozgoemen S. Classification criteria for spondyloarthropathies. *World J Orthop*. Dec 18; 2011; 2(12):107-115. doi: 10.5312/wjo.v2.i12.07. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3302034/>. Retrieved March 29, 2018.
- Alexandru D. Evaluation and management of vertebral compression fractures. *Perm J*. Published online October 30, 2012:46-51. doi:10.7812/TPP/12-037.
- Allegri M, Montella S, Salici F, et al. Mechanisms of low back pain: A guide for diagnosis and therapy. *F1000Res*. 2016 Oct; 2.
- American Association of Neurological Surgeons (AANS). Tethered spinal cord syndrome. 2019.
- American Association of Neurological Surgeons and Congress of Neurological Surgeons (AANSCNS). Choosing Wisely®. <http://www.choosingwisely.org/clinician-lists/american-association-neurological-surgeons-imaging-for-nonspecific-acute-low-back-pain/>. Released June 24, 2014.
- American College of Radiology (ACR). ACR Appropriateness Criteria®. Published 2015.
- American College of Radiology (ACR). ACR Appropriateness Criteria®. Inflammatory Back Pain: Known or Suspected Axial Spondyloarthritis. <https://acsearch.acr.org/docs/3094107/Narrative/>. Revised 2021.
- Ando K, Imagama S, Kobayashi K, et al. Clinical features of thoracic myelopathy: A single-center study. *J Am Acad Orthop Surg Glob Res Rev*. 2019 Nov; 3(11):e10.5435.
- Behrbalk E, Salame K, Regev GJ, et al. Delayed diagnosis of cervical spondylotic myelopathy by primary care physicians. *Neurosurg Focus*. July 2013; 35(1):1-6.
- Bennett AN, Marzo-Ortega H, Rehman A, et al. The evidence for whole spine MRI in the assessment of axial spondyloarthropathy. *Rheumatology*. March 2010; 49(3):426-432. <https://doi.org/10.1093/rheumatology/kep427>. <https://academic.oup.com/rheumatology/article/49/3/426/1787410>. Retrieved May 12, 2018.
- Berberi EF, Kanj SS, Kowalski TJ, et al. 2015 Infectious Diseases Society of America (IDSA) Clinical Practice Guidelines for the Diagnosis and Treatment of Native Vertebral Osteomyelitis in Adults. *Clin Infect Dis*. 2015 Sep 15; 61(6):e26-e46.

Bernstein RM, Cozen H. Evaluation of back pain in children and adolescents. *Am Fam Physician*. 2007;76(11):1669–1676.

Bond A, Manian FA. Spinal epidural abscess: A review with special emphasis on earlier diagnosis. *Biomed Res Int*. 2016; 2016:1614328.

Budoff MJ, Khairallah W, Li D, et al. Trabecular bone mineral density measurement using thoracic and lumbar quantitative computed tomography. *Acra Radiology*. 2012; 19(2):179–183. doi: 10.1016/j.acra.2011.10.006.

Chhetri SK, Gow D, Shaunak S, Varma A. Clinical assessment of the sensory ataxias; diagnostic algorithm with illustrative cases. *Pract Neurol*. 2014;14(4):242–251. doi:10.1136/practneurol-2013-000764.

Choi BW, Song KJ, Chang H. Ossification of the posterior longitudinal ligament: A review of literature. *Asian Spine J*. 2011; 5(4):267–276. doi:10.4184/asj.2011.5.4.267

Choosing Wisely®. Lists. <http://www.choosingwisely.org/clinician-lists/#keyword=spine&topic-area=Radiology&service=Imaging>. Retrieved March 27, 2018.

Clinch J, Wood M, Driscoll S. Evaluation of gait disorders in children. *BMJ Best Practice*. Published February 23, 2021. Accessed July 14, 2021. <https://bestpractice.bmj.com/topics/en-us/709>.

D’Alessandro. Does This Sacral Dimple Need to be Evaluated? *PediatricEducation.org™*. <https://pediatriceducation.org/2009/07/20/does-this-sacral-dimple-need-to-be-evaluated/>. Posted July 20, 2009. Retrieved March 29, 2018.

Davies BM, Mowforth OD, Smith EK, et al. Degenerative cervical myelopathy. *BMJ*. 2018; 360. doi: <https://doi.org/10.1136/bmj.k186>.

Dias M, Partington M. Congenital brain and spinal cord malformations and their associated cutaneous markers. *Pediatrics*. 2015; 136(4):e1105–19.

Duz B, Gocmen S, Secer HI, et al. Tethered cord syndrome in adulthood. *J Spinal Cord Med*. 2008; 31(3):272–278. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2565560/>. Retrieved March 29, 2018.

Eubanks JD. Cervical radiculopathy: Nonoperative management of neck pain and radicular symptoms. *Am Fam Physician*. 2010 Jan 1; 81(1):33–40. <https://www.aafp.org/afp/2010/0101/p33.pdf>.

Feldman DS, Straight JJ, Badra MI, Mohaideen A, Madan SS. Evaluation of an algorithmic approach to pediatric back pain. *J Pediatr Orthop*. 2006;26(3):353-357. doi:10.1097/01.bpo.0000214928.25809.f9.

Fisher BM, Cowles S, Matulich JR, Evanson BG, Vega D, Dissanaik S. Is magnetic resonance imaging in addition to a computed tomographic scan necessary to identify clinically significant cervical spine injuries in obtunded blunt trauma patients? *Am J Surg*. 2013;206(6):987-993; discussion 993-994. doi:10.1016/j.amjsurg.2013.08.021.

Gait abnormalities. Stanford Medicine 25. Published 2021. Accessed July 14, 2021. <https://stanfordmedicine25.stanford.edu/the25/gait.html>.

Graeber A, Cecava ND. Vertebral Osteomyelitis. [Updated 2019 Jun 3]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan.

Haynes KB, Wimberly RL, VanPelt JM, Jo C-H, Riccio AI, Delgado MR. Toe walking: A neurological perspective after referral from pediatric orthopaedic surgeons. *Journal of Pediatric Orthopaedics*. 2018;38(3):152-156. doi:10.1097/BPO.0000000000001115.

Hertzler DA, DePowell JJ, Stevenson CB, Mangano FT. Tethered cord syndrome: A review of the literature from embryology to adult presentation. *Neurosurg Focus*. 2010;29(1):E1. doi:10.3171/2010.3.FOCUS1079.

Hitson WJ, Lane JR, Bauer DF, et al. A prospective natural history study of nonoperatively managed Chiari I malformation: Does follow-up MRI surveillance alter surgical decision making? *J Neurosurg Pediatr*. 2015 Aug; 16(2):159-66.

Hou X, Sun C, Liu X, et al. Clinical features of thoracic spinal stenosis-associated myelopathy: A retrospective analysis of 427 cases. *Clin Spine Surg*. 2016 Mar; 29(2):86-9. doi: 10.1097/BSD.0000000000000081.

Jarvik JG, Gold LS, Comstock BA, et al. Association of early imaging for back pain with clinical outcomes in older adults. *JAMA*. 2015; 313(11):1143-1153. doi: 10.1001/jama.2015.1871.

Kim YS, Han IH, Lee IS, et al. Imaging findings of solitary spinal bony lesions and the differential diagnosis of benign and malignant lesions. *J Korean Neurosurg Soc*. August 2012; 52(2):126-132. doi: 10.3340/jkns.2012.52.2. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3467370/>. Retrieved March 29, 2018.

Koivikko MP, Koskinen SK. MRI of cervical spine injuries complicating ankylosing spondylitis. *Skeletal Radiol*. 2008 Sep; 37(9):813-9. doi:10.1007/s00256-008-0484-x. Epub 2008 Apr 18.

Last AR, Hulbert K. Chronic low back pain: Evaluation and management. *Am Fam Physician*. 2009; 79(12):1067-74.

Lerner S, Hartmann S, Barbagallo GMV, et al. Management of spinal infection: A review of the literature. *Acta Neurochir (Wien)*. 2018; 160(3):487–496. doi: 10.1007/s00701-018-3467-2.

Marshall FJ. Approach to the elderly patient with gait disturbance. *Neurol Clin Pract*. 2012;2(2):103–111. doi:10.1212/CPJ.0b013e31825a7823.

Milhorat TH, Bolognese PA, Nishikawa M, et al. Association of Chiari malformation type I and tethered cord syndrome: Preliminary results of sectioning filum terminale. *Surg Neurol*. July 1, 2009; 72(1):20–35. <http://europepmc.org/abstract/med/19559924>. Retrieved March 29, 2018.

Muller D, Bauer JS, Zeile M, et al. Significance of sagittal reformations in routine thoracic and abdominal multislice CT studies for detecting osteoporotic fractures and other spine abnormalities. *Eur Radiol*. 2008; 18(8):1696–1702. doi: 10.1007/s00330-008-0920-2.

North American Spine Society (NASS). Clinician Lists. Choosing Wisely®. <http://www.choosingwisely.org/clinician-lists/nass-emg-nerve-conduction-studies-to-determine-cause-of-spine-pain/>. Released October 9, 2013.

North American Spine Society (NASS). Evidence-based Clinical Guidelines for Multidisciplinary Spine Care. Diagnosis and Treatment of Cervical Radiculopathy from Degenerative Disorders. 2010 North American Spine Society. <https://www.spine.org/Portals/0/Assets/Downloads/ResearchClinicalCare/Guidelines/CervicalRadiculopathy.pdf>

North American Spine Society (NASS). Five Things Physicians and Patients Should Question. Choosing Wisely®. <http://www.choosingwisely.org/doctor-patient-lists/north-american-spine-society/>. 2014.

Ostergaard M, Lambert RG. Imaging in ankylosing spondylitis. *Ther Adv Musculoskelet Dis*. August 2012; 4(4):301–311. doi: 10.1177/1759720X11436240. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3403247/>. Retrieved March 29, 2018.

Ozturk C, Karadereler S, Ornek I, Enercan M, Ganiyusufoglu K, Hamzaoglu A. The role of routine magnetic resonance imaging in the preoperative evaluation of adolescent idiopathic scoliosis. *Int Orthop*. 2010;34(4):543–546. doi:10.1007/s00264-009-0817-y.

Pirker W, Katzenschlager R. Gait disorders in adults and the elderly: A clinical guide. *Wien Klin Wochenschr*. 2017;129(3–4):81–95. doi:10.1007/s00508-016-1096-4.

Pomerantz SR. Myelography: Modern technique and indications. *Handb Clin Neurol*. 2016; 135:193–208.

~~Radic JAE, Cochrane DD. Choosing wisely canada: pediatric neurosurgery recommendations. *Paediatr Child Health*. 2018;23(6):383-387. doi:10.1093/pch/pxy012.~~

~~Rao D, Scuderi G, Scuderi C, Grewal R, et al. The use of imaging in management of patients with low back pain. *J Clin Imaging Sci*. 2018 Aug 24; 8:30.~~

~~Scoliosis Research Society (SRS). Conditions and treatments: Juvenile scoliosis. 2019.~~

~~Shah LM, Salzman KL. Imaging of spinal metastatic disease. *Int J Surg Oncol*. 2011; 2011:769753.~~

~~Sieper J, Rudwaleit M, Baraliakos X, et al. The assessment of SpondyloArthritis international Society (ASAS) handbook: A guide to assess spondyloarthritis. *Ann Rheum Dis*. June 2009; 68 Suppl 2:ii1-44.~~

~~Starling A, Hernandez F, Hoxworth JM, et al. Sensitivity of MRI of the spine compared with CT myelography in orthostatic headache with CSF leak. *Neurology*. 2013;81(20):1789-1792. doi:10.1212/01.wnl.0000435555.13695.22.~~

~~Stolper K, Haug JC, Christensen CT, et al. Prevalence of thoracic spine lesions masquerading as cauda equina syndrome: yield of a novel magnetic resonance imaging protocol. *Intern Emerg Med*. 2017 Dec; 12(8):1259-1264. doi: 10.1007/s11739-016-1565-9.~~

~~Strahle J, Muraszko KM, Kapurch J, Bapuraj JR, Garton HJL, Maher CO. Chiari malformation Type I and syrinx in children undergoing magnetic resonance imaging. *J Neurosurg-Pediatr*. 2011;8(2):205-213. doi:10.3171/2011.5.PEDS1121.~~

~~Strahle J, Smith BW, Martinez M, et al. The association between Chiari malformation Type I, spinal syrinx, and scoliosis. *J Neurosurg-Pediatr*. 2015; 15:607-611. <http://thejns.org/doi/pdf/10.3171/2014.11.PEDS14135>. Retrieved March 29, 2018.~~

~~Taljanovic MS, Hunter TB, Wisneski RJ, et al. Imaging characteristics of diffuse idiopathic skeletal hyperostosis with an emphasis on acute spinal fractures: Review. *AJR Am J Roentgenol*. 2009;193(3 Suppl):S10-19, Quiz S20-24. doi:10.2214/AJR.07.7102.~~

~~Teoli D, Rocha-Cabrero F, Ghassemzadeh S. Lhermitte Sign. In: *StatPearls*. StatPearls Publishing; 2021. Accessed July 15, 2021. <http://www.ncbi.nlm.nih.gov/books/NBK493237/>~~

~~Timpone V, Patel SH. MRI of a syrinx: Is contrast material always necessary? *Am J Roentgenol*. 2015; 204:1082-1085. 10.2214/AJR.14.13310.~~

~~Trenga AP, Singla A, Feger MA, et al. Patterns of congenital bony spinal deformity and associated neural anomalies on X-ray and magnetic resonance imaging. *J Child Orthop*. August 2016; 10(4):343-352. doi: 10.1007/s11832-016-0752-6. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4940247/>. Retrieved March 29, 2018.~~

~~Vilaca C, Orsini M, Leite MAA, et al. Cervical spondylotic myelopathy: What the neurologist should know. *Neurol Int.* 2016 Nov 2; 8(4):6330. doi: 10.4081/ni.2016.6330.~~

~~Vitzthum H, Dalitz K. Analysis of five specific scores of cervical spondylogenic myelopathy. *Eur Spine J.* December 2007; 16(12):2096-2103.
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2140133/>.~~

~~Waly FJ, Abduljabbar FH, Fortin M, et al. Preoperative computed tomography myelography parameters as predictors of outcome in patients with degenerative cervical myelopathy: Results of a systematic review. *Global Spine J.* 2017; 7(6):521-528. doi:10.1177/2192568217701101.~~

~~Ziu E, Viswanathan VK, Mesfin FB. Cancer, Spinal Metastasis. [Updated 2020 Mar 30]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK441950/>.~~

Reviewed / Approved by NIA Clinical Guideline Committee

GENERAL INFORMATION

~~It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.~~

Disclaimer: Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.

1. Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis and Treatment of Cervical Radiculopathy from Degenerative Disorders. North American Spine Society (NASS). Updated 2010. Accessed November 9, 2021. <https://www.spine.org/Portals/0/Assets/Downloads/ResearchClinicalCare/Guidelines/CervicalRadiculopathy.pdf>
2. Acharya AB, Fowler JB. Chaddock Reflex. StatPearls Publishing. Updated August 1, 2021. Accessed November 9, 2021. <https://www.ncbi.nlm.nih.gov/books/NBK519555/>
3. Albert TJ, Murrell SE. Surgical management of cervical radiculopathy. *J Am Acad Orthop Surg*. Nov-Dec 1999;7(6):368-76. doi:10.5435/00124635-199911000-00003
4. Moore KR, Tsuruda JS, Dailey AT. The value of MR neurography for evaluating extraspinal neuropathic leg pain: a pictorial essay. *AJNR Am J Neuroradiol*. Apr 2001;22(4):786-94.
5. Dydyk AM, Hameed S. Lumbosacral Plexopathy. StatPearls Publishing Copyright © 2022, StatPearls Publishing LLC. Updated November 7, 2021. Accessed February 25, 2022. <https://www.ncbi.nlm.nih.gov/books/NBK556030/>
6. Teoli D, Rocha Cabrero F, Ghassemzadeh S. Lhermitte Sign. StatPearls Publishing. Updated June 29, 2021. Accessed November 9, 2021. <https://www.ncbi.nlm.nih.gov/books/NBK493237/>
7. Allegri M, Montella S, Salici F, et al. Mechanisms of low back pain: a guide for diagnosis and therapy. *F1000Res*. 2016;5doi:10.12688/f1000research.8105.2
8. American Association of Neurological Surgeons, Congress of Neurological Surgeons. Five things physicians and patients should question: Don’t obtain imaging (plain radiographs,

magnetic resonance imaging, computed tomography [CT], or other advanced imaging) of the spine in patients with non-specific acute low back pain and without red flags. Choosing Wisely Initiative ABIM Foundation. Updated 2020. Accessed November 9, 2021.

<https://www.choosingwisely.org/clinician-lists/american-association-neurological-surgeons-imaging-for-nonspecific-acute-low-back-pain/>

9. Jarvik JG, Gold LS, Comstock BA, et al. Association of early imaging for back pain with clinical outcomes in older adults. *Jama*. Mar 17 2015;313(11):1143-53. doi:10.1001/jama.2015.1871

10. Last AR, Hulbert K. Chronic low back pain: evaluation and management. *Am Fam Physician*. Jun 15 2009;79(12):1067-74.

11. Eubanks JD. Cervical radiculopathy: nonoperative management of neck pain and radicular symptoms. *Am Fam Physician*. Jan 1 2010;81(1):33-40.

12. North American Spine Society. Five things physicians and patients should question: Don't use electromyography (EMG) and nerve conduction studies (NCS) to determine the cause of axial lumbar, thoracic or cervical spine pain. Choosing Wisely Initiative ABIM Foundation. Updated 2021. Accessed November 9, 2021.

<https://www.choosingwisely.org/clinician-lists/nass-emg-nerve-conduction-studies-to-determine-cause-of-spine-pain/>

13. American College of Radiology. ACR Appropriateness Criteria® Back Pain—Child. American College of Radiology (ACR). Updated 2016. Accessed November 10, 2021.

<https://acsearch.acr.org/docs/3099011/Narrative/>

14. Bernstein RM, Cozen H. Evaluation of back pain in children and adolescents. *Am Fam Physician*. Dec 1 2007;76(11):1669-76.

15. Feldman DS, Straight JJ, Badra MI, Mohaideen A, Madan SS. Evaluation of an algorithmic approach to pediatric back pain. *J Pediatr Orthop*. May-Jun 2006;26(3):353-7. doi:10.1097/01.bpo.0000214928.25809.f9

16. Rao D, Scuderi G, Scuderi C, Grewal R, Sandhu SJ. The Use of Imaging in Management of Patients with Low Back Pain. *J Clin Imaging Sci*. 2018;8:30. doi:10.4103/jcis.JCIS_16_18

17. American College of Radiology. ACR Appropriateness Criteria® Low Back Pain. American College of Radiology (ACR). Updated 2021. Accessed November 10, 2021.

<https://acsearch.acr.org/docs/69483/Narrative/>

18. Starling A, Hernandez F, Hoxworth JM, et al. Sensitivity of MRI of the spine compared with CT myelography in orthostatic headache with CSF leak. *Neurology*. Nov 12 2013;81(20):1789-92. doi:10.1212/01.wnl.0000435555.13695.22

19. Corona-Cedillo R, Saavedra-Navarrete MT, Espinoza-Garcia JJ, Mendoza-Aguilar AN, Ternovoy SK, Roldan-Valadez E. Imaging Assessment of the Postoperative Spine: An Updated Pictorial Review of Selected Complications. *Biomed Res Int*. 2021;2021:9940001. doi:10.1155/2021/9940001

20. Fisher BM, Cowles S, Matulich JR, Evanson BG, Vega D, Dissanaik S. Is magnetic resonance imaging in addition to a computed tomographic scan necessary to identify clinically significant cervical spine injuries in obtunded blunt trauma patients? *Am J Surg*. Dec 2013;206(6):987-93; discussion 993-4. doi:10.1016/j.amjsurg.2013.08.021

21. Choi BW, Song KJ, Chang H. Ossification of the posterior longitudinal ligament: a review of literature. *Asian Spine J*. Dec 2011;5(4):267-76. doi:10.4184/asj.2011.5.4.267

22. American College of Radiology. ACR Appropriateness Criteria® Myelopathy. American College of Radiology (ACR). Updated 2020. Accessed November 9, 2021. <https://acsearch.acr.org/docs/69484/Narrative/>
23. Behrbalk E, Salame K, Regev GJ, Keynan O, Boszczyk B, Lidar Z. Delayed diagnosis of cervical spondylotic myelopathy by primary care physicians. *Neurosurg Focus*. Jul 2013;35(1):E1. doi:10.3171/2013.3.Focus1374
24. Davies BM, Mowforth OD, Smith EK, Kotter MR. Degenerative cervical myelopathy. *Bmj*. Feb 22 2018;360:k186. doi:10.1136/bmj.k186
25. de Oliveira Vilaça C, Orsini M, Leite MA, et al. Cervical Spondylotic Myelopathy: What the Neurologist Should Know. *Neurol Int*. Nov 2 2016;8(4):6330. doi:10.4081/ni.2016.6330
26. Waly FJ, Abduljabbar FH, Fortin M, Nooh A, Weber M. Preoperative Computed Tomography Myelography Parameters as Predictors of Outcome in Patients With Degenerative Cervical Myelopathy: Results of a Systematic Review. *Global Spine J*. Sep 2017;7(6):521-528. doi:10.1177/2192568217701101
27. American College of Radiology. ACR Appropriateness Criteria® Suspected Spine Trauma American College of Radiology. Updated 2018. Accessed November 9, 2021. <https://acsearch.acr.org/docs/69359/Narrative/>
28. American College of Radiology. ACR Appropriateness Criteria® Inflammatory Back Pain: Known or Suspected Axial Spondyloarthritis. American College of Radiology (ACR). Updated 2021. Accessed November 10, 2021. <https://acsearch.acr.org/docs/3094107/Narrative/>
29. Koivikko MP, Koskinen SK. MRI of cervical spine injuries complicating ankylosing spondylitis. *Skeletal Radiol*. Sep 2008;37(9):813-9. doi:10.1007/s00256-008-0484-x
30. Taljanovic MS, Hunter TB, Wisneski RJ, et al. Imaging characteristics of diffuse idiopathic skeletal hyperostosis with an emphasis on acute spinal fractures: review. *AJR Am J Roentgenol*. Sep 2009;193(3 Suppl):S10-9, Quiz S20-4. doi:10.2214/ajr.07.7102
31. American College of Radiology. ACR Appropriateness Criteria® Management of Vertebral Compression Fractures. American College of Radiology. Updated 2018. Accessed November 10, 2021. <https://acsearch.acr.org/docs/70545/Narrative/>
32. Alexandru D, So W. Evaluation and management of vertebral compression fractures. *Perm J*. Fall 2012;16(4):46-51. doi:10.7812/tpj/12-037
33. Grams AE, Gempt J, Förschler A. Comparison of spinal anatomy between 3-Tesla MRI and CT-myelography under healthy and pathological conditions. *Surg Radiol Anat*. Jul 2010;32(6):581-5. doi:10.1007/s00276-009-0601-0
34. Morita M, Miyauchi A, Okuda S, Oda T, Iwasaki M. Comparison between MRI and myelography in lumbar spinal canal stenosis for the decision of levels of decompression surgery. *J Spinal Disord Tech*. Feb 2011;24(1):31-6. doi:10.1097/BSD.0b013e3181d4c993
35. Naganawa T, Miyamoto K, Ogura H, Suzuki N, Shimizu K. Comparison of magnetic resonance imaging and computed tomogram-myelography for evaluation of cross sections of cervical spinal morphology. *Spine (Phila Pa 1976)*. Jan 1 2011;36(1):50-6. doi:10.1097/BRS.0b013e3181cb469c
36. Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis and Treatment of Lumbar Disc Herniation with Radiculopathy. North American Spine Society (NASS). Updated 2012. Accessed November 11, 2021.

<https://www.spine.org/Portals/0/Assets/Downloads/ResearchClinicalCare/Guidelines/LumbarDiscHerniation.pdf>

37. Ozdoba C, Gralla J, Rieke A, Binggeli R, Schroth G. Myelography in the Age of MRI: Why We Do It, and How We Do It. *Radiol Res Pract*. 2011;2011:329017. doi:10.1155/2011/329017
38. Kim YS, Han IH, Lee IS, Lee JS, Choi BK. Imaging findings of solitary spinal bony lesions and the differential diagnosis of benign and malignant lesions. *J Korean Neurosurg Soc*. 2012;52(2):126-132. doi:10.3340/jkns.2012.52.2.126
39. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Central Nervous System Cancers Version 2.2021. National Comprehensive Cancer Network (NCCN). Updated September 8, 2021. Accessed February 22, 2022.
https://www.nccn.org/professionals/physician_gls/pdf/cns.pdf
40. Ziu E, Viswanathan VK, Mesfin FB. Spinal Metastasis. StatPearls Publishing. Updated August 27, 2021. Accessed November 10, 2021. <https://www.ncbi.nlm.nih.gov/books/NBK441950/>
41. McDonald MA, Kirsch CFE, Amin BY, et al. ACR Appropriateness Criteria® Cervical Neck Pain or Cervical Radiculopathy. *J Am Coll Radiol*. May 2019;16(5s):S57-s76. doi:10.1016/j.jacr.2019.02.023
42. American College of Radiology. ACR Appropriateness Criteria® Suspected Spine Infection. American College of Radiology (ACR). Updated 2021. Accessed November 10, 2021.
<https://acsearch.acr.org/docs/3148734/Narrative/>
43. Lener S, Hartmann S, Barbagallo GMV, Certo F, Thomé C, Tschugg A. Management of spinal infection: a review of the literature. *Acta Neurochir (Wien)*. Mar 2018;160(3):487-496. doi:10.1007/s00701-018-3467-2
44. Bond A, Manian FA. Spinal Epidural Abscess: A Review with Special Emphasis on Earlier Diagnosis. *Biomed Res Int*. 2016;2016:1614328. doi:10.1155/2016/1614328
45. Berbari EF, Kanj SS, Kowalski TJ, et al. 2015 Infectious Diseases Society of America (IDSA) Clinical Practice Guidelines for the Diagnosis and Treatment of Native Vertebral Osteomyelitis in Adults. *Clin Infect Dis*. Sep 15 2015;61(6):e26-46. doi:10.1093/cid/civ482
46. Düz B, Gocmen S, Secer HI, Basal S, Gönül E. Tethered cord syndrome in adulthood. *J Spinal Cord Med*. 2008;31(3):272-8. doi:10.1080/10790268.2008.11760722
47. Milhorat TH, Bolognese PA, Nishikawa M, et al. Association of Chiari malformation type I and tethered cord syndrome: preliminary results of sectioning filum terminale. *Surg Neurol*. Jul 2009;72(1):20-35. doi:10.1016/j.surneu.2009.03.008
48. Zalatimo O. Tethered Spinal Cord Syndrome. American Association of Neurological Surgeons (AANS). Updated 2021. Accessed November 10, 2021.
<https://www.aans.org/Patients/Neurosurgical-Conditions-and-Treatments/Tethered-Spinal-Cord-Syndrome>
49. Whitson WJ, Lane JR, Bauer DF, Durham SR. A prospective natural history study of nonoperatively managed Chiari I malformation: does follow-up MRI surveillance alter surgical decision making? *J Neurosurg Pediatr*. Aug 2015;16(2):159-66. doi:10.3171/2014.12.Peds14301
50. Timpone VM, Patel SH. MRI of a syrinx: is contrast material always necessary? *AJR Am J Roentgenol*. May 2015;204(5):1082-5. doi:10.2214/ajr.14.13310
51. American College of Radiology. ACR Appropriateness Criteria® Headache. American College of Radiology. Updated 2019. Accessed November 1, 2021.
<https://acsearch.acr.org/docs/69482/Narrative/>

52. American College of Radiology. ACR Appropriateness Criteria® Headache-Child. American College of Radiology. Updated 2017. Accessed November 10, 2021.
<https://acsearch.acr.org/docs/69439/Narrative/>
53. Strahle J, Smith BW, Martinez M, et al. The association between Chiari malformation Type I, spinal syrinx, and scoliosis. *J Neurosurg Pediatr*. Jun 2015;15(6):607-11.
doi:10.3171/2014.11.Peds14135
54. Juvenile Scoliosis. Scoliosis Research Society (SRS). Updated 2021. Accessed November 10, 2021. <https://www.srs.org/professionals/online-education-and-resources/conditions-and-treatments/juvenile-scoliosis>
55. American College of Radiology. ACR Appropriateness Criteria® Scoliosis-Child. American College of Radiology. Updated 2018. Accessed November 10, 2021.
<https://acsearch.acr.org/docs/3101564/Narrative/>
56. Trenga AP, Singla A, Feger MA, Abel MF. Patterns of congenital bony spinal deformity and associated neural anomalies on X-ray and magnetic resonance imaging. *J Child Orthop*. Aug 2016;10(4):343-52. doi:10.1007/s11832-016-0752-6
57. Ozturk C, Karadereler S, Ornek I, Enercan M, Ganiyusufoglu K, Hamzaoglu A. The role of routine magnetic resonance imaging in the preoperative evaluation of adolescent idiopathic scoliosis. *Int Orthop*. Apr 2010;34(4):543-6. doi:10.1007/s00264-009-0817-y
58. Strahle J, Muraszko KM, Kapurch J, Bapuraj JR, Garton HJ, Maher CO. Chiari malformation Type I and syrinx in children undergoing magnetic resonance imaging. *J Neurosurg Pediatr*. Aug 2011;8(2):205-13. doi:10.3171/2011.5.Peds1121
59. Radic JAE, Cochrane DD. Choosing Wisely Canada: Pediatric Neurosurgery Recommendations. *Paediatr Child Health*. Sep 2018;23(6):383-387. doi:10.1093/pch/pxy012
60. Hertzler DA, 2nd, DePowell JJ, Stevenson CB, Mangano FT. Tethered cord syndrome: a review of the literature from embryology to adult presentation. *Neurosurg Focus*. Jul 2010;29(1):E1. doi:10.3171/2010.3.Focus1079
61. Shah LM, Salzman KL. Imaging of spinal metastatic disease. *Int J Surg Oncol*. 2011;2011:769753. doi:10.1155/2011/769753
62. Akgul O, Ozgocmen S. Classification criteria for spondyloarthropathies. *World J Orthop*. Dec 18 2011;2(12):107-15. doi:10.5312/wjo.v2.i12.07
63. Bennett AN, Marzo-Ortega H, Rehman A, Emery P, McGonagle D. The evidence for whole-spine MRI in the assessment of axial spondyloarthritis. *Rheumatology (Oxford)*. Mar 2010;49(3):426-32. doi:10.1093/rheumatology/kep427
64. Ostergaard M, Lambert RG. Imaging in ankylosing spondylitis. *Ther Adv Musculoskelet Dis*. Aug 2012;4(4):301-11. doi:10.1177/1759720x11436240
65. Sieper J, Rudwaleit M, Baraliakos X, et al. The Assessment of SpondyloArthritis international Society (ASAS) handbook: a guide to assess spondyloarthritis. *Ann Rheum Dis*. Jun 2009;68 Suppl 2:ii1-44. doi:10.1136/ard.2008.104018
66. Chhetri SK, Gow D, Shaunak S, Varma A. Clinical assessment of the sensory ataxias; diagnostic algorithm with illustrative cases. *Pract Neurol*. Aug 2014;14(4):242-51.
doi:10.1136/practneurol-2013-000764
67. Foster H, Drummond P, Jandial S, Clinch J, Wood M, Driscoll S. Evaluation of gait disorders in children. BMJ Best Practice. Updated February 23, 2021. Accessed November 2, 2021.
<https://bestpractice.bmj.com/topics/en-us/709>

68. Stanford Medicine. Gait Abnormalities. Stanford University. Updated 2021. Accessed November 2, 2021. <https://stanfordmedicine25.stanford.edu/the25/gait.html>
69. Haynes KB, Wimberly RL, VanPelt JM, Jo CH, Riccio AI, Delgado MR. Toe Walking: A Neurological Perspective After Referral From Pediatric Orthopaedic Surgeons. *J Pediatr Orthop*. Mar 2018;38(3):152-156. doi:10.1097/bpo.0000000000001115
70. Marshall FJ. Approach to the elderly patient with gait disturbance. *Neurol Clin Pract*. Jun 2012;2(2):103-111. doi:10.1212/CPJ.0b013e31825a7823
71. Pirker W, Katzenschlager R. Gait disorders in adults and the elderly : A clinical guide. *Wien Klin Wochenschr*. Feb 2017;129(3-4):81-95. doi:10.1007/s00508-016-1096-4
72. Vitzthum HE, Dalitz K. Analysis of five specific scores for cervical spondylogenic myelopathy. *Eur Spine J*. Dec 2007;16(12):2096-103. doi:10.1007/s00586-007-0512-x
73. Pomerantz SR. Myelography: modern technique and indications. *Handb Clin Neurol*. 2016;135:193-208. doi:10.1016/b978-0-444-53485-9.00010-6
74. Ahmed A. MRI features of disseminated 'drop metastases'. *S Afr Med J*. Jul 2008;98(7):522-3.
75. Batool A, Kasi A. Leptomeningeal Carcinomatosis. StatPearls Publishing Copyright © 2022, StatPearls Publishing LLC. Updated March 25, 2021. Accessed February 25, 2022. <https://www.ncbi.nlm.nih.gov/books/NBK499862/>
76. Dias M, Partington M. Congenital Brain and Spinal Cord Malformations and Their Associated Cutaneous Markers. *Pediatrics*. Oct 2015;136(4):e1105-19. doi:10.1542/peds.2015-2854

ADDITIONAL RESOURCES

1. Ando K, Imagama S, Kobayashi K, et al. Clinical Features of Thoracic Myelopathy: A Single-Center Study. *J Am Acad Orthop Surg Glob Res Rev*. Nov 2019;3(11)doi:10.5435/JAAOSGlobal-D-18-00090
2. Budoff MJ, Khairallah W, Li D, et al. Trabecular bone mineral density measurement using thoracic and lumbar quantitative computed tomography. *Acad Radiol*. Feb 2012;19(2):179-83. doi:10.1016/j.acra.2011.10.006
3. Choosing Wisely Clinician Lists: Spine. ABIM Foundation. Updated 2021. Accessed January 3, 2022. <https://www.choosingwisely.org/clinician-lists/#keyword=spine&topic-area=Radiology&service=Imaging>
4. D'Alessandro DM. Does This Sacral Dimple Need to be Evaluated? *PediatricEducation.org™*. Updated July 20, 2009. Accessed November 10, 2021. <https://pediatriceducation.org/2009/07/20/does-this-sacral-dimple-need-to-be-evaluated/>
5. Graeber A, Cecava ND. *Vertebral Osteomyelitis*. StatPearls Publishing. Updated July 22, 2021. Accessed November 10, 2021. <https://www.ncbi.nlm.nih.gov/books/NBK532256/>
6. Hou X, Sun C, Liu X, et al. Clinical Features of Thoracic Spinal Stenosis-associated Myelopathy: A Retrospective Analysis of 427 Cases. *Clin Spine Surg*. Mar 2016;29(2):86-9. doi:10.1097/bsd.0000000000000081

7. North American Spine Society. Five things physicians and patients should question. Choosing Wisely Initiative ABIM Foundation. Updated 2021. Accessed November 9, 2021. <https://www.choosingwisely.org/societies/north-american-spine-society/>
8. Müller D, Bauer JS, Zeile M, Rummeny EJ, Link TM. Significance of sagittal reformations in routine thoracic and abdominal multislice CT studies for detecting osteoporotic fractures and other spine abnormalities. *Eur Radiol*. Aug 2008;18(8):1696-702. doi:10.1007/s00330-008-0920-2
9. Stolper K, Haug JC, Christensen CT, Samsey KM, April MD. Prevalence of thoracic spine lesions masquerading as cauda equina syndrome: yield of a novel magnetic resonance imaging protocol. *Intern Emerg Med*. Dec 2017;12(8):1259-1264. doi:10.1007/s11739-016-1565-9

Reviewed / Approved by NIA Clinical Guideline Committee

GENERAL INFORMATION

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

Disclaimer: Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.