

*National Imaging Associates, Inc.*	
Clinical guidelines THORACIC SPINE MRI	Original Date: September 1997
CPT Codes: 72146, 72147, 72157, +0698T	Last Revised Date: <del>May 2023</del> March 2022
Guideline Number: NIA_CG_042	Implementation Date: January 20 <del>24</del> 23

### GENERAL INFORMATION

- *It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.*
- *Where a specific clinical indication is not directly addressed in this guideline, medical necessity determination will be made based on widely accepted standard of care criteria. These criteria are supported by evidence-based or peer-reviewed sources such as medical literature, societal guidelines and state/national recommendations.*

### INDICATIONS FOR THORACIC SPINE MRI

\*If there is a combination request\* for an overlapping body part, either requested at the same time or sequentially (within the past 3 months) the results of the prior study should be:

- Inconclusive or show a need for additional or follow-up imaging evaluation OR
- The office notes should clearly document an indication why overlapping imaging is needed and how it will change management for the patient (the entire spinal cord and/or autonomic postganglionic chain must be assessed)

(\*Unless approvable in the [combination section](#) as noted in the guidelines)

### For evaluation of neurologic deficits<sup>1-4</sup>

- With any of the following new neurological deficits documented on physical exam
  - Extremity muscular weakness (and not likely caused by plexopathy or peripheral neuropathy)<sup>5, 6</sup>

~~\*National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.~~

- ~~Pathologic (e.g., Babinski, Lhermitte's sign, Chaddock Sign, Hoffman's and other upper motor neuron signs); OR abnormal deep tendon reflexes (and not likely caused by plexopathy, or peripheral neuropathy) Pathologic (e.g., Babinski, Chaddock Sign) or abnormal reflexes<sup>7</sup>~~
- ~~Abnormal reflexes (decreased or absent (and not likely caused by plexopathy, or peripheral neuropathy)~~
- Absent/decreased sensory changes along a particular thoracic dermatome (nerve distribution): pin prick, touch, vibration, proprioception, or temperature (and not likely caused by plexopathy, or peripheral neuropathy)
- Upper or lower extremity increase muscle tone/spasticity, and likely localized to the thoracic spinal cord
- New onset bowel or bladder dysfunction (e.g., retention or incontinence)- not related to an inherent bowel or bladder process
- Gait abnormalities, most likely cause by a suspected or known myelopathy (see Table 1 for more details)
- Suspected thoracic cord compression with any neurological deficits as listed above

#### For evaluation of back pain with any of the following<sup>7-9</sup>

- With new or worsening objective neurologic deficits (as listed above) on exam
- Failure of conservative treatment\* for at least six (6) weeks within the last six (6) months
- With progression or worsening of symptoms during the course of conservative treatment\*
- With an abnormal electromyography (EMG) or nerve conduction study (if performed) indicating a thoracic radiculopathy. (EMG is not recommended to determine the cause of axial lumbar, thoracic, or cervical spine pain)<sup>10</sup>
- ~~Isolated thoracic~~ back pain in pediatric population<sup>11</sup> – conservative care not required if red flags present ~~AND initial radiographs have been performed.:~~
- Red flags that prompt imaging ~~should~~ include any the presence of the following:
  - ~~A~~ge 5 or younger, OR
  - ~~C~~onstant pain, OR
  - ~~P~~ain lasting > 4 weeks, OR
  - ~~A~~bnormal neurologic examination, OR
  - ~~E~~arly morning stiffness and/or gelling, OR
  - ~~N~~ight pain that prevents or disrupts sleep, OR
  - Radicular pain, OR
  - ~~radicular pain; F~~ever or w
  - ~~;~~ Weight loss or malaise, OR
  - ~~malaise; P~~ostural changes (e.g., kyphosis or scoliosis), OR ~~and~~

- Limp (or refusal to walk in a younger child ~~child < 5yo~~) ~~AND initial radiographs have been performed~~<sup>12, 13</sup>

**As part of initial pre-operative / post-operative / procedural evaluation (“CT best examination to assess for hardware complication, extent of fusion and pseudoarthrosis”<sup>14, 15</sup> and MRI for cord, nerve root compression, disc pathology or post-op infection)**

- For preoperative evaluation/planning
- Prior to spinal cord stimulator to exclude canal stenosis if no prior MRI imaging of the thoracic spine has been done recently<sup>16, 17</sup>
- CSF leak highly suspected and supported by patient history and/or physical exam findings (leak (known or suspected spontaneous (idiopathic) intracranial hypotension (SIH), post lumbar puncture headache, post spinal surgery headache, orthostatic headache, rhinorrhea or otorrhea, or cerebrospinal-venous fistula or dural fistula))
- A follow-up study may be needed to help evaluate a patient’s progress after treatment, procedure, intervention, or surgery in the last 6 months. Documentation requires a medical reason that clearly indicates why additional imaging is needed for the type and area(s) requested (routine surveillance post-op not indicated without symptoms)
- Surgical infection as evidenced by signs/symptoms, laboratory, or prior imaging findings
- New or changing neurological deficits or symptoms post-operatively<sup>14, 18</sup>- see neurological deficit section above
- When combo requests (see above statement<sup>+</sup>) are submitted (i.e., MRI and CT of the spine), the office notes should clearly document the need for both studies to be done simultaneously, i.e., the need for both soft tissue and bony anatomy is required<sup>19</sup>
  - Combination requests where both thoracic spine CT and MRI thoracic spine are both approvable (not an all-inclusive list):
    - OPLL (Ossification of posterior longitudinal ligament)-
      - Most common in cervical spine (rare but more severe in thoracic spine)<sup>20</sup>
    - Pathologic or complex fractures
    - Malignant process of spine with both bony and soft tissue involvement
    - Clearly documented indication for bony and soft tissue abnormality where assessment will change management for the patient

#### **For evaluation of suspected myelopathy<sup>21-25</sup>**

- Does **NOT** require conservative care
- Progressive symptoms including unsteadiness, broad-based gait, increased muscle tone, pins and needles sensation, weakness and wasting of the lower limbs, and diminished sensation to light touch, temperature, proprioception, and vibration; limb hyperreflexia and pathologic reflexes; bowel and bladder dysfunction in more severe cases
- Any of the neurological deficits as noted above

### For evaluation of known or suspected multiple sclerosis (MS)<sup>25-28</sup>

- Suspected or known MS with new or changing symptoms suggesting underlying thoracic spinal cord disease (i.e., transverse myelitis, progressive myelopathy)
- Suspected or known pediatric demyelinating diseases (MS/ADEM)

### Combination studies for MS

- **These body regions might be evaluated separately or in combination as guided by physical examination findings (e.g., localization to a particular segment of the spinal cord), patient history (e.g., symptom(s), time course, and where in the CNS the likely localization(s) is/are), and other available information, including prior imaging.**
  - Cervical **and/or** Thoracic MRI for evaluation of highly suspected multiple sclerosis (MS) when Brain MRI has indeterminate findings and/or does not fulfill the McDonald criteria for the diagnosis of MS<sup>27</sup>
  - Cervical **and/or** Thoracic MRI with suspected transverse myelitis-with appropriate clinical symptoms (e.g., bilateral weakness, sensory disturbance, and autonomic dysfunction which typically evolve over hours or days)
  - Brain MRI with Cervical **and/or** Thoracic MRI for evaluation of neuromyelitis optica spectrum disorders (recurrent or bilateral optic neuritis; recurrent transverse myelitis)<sup>29</sup>
  - Known MS- entire CNS axis (Brain, **and/or** Cervical **and/or** Thoracic spine) is approvable prior to the initiation or change of disease modification treatments and assess disease burden (to establish a new baseline)
  - Known MS- Follow-up scans, including brain and spine imaging, if patients have known spine disease:
    - 6-12 months after starting/changing treatment
    - Every 1-2 years while on disease-modifying therapy to assess for subclinical disease activity, less frequently when stable for 2-3 years

### For evaluation of trauma or acute injury<sup>30</sup>

- Presents with any of the following [neurological deficits](#) as above
- With progression or worsening of symptoms during the course of a trial of conservative treatment\*
- History of underlying spinal abnormalities (i.e., ankylosing spondylitis, diffuse idiopathic skeletal hyperostosis) (Both MRI and CT would be approvable)<sup>31-33</sup>
- When the patient is clinically unevaluable or there are preliminary imaging findings (x-ray or CT) needing further evaluation

("MRI and CT provide complementary information. When indicated it is appropriate to perform both examinations").<sup>30</sup>

## For evaluation of known or new compression fractures with worsening back pain<sup>30, 34</sup>

- With history of malignancy
  - To aid in differentiation of benign osteoporotic fractures from metastatic disease
    - A follow-up MRI in 6-8 weeks after initial MRI when initial imaging cannot decipher (indeterminate) benign osteoporotic fracture from metastatic disease<sup>35</sup>
- With an associated new focal neurologic deficit as above
- Prior to a planned surgery/intervention or if the results of the MRI will change management

## For evaluation of tumor, cancer, or metastasis with any of the following, as per NCCN: (MRI is usually the preferred study, but CT may be needed to further characterize solitary indeterminate lesions seen on MRI)<sup>36-38</sup>

- **Primary tumor**
  - Initial staging ~~or re-staging of a known~~ primary spinal tumor<sup>39</sup> as per NCCN
  - Follow-up of known primary cancer of patient undergoing active treatment within the past year or as per surveillance imaging guidance for that cancer
  - Known primary tumor with new signs or symptoms (e.g., new or increasing nontraumatic pain, physical, laboratory, and/or imaging findings)
  - With an associated new focal neurologic deficit as above<sup>40</sup>
- **Metastatic tumor**
  - With evidence of metastasis on bone scan needing further clarification OR inconclusive findings on a prior imaging exam
  - With an associated new focal neurologic deficit<sup>40</sup>
  - Known malignancy with new signs or symptoms (e.g., new or increasing nontraumatic pain, radiculopathy or back pain that occurs at night and wakes the patient from sleep with known active cancer, physical, laboratory, and/or imaging findings) in a tumor that tends to metastasize to the spine<sup>33-35</sup>

## Further evaluation of indeterminate ~~or questionable~~ findings on prior imaging (unless follow up is otherwise specified within the guideline): :

- For initial evaluation of an inconclusive finding on a prior imaging report that requires further clarification:
- One follow-up exam of a prior indeterminate MR/CT finding to ensure no suspicious interval change has occurred. (No further surveillance unless specified as highly suspicious or change was found on last follow-up exam.<sup>42</sup>)

- ~~For evaluation of inconclusive finding on prior imaging that requires further clarification~~
  - ~~One follow-up exam to ensure no suspicious change has occurred in prior imaging finding. No further surveillance unless specified as highly suspicious or change was found on last follow-up exam<sup>39</sup>~~

**Indication for combination studies for the initial pre-therapy staging of cancer, OR active monitoring for recurrence as clinically indicated, OR evaluation of suspected metastases**

- $\leq 5$  concurrent studies to include CT or MRI of any of the following areas as appropriate depending on the cancer: Neck, Abdomen, Pelvis, Chest, Brain, Cervical Spine, Thoracic Spine or Lumbar Spine

**For evaluation of known or suspected infection (osteomyelitis), abscess, or inflammatory disease<sup>41, 42</sup>**

- **Infection**
  - As evidenced by signs and/or symptoms, laboratory (i.e., abnormal white blood cell count, ESR and/or CRP) or prior imaging findings<sup>43</sup>
  - Follow-up imaging of infection
    - With worsening symptoms/laboratory values (i.e., white blood cell count, ESR/CRP) or radiographic findings<sup>44</sup>
- **Spondyloarthropathies**
  - Ankylosing Spondylitis/Spondyloarthropathies with non-diagnostic or indeterminate x-ray and appropriate rheumatology workup

**For evaluation of spine abnormalities related to immune system suppression, e.g., HIV, chemotherapy, leukemia, or lymphoma<sup>42</sup>**

- As evidenced by signs/symptoms, laboratory, or prior imaging findings

### **Other Indications for a Thoracic Spine MRI**

(Note- See combination requests, below, for initial advanced imaging assessment and pre-operatively)

- Tethered cord, or spinal dysraphism (known or suspected) based on preliminary imaging, neurological exam, and/or high-risk cutaneous stigmata<sup>45-47</sup>
- Known Arnold-Chiari syndrome (For initial imaging (one-time initial MRI-modality assessment) see combination below)
  - Known Chiari I malformation without syrinx or hydrocephalus, follow-up imaging after initial diagnosis with new or changing signs/symptoms or exam findings consistent with spinal cord pathology<sup>48</sup>
  - Known Chiari II ( Arnold-Chiari syndrome), III, or IV malformation
- Syrinx or syringomyelia (known or suspected)

- With neurologic findings and/or predisposing conditions (e.g., Chiari malformation, prior trauma, neoplasm, arachnoiditis, severe spondylosis<sup>49</sup>)
- To further characterize a suspicious abnormality seen on prior imaging
- Known syrinx with new/worsening symptoms
- Toe walking in a child with signs/symptoms of myelopathy localized to the Thoracic Spine
- Suspected neuroinflammatory Conditions/Diseases (e.g., sarcoidosis, Behcet's)
  - After detailed neurological exam and ~~basic testing~~appropriate initial work up completed

## COMBINATION STUDIES WITH THORACIC SPINE MRI

### Cervical and Thoracic MRI

- Initial evaluation of known or suspected syrinx or syringomyelia
  - With neurologic findings and/or predisposing conditions (e.g., Chiari malformation, prior trauma, neoplasm, arachnoiditis, severe spondylosis<sup>49</sup>)
  - To further characterize a suspicious abnormality seen on prior imaging
  - Known syrinx with new/worsening symptom

### Any combination of Cervical and/or Thoracic and/or Lumbar MRIs

Note: These body regions might be evaluated separately or in combination as documented in the clinical notes by physical examination findings (e.g., localization to a particular segment of the spinal cord), patient history, and other available information, including prior imaging.

**Exception-** Indications for combination studies<sup>50, 51</sup>: Are approved indications as noted below and being performed in children who will need anesthesia for the procedure

- Any combination of these studies for:
  - Survey/complete initial assessment of infant/child with congenital scoliosis or juvenile idiopathic scoliosis under the age of 10<sup>52-54</sup> (e.g., congenital scoliosis, idiopathic scoliosis, scoliosis with vertebral anomalies)
  - In the presence of neurological deficit, progressive spinal deformity, or for preoperative planning<sup>55</sup>
  - Back pain with known vertebral anomalies (hemivertebrae, hypoplasia, agenesis, butterfly, segmentation defect, bars, or congenital wedging) in a child on preliminary imaging
  - Scoliosis with any of the following<sup>56</sup>:
    - Progressive spinal deformity;
    - Neurologic deficit (new or unexplained);
    - Early onset;

- Atypical curve (e.g., short segment, >30° kyphosis, left thoracic curve, associated organ anomalies);
    - Pre-operative planning; OR
    - When office notes clearly document how imaging will change management
  - Arnold-Chiari malformations<sup>57, 58</sup>
    - Arnold-Chiari I
      - For evaluation of spinal abnormalities associated with initial diagnosis of Arnold-Chiari Malformation. (C/T/L spine due to association with tethered cord and syringomyelia), and initial imaging has not been completed<sup>46, 52</sup>
    - Arnold-Chiari II-IV - For initial evaluation and follow-up as appropriate
      - Usually associated with open and closed spinal dysraphism, particularly meningocele
  - Tethered cord, or spinal dysraphism (known or suspected) based on preliminary imaging, neurological exam, and/or high-risk cutaneous stigmata,<sup>45-47</sup> when anesthesia required for imaging<sup>59</sup> (e.g., meningocele, lipomeningocele, diastematomyelia, fatty/thickened filum terminale, and other spinal cord malformations)
  - Oncological Applications (e.g., primary nervous system, metastatic)
    - Drop metastasis from brain or spine (imaging also includes brain)- see [Overview](#)
    - Suspected leptomeningeal carcinomatosis (LC)<sup>60</sup> -see [Overview](#)
    - Any combination of these for spinal survey in patient with metastases
    - Tumor evaluation and monitoring in neurocutaneous syndromes
  - CSF leak highly suspected and supported by patient history and/or physical exam findings (leak (known or suspected spontaneous (idiopathic) intracranial hypotension (SIH), post lumbar puncture headache, post spinal surgery headache, orthostatic headache, rhinorrhea or otorrhea, or cerebrospinal-venous fistula))
- 

## BACKGROUND

Magnetic resonance imaging (MRI) produces high quality multiplanar images of organs and structures within the body without using ionizing radiation. It is used for evaluation, assessment of severity, and follow-up of diseases of the spine and is the preferred modality for imaging intervertebral disc degeneration. High contrast resolution (soft tissue contrast) and multiplanar imaging (sagittal as well as axial planes) are helpful in the evaluation of possible disc herniation and detecting nerve root compression. MRI is one of the most useful techniques to evaluate spine infection and is also used to evaluate tumors, cancer, and immune system suppression.



## OVERVIEW

~~Ankylosing Spondylitis/Spondyloarthropathies is a cause of back or sacroiliac pain of insidious onset (usually > 3 month), associated with morning stiffness not relieved with rest (usually age at onset < 40). It is associated with any of the following<sup>62-65</sup>:~~

- ~~• Sedimentation rate and/or C-reactive protein (not an essential criteria)~~
- ~~• HLA B27 (not an essential criteria)~~
- ~~• Non diagnostic or indeterminate x ray~~
- ~~• Personal or family history of sacroiliitis, peripheral inflammatory arthritis, and/or inflammatory bowel disease~~

**\*Conservative Therapy** – (Spine) should include a multimodality approach consisting of a **combination of active and inactive components**. Inactive components, such as rest, ice, heat, modified activities, medical devices, acupuncture and/or stimulators, medications, injections (epidural, facet, bursal, and/or joint, not including trigger point), and diathermy can be utilized. Active modalities may consist of physical therapy, a physician-supervised home exercise program\*\*, and/or osteopathic manipulative medicine (OMT) or chiropractic care when considered safe and appropriate.

**\*\*Home Exercise Program - (HEP)/Therapy** – the following elements are required to meet guidelines for completion of conservative therapy<sup>15, 61</sup>:

- Information provided on exercise prescription/plan AND
- Follow-up with member with documentation provided regarding lack of improvement (failed) after completion of HEP (after suitable 6-week period), or inability to complete HEP due to physical reason- i.e., increased pain, inability to physically perform exercises. (Patient inconvenience or noncompliance without explanation does not constitute “inability to complete” HEP).
- Dates and duration of failed PT, physician-supervised HEP, or chiropractic treatment should be documented in the original office notes or an addendum to the notes.

### **Infection, Abscess, or Inflammatory disease**

~~Most common site is the lumbar spine (58%), followed by the thoracic spine (30%) and the cervical spine (11%)<sup>[66]</sup>~~

~~High risk populations (indwelling hardware, history of endocarditis, IVDA, recent procedures) with appropriate signs/symptoms~~

**Table 1: Gait and spine imaging<sup>62-67</sup>**

Gait	Characteristic	Work up/Imaging
Hemiparetic	Spastic unilateral, circumduction	Brain and/or, Cervical spine imaging based on associated symptoms
Diplegic	Spastic bilateral, circumduction	Brain, Cervical and Thoracic Spine imaging
Myelopathic	Wide based, stiff, unsteady	Cervical and/or Thoracic spine MRI based on associated symptoms
<b>Cerebellar</b> Ataxic	Broad based, clumsy, staggering, lack of coordination, usually also with limb ataxia	Brain imaging <a href="#">see Brain MRI Guideline</a>
Apraxic	Magnetic, shuffling, difficulty initiating	Brain imaging <a href="#">see Brain MRI Guideline</a>
Parkinsonian	Stooped, small steps, rigid, turning en bloc, decreased arm swing	Brain Imaging <a href="#">see Brain MRI Guideline</a>
Choreiform	Irregular, jerky, involuntary movements	Medication review, consider brain imaging as per movement disorder Brain MR guidelines
Sensory ataxic	Cautious, stomping, worsening without visual input (ie + Romberg)	EMG, blood work, consider spinal (cervical or thoracic cord imaging) imaging based on EMG
Neurogenic	Steppage, dragging of toes	<ul style="list-style-type: none"> <li>• EMG initial testing;</li> <li>• BUT if there is a foot drop, lumbar spine MRI is appropriate without EMG</li> <li>• Pelvis MR if there is evidence of plexopathy</li> </ul>
Vestibular	Insecure, veer to one side, worse when eyes closed, vertigo	Consider Brain/IAC MRI <a href="#">see Brain MRI Guidelines as per GL</a>

**Table 2: MRI and Cutaneous Stigmata**

Risk Stratification for Various Cutaneous Markers		
High Risk	Intermediate Risk	Low Risk
<ul style="list-style-type: none"> <li>• Hypertrichosis</li> <li>• Infantile hemangioma</li> <li>• Atretic meningocele</li> <li>• DST</li> <li>• Subcutaneous lipoma</li> <li>• Caudal appendage</li> <li>• Segmental hemangiomas in association with LUMBAR<sup>‡</sup> syndrome</li> </ul>	<ul style="list-style-type: none"> <li>• Capillary malformations (also referred to as NFS or salmon patch when pink and poorly defined or PWS when darker red and well-defined)</li> </ul>	<ul style="list-style-type: none"> <li>• Coccygeal dimple</li> <li>• Light hair</li> <li>• Isolated café au lait spots</li> <li>• Mongolian spots</li> <li>• Hypo- and hypermelanotic macules or papules</li> <li>• Deviated or forked gluteal cleft</li> <li>• Nonmidline lesions</li> </ul>
<sup>‡</sup> LUMBAR, lower body hemangioma and other cutaneous defects, urogenital abnormalities, ulcerations, myelopathy, bony defects, anorectal malformations, arterial anomalies, and renal anomalies.		

**Myelopathy** – Symptom severity varies, and a high index of suspicion is essential for making the proper diagnosis in early cases. Symptoms of pain and radiculopathy may not be present. The natural history of myelopathy is characterized by neurological deterioration. The most frequently encountered symptom is gait abnormality (86%) followed by increased muscular reflexes (79.1%), pathological reflexes (65.1%), paresthesia of upper limb (69.8%), and pain (67.4%).<sup>68</sup>

**Ossification Posterior Longitudinal Ligament (OPLL)<sup>20</sup>** – Most common in cervical spine (rare but more severe in thoracic spine)

**Tethered spinal cord syndrome**—a neurological disorder caused by tissue attachments that limit the movement of the spinal cord within the spinal column. Although this condition is rare, it can continue undiagnosed into adulthood. The primary cause is myelomeningocele and lipomyelomeningocele; the following are other associations that vary in severity of symptoms and treatment.

- Dermal sinus tract (a rare congenital deformity)
- Diastematomyelia (split spinal cord)
- Lipoma
- Tumor
- Thickened/tight filum terminale
- History of spine trauma/surgery
- Arnold-Chiari Malformation

Magnetic resonance imaging (MRI) can display the low level of the spinal cord and a thickened filum terminale, the thread-like extension of the spinal cord in the lower back. Treatment depends upon the underlying cause of the tethering. If the only abnormality is a thickened, shortened filum, then limited surgical treatment may suffice.

**MRI and Spinal Infections**—Infection of the spine is not easy to differentiate from other spinal disorders, e.g., degenerative disease, spinal neoplasms, and noninfectious inflammatory lesions. Infections may affect different parts of the spine, e.g., vertebrae, intervertebral discs and paraspinal tissues, or the spinal cord tissue and/or roots themselves. Imaging is important in obtaining an early diagnosis and treatment to avoid permanent neurologic deficits. MRI is the preferred imaging technique to evaluate infections of the spine. With its high contrast resolution and direct multiplanar imaging, MRI can detect and delineate infective lesions irrespective of their spinal location.

**Neck and Back Pain with Cancer History** – Bone is the third most common site of metastases after the liver and the lungs, and approximately two-thirds of all osseous metastases occur in the spine. Approximately 60–70% of patients with systemic cancer will have spinal metastasis. Radiographic (x-ray) examination should be performed in cases of back pain when a patient has a cancer history, but without known active cancer or a tumor that tends to metastasize to the spine. This can make a diagnosis in many cases. This may occasionally allow for selection of bone scan in lieu of MRI in some cases. When radiographs do not answer the clinical question, then MRI may be appropriate after a consideration of conservative care.

76

Most common spine metastasis involving primary metastasis originate from the following tumors in descending order: breast (21%), lung (19%), prostate (7.5%), renal (5%), gastrointestinal (4.5%), and thyroid (2.5%). While all tumors can seed to the spine, the cancers mentioned above metastasize to the spinal column early in the disease process. Spinal metastasis is more commonly found in the thoracic region, followed by the lumbar region, while the cervical region is the least likely site of metastasis.<sup>69</sup>

**Back Pain with Cancer History**—Radiographic (x-ray) examination should be performed in cases of back pain when a patient has a cancer history, but without known active cancer or a tumor that tends to metastasize to the spine. This can make a diagnosis in many cases. This may occasionally allow for selection of bone scan in lieu of MRI in some cases. When radiographs do not answer the clinical question, then MRI may be appropriate after a consideration of conservative care.

“Neoplasms causing VCF (vertebral compression fractures) include: 1) primary bone neoplasms, such as hemangioma (aggressive type) or giant cell tumors, and tumor-like conditions causing bony and cellular remodeling, such as aneurysmal bone cysts, or Paget’s disease (osteitis deformans); 2) primary malignant neoplasms including but not limited to multiple myeloma and lymphoma; and 3) infiltrative metastatic neoplasms, including and not limited to, multiple myeloma and lymphoma, and metastatic neoplasms.”<sup>35</sup>

Most common spine metastasis involving primary metastasis originate from the following tumors in descending order: breast (21%), lung (19%), prostate (7.5%), renal (5%), gastrointestinal (4.5%), and thyroid (2.5%). While all tumors can seed to the spine, the cancers mentioned above metastasize to the spinal column early in the disease process.<sup>76</sup>

**Cauda Equina Syndrome**—Symptoms include severe back pain or sciatica along with one or more of the following:

- Saddle anesthesia—loss of sensation restricted to the area of the buttocks, perineum, and inner surfaces of the thighs (areas that would sit on a saddle)
- Recent bladder/bowel dysfunction
- Achilles reflex absent on both sides
- Sexual dysfunction that can come on suddenly
- Absent anal reflex and bulbocavernosus reflex

**Spinal MRI and Neuromyelitis optica spectrum disorders (NMOSD)**—NMOSD are inflammatory disorders of the central nervous system characterized by severe, immune-mediated demyelination and axonal damage predominantly affecting the optic nerves and spinal cord, but NMOSD may also affect the brain and brainstem. NMOSD can typically be distinguished from multiple sclerosis and other inflammatory disorders by the presence of the aquaporin 4 (AQP4) antibody; although, up to 10% of patients with NMOSD can be seronegative.<sup>77,78</sup> Features of NMOSD include attacks of bilateral or sequential optic neuritis acute transverse myelitis and the area postrema syndrome (with intractable hiccups or nausea

and vomiting). The evaluation of suspected NMOSD entails brain and spinal cord neuroimaging. In contrast to MS (in which spinal cord involvement tends to be incomplete and asymmetric), NMOSD have a longer extent of spinal cord demyelination generally involving three or more vertebral segments.

## MRI and Neurocutaneous Syndromes

- In NF-1, clinical evaluation appears to be more useful to detect complications than is screening imaging in asymptomatic patients. Imaging is indicated in evaluation of suspected tumors based on clinical evaluation and for follow-up of known intracranial and intraspinal tumors.<sup>70</sup>
- Conversely in NF-2, routine MR imaging screening is always indicated, given the high prevalence of CNS tumors, especially vestibular schwannomas. In patients with NF-2, routine screening brain/IAC imaging is indicated annually starting from age 10, if asymptomatic, or earlier with clinical signs/symptoms. Most individuals with NF2 eventually develop a spinal tumor, mostly commonly schwannomas, but meningioma and ependymomas are also seen. Spinal imaging at baseline and every 2 to 3 years is also advised with more frequent imaging, if warranted, based on sites of tumor involvement.<sup>71</sup>
- In patients with Tuberous Sclerosis, Brain MRI should be obtained every 1-3 years up until age 25 for surveillance for CNS abnormalities.<sup>72</sup>
- In Von Hippel Lindau Syndrome, imaging of the brain and spinal cord for hemangioblastomas is recommended every 2 years.<sup>73</sup>
- In Sturge Weber Syndrome, Brain MRI can rule out intracranial involvement after only age 1 and is recommended in patients < 1 year old only if symptomatic.<sup>74</sup>

**Drop Metastases<sup>75</sup>** – Drop metastases are intradural extramedullary spinal metastases that arise from intracranial lesions. Common examples of intracranial neoplasms that result in drop metastases include pineal tumors, ependymomas, medulloblastomas, germinomas, primitive neuroectodermal tumors (PNET), glioblastomas multiform, anaplastic astrocytomas, oligodendrogliomas and less commonly choroid plexus neoplasms and teratomas.

**Leptomeningeal Carcinomatosis<sup>76</sup>** – Leptomeningeal carcinomatosis is complication of cancer in which cancerous cells spread to the membranes (meninges) that covers the brain and spinal cord. The most common solid tumors that involve the leptomeninges are breast, lung, and melanoma, gastrointestinal, and primary central nervous system tumors.

## POLICY HISTORY

Date	Summary
<u>2023</u>	<ul style="list-style-type: none"> <li><del>Updated references</del></li> <li><del>Updated background section</del></li> <li><del>Clarified pathological reflexes</del></li> <li><del>Added Further evaluation of indeterminate or questionable findings on prior imaging</del></li> <li><del>Clarified cerebellar ataxia in gait table</del></li> </ul>
March 2022	<p>Added</p> <ul style="list-style-type: none"> <li><del>Combination request for overlapping body part statement</del></li> <li><del>Clarified muscle weakness not related to plexopathy or peripheral neuropathy</del></li> <li><del>Clarified bowel and bladder dysfunction — not related to an inherent bowel or bladder problem</del></li> <li><del>Clarified combination MS for cervical and/or thoracic spine combination requests</del></li> <li><del>Descriptions for tethered cord</del></li> <li><del>Background section of Drop Metastases</del></li> <li><del>Background section of Leptomeningeal Carcinomatosis</del></li> <li><del>Clarified toe walking in pediatric patient with myelopathy for thoracic spine</del></li> </ul> <p>Removed</p> <ul style="list-style-type: none"> <li><del>Removed from combination section syrinx and syringomyelia and added subsection for cervical and thoracic spine section</del></li> <li><del>Removed pediatric back pain from the total spine combination section</del></li> </ul>
November 2021	<ul style="list-style-type: none"> <li><del>Added +0698T</del></li> </ul>
April 2021	<ul style="list-style-type: none"> <li><del>Added/modified</del> <ul style="list-style-type: none"> <li><del>Modified section on neurological deficits</del></li> <li><del>Back pain in a child added/modified red flags</del></li> <li><del>Gait table in background</del></li> <li><del>Post-surgical modified/clarified surgical criteria for combination exams</del></li> <li><del>Removed myelopathy combination studies</del></li> <li><del>Updated/added MS Criteria</del> <ul style="list-style-type: none"> <li><del>Combination section for initial imaging and follow up</del></li> <li><del>Added pediatric MS</del></li> </ul> </li> <li><del>Modified known tumor imaging into primary and metastatic disease</del></li> <li><del>Added toe walking for pediatric patients</del></li> </ul> </li> </ul>

	<ul style="list-style-type: none"> <li>○ <del>Modified Combination exam wording</del></li> </ul>
May 2020	<ul style="list-style-type: none"> <li>● <del>Added</del> <ul style="list-style-type: none"> <li>○ <del>For evaluation of neurologic deficits when new deficits are present</del></li> <li>○ <del>Removed pars defect section</del></li> <li>○ <del>Added ankylosing spondylitis for evaluation of trauma/acute injury</del></li> <li>○ <del>Added follow up of osteoporotic fracture from metastatic disease</del></li> <li>○ <del>Added transverse myelitis</del></li> <li>○ <del>Modified Initial imaging of new or increasing non-traumatic back pain or radiculopathy or back pain that occurs at night and wakes the patient from sleep with known active cancer and a tumor that tends to metastasize to the spine</del></li> <li>○ <del>Added Imaging of Ossification of the Posterior Longitudinal Ligament (OPPL)</del></li> <li>○ <del>Added Osteopathic Manipulative medicine to conservative care therapy</del></li> </ul> </li> </ul>
June 2019	<ul style="list-style-type: none"> <li>● <del>Added:</del> <ul style="list-style-type: none"> <li>○ <del>new or worsening objective neuro deficits for chronic and acute back pain</del></li> <li>○ <del>CSF leak</del></li> <li>○ <del>last 6 months for allowable post op f/u period and removed EMG comment</del></li> <li>○ <del>red flags specifically for peds back pain and pain related to malignancy, infection, inflammation</del></li> <li>○ <del>new sections: pars defect; compression fractures; congenital abnormalities including section on scoliosis and vertebral anomalies in children w/back pain;</del></li> <li>○ <del>For combination studies cervical/thoracic/lumbar added drop metastasis, tumor evaluation for neurocutaneous syndromes, and abnormalities associated w/Arnold Chiari, as well as separate indication for tethered cord or spinal dysraphism</del></li> <li>○ <del>Myelopathy</del></li> <li>○ <del>Pre op for spinal cord stimulator</del></li> <li>○ <del>Evaluation of MS including NMO disorders and recurrent transverse myelitis</del></li> </ul> </li> </ul>



	<ul style="list-style-type: none"> <li>○ <del>Back pain in cancer patients with known active cancer in tumors that tend to metastasize</del></li> <li>○ <del>Expanded on tethered cord in Other Indications for Imaging and added content on sacral dimple</del></li> </ul>
--	---

## REFERENCES

1. Acharya AB, Fowler JB. Chaddock Reflex. StatPearls Publishing. Updated June 27, 2022. Accessed December 1, 2022. <https://www.ncbi.nlm.nih.gov/books/NBK519555/>
2. Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis and Treatment of Cervical Radiculopathy from Degenerative Disorders. North American Spine Society (NASS). Updated 2010. Accessed December 1, 2022. <https://www.spine.org/Portals/0/Assets/Downloads/ResearchClinicalCare/Guidelines/CervicalRadiculopathy.pdf>
3. Stolper K, Haug JC, Christensen CT, Samsey KM, April MD. Prevalence of thoracic spine lesions masquerading as cauda equina syndrome: yield of a novel magnetic resonance imaging protocol. *Intern Emerg Med*. Dec 2017;12(8):1259-1264. doi:10.1007/s11739-016-1565-9
4. Albert TJ, Murrell SE. Surgical management of cervical radiculopathy. *J Am Acad Orthop Surg*. Nov-Dec 1999;7(6):368-76. doi:10.5435/00124635-199911000-00003
5. Moore KR, Tsuruda JS, Dailey AT. The value of MR neurography for evaluating extraspinal neuropathic leg pain: a pictorial essay. *AJNR Am J Neuroradiol*. Apr 2001;22(4):786-94.
6. Dydyk AM, Hameed S. Lumbosacral Plexopathy. StatPearls Publishing Copyright © 2022, StatPearls Publishing LLC. Updated March 26, 2022. Accessed November 16, 2022. <https://www.ncbi.nlm.nih.gov/books/NBK556030/>
7. Jarvik JG, Gold LS, Comstock BA, et al. Association of early imaging for back pain with clinical outcomes in older adults. *Jama*. Mar 17 2015;313(11):1143-53. doi:10.1001/jama.2015.1871
8. American Association of Neurological Surgeons, Congress of Neurological Surgeons. Five things physicians and patients should question: Don't obtain imaging (plain radiographs, magnetic resonance imaging, computed tomography [CT], or other advanced imaging) of the spine in patients with non-specific acute low back pain and without red flags. Choosing Wisely Initiative ABIM Foundation. Updated 2014. Accessed November 19, 2022. <https://www.choosingwisely.org/clinician-lists/american-association-neurological-surgeons-imaging-for-nonspecific-acute-low-back-pain/>
9. Allegri M, Montella S, Salici F, et al. Mechanisms of low back pain: a guide for diagnosis and therapy. *F1000Res*. 2016;5doi:10.12688/f1000research.8105.2
10. North American Spine Society. Five things physicians and patients should question: Don't use electromyography (EMG) and nerve conduction studies (NCS) to determine the cause of axial lumbar, thoracic or cervical spine pain. Choosing Wisely Initiative ABIM Foundation. Updated 2019. Accessed December 1, 2022. <https://www.choosingwisely.org/clinician-lists/nass-emg-nerve-conduction-studies-to-determine-cause-of-spine-pain/>
11. American College of Radiology. ACR Appropriateness Criteria® Back Pain—Child. American College of Radiology (ACR). Updated 2016. Accessed December 1, 2022. <https://acsearch.acr.org/docs/3099011/Narrative/>
12. Bernstein RM, Cozen H. Evaluation of back pain in children and adolescents. *Am Fam Physician*. Dec 1 2007;76(11):1669-76.

13. Feldman DS, Straight JJ, Badra MI, Mohaideen A, Madan SS. Evaluation of an algorithmic approach to pediatric back pain. *J Pediatr Orthop*. May-Jun 2006;26(3):353-7. doi:10.1097/01.bpo.0000214928.25809.f9
14. Rao D, Scuderi G, Scuderi C, Grewal R, Sandhu SJ. The Use of Imaging in Management of Patients with Low Back Pain. *J Clin Imaging Sci*. 2018;8:30. doi:10.4103/jcis.JCIS\_16\_18
15. American College of Radiology. ACR Appropriateness Criteria® Low Back Pain. American College of Radiology (ACR). Updated 2021. Accessed January 29, 2023. <https://acsearch.acr.org/docs/69483/Narrative/>
16. Villavicencio AT, Burneikiene S. Elements of the Pre-Operative Workup, Case Examples. *Pain Medicine*. 2006;7(suppl\_1):S35-S46. doi:10.1111/j.1526-4637.2006.00121.x
17. Carayannopoulos A, Han A, Telfeian AE, Scarfo K. Preoperative surveillance thoracic MRI for thoracic dorsal column stimulation: case series. *IPM Reports*. 2019;3(1):1-7. doi:10.36076/pmcr.2019/3/1
18. Corona-Cedillo R, Saavedra-Navarrete MT, Espinoza-Garcia JJ, Mendoza-Aguilar AN, Ternovoy SK, Roldan-Valadez E. Imaging Assessment of the Postoperative Spine: An Updated Pictorial Review of Selected Complications. *Biomed Res Int*. 2021;2021:9940001. doi:10.1155/2021/9940001
19. Fisher BM, Cowles S, Matulich JR, Evanson BG, Vega D, Dissanaik S. Is magnetic resonance imaging in addition to a computed tomographic scan necessary to identify clinically significant cervical spine injuries in obtunded blunt trauma patients? *Am J Surg*. Dec 2013;206(6):987-93; discussion 993-4. doi:10.1016/j.amjsurg.2013.08.021
20. Choi BW, Song KJ, Chang H. Ossification of the posterior longitudinal ligament: a review of literature. *Asian Spine J*. Dec 2011;5(4):267-76. doi:10.4184/asj.2011.5.4.267
21. Behrbalk E, Salame K, Regev GJ, Keynan O, Boszczyk B, Lidar Z. Delayed diagnosis of cervical spondylotic myelopathy by primary care physicians. *Neurosurg Focus*. Jul 2013;35(1):E1. doi:10.3171/2013.3.Focus1374
22. Davies BM, Mowforth OD, Smith EK, Kotter MR. Degenerative cervical myelopathy. *Bmj*. Feb 22 2018;360:k186. doi:10.1136/bmj.k186
23. Sarbu N, Lolli V, Smirniotopoulos JG. Magnetic resonance imaging in myelopathy: a pictorial review. *Clin Imaging*. Sep-Oct 2019;57:56-68. doi:10.1016/j.clinimag.2019.05.002
24. de Oliveira Vilaça C, Orsini M, Leite MA, et al. Cervical Spondylotic Myelopathy: What the Neurologist Should Know. *Neurol Int*. Nov 2 2016;8(4):6330. doi:10.4081/ni.2016.6330
25. American College of Radiology. ACR Appropriateness Criteria® Myelopathy. American College of Radiology (ACR). Updated 2020. Accessed January 29, 2023. <https://acsearch.acr.org/docs/69484/Narrative/>
26. Consortium of Multiple Sclerosis Centers. 2018 MRI Protocol and Clinical Guidelines for MS. Consortium of Multiple Sclerosis Centers (CMSC). Updated May 22, 2018. Accessed January 23, 2023. [https://www.mscares.org/page/MRI\\_protocol](https://www.mscares.org/page/MRI_protocol)
27. Filippi M, Rocca MA, Ciccarelli O, et al. MRI criteria for the diagnosis of multiple sclerosis: MAGNIMS consensus guidelines. *Lancet Neurol*. Mar 2016;15(3):292-303. doi:10.1016/s1474-4422(15)00393-2

28. Kaunzner UW, Gauthier SA. MRI in the assessment and monitoring of multiple sclerosis: an update on best practice. *Ther Adv Neurol Disord*. Jun 2017;10(6):247-261. doi:10.1177/1756285617708911
29. Wingerchuk DM, Banwell B, Bennett JL, et al. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology*. Jul 14 2015;85(2):177-89. doi:10.1212/wnl.0000000000001729
30. American College of Radiology. ACR Appropriateness Criteria® Suspected Spine Trauma American College of Radiology. Updated 2018. Accessed December 1, 2022. <https://acsearch.acr.org/docs/69359/Narrative/>
31. American College of Radiology. ACR Appropriateness Criteria® Inflammatory Back Pain: Known or Suspected Axial Spondyloarthritis. American College of Radiology (ACR). Updated 2021. Accessed December 1, 2022. <https://acsearch.acr.org/docs/3094107/Narrative/>
32. Koivikko MP, Koskinen SK. MRI of cervical spine injuries complicating ankylosing spondylitis. *Skeletal Radiol*. Sep 2008;37(9):813-9. doi:10.1007/s00256-008-0484-x
33. Taljanovic MS, Hunter TB, Wisneski RJ, et al. Imaging characteristics of diffuse idiopathic skeletal hyperostosis with an emphasis on acute spinal fractures: review. *AJR Am J Roentgenol*. Sep 2009;193(3 Suppl):S10-9, Quiz S20-4. doi:10.2214/ajr.07.7102
34. American College of Radiology. ACR Appropriateness Criteria® Management of Vertebral Compression Fractures. American College of Radiology. Updated 2022. Accessed December 1, 2022. <https://acsearch.acr.org/docs/70545/Narrative/>
35. Kumar Y, Hayashi D. Role of magnetic resonance imaging in acute spinal trauma: a pictorial review. *BMC Musculoskelet Disord*. Jul 22 2016;17:310. doi:10.1186/s12891-016-1169-6
36. Kim YS, Han IH, Lee IS, Lee JS, Choi BK. Imaging findings of solitary spinal bony lesions and the differential diagnosis of benign and malignant lesions. *J Korean Neurosurg Soc*. 2012;52(2):126-132. doi:10.3340/jkns.2012.52.2.126
37. Roberts CC, Daffner RH, Weissman BN, et al. ACR appropriateness criteria on metastatic bone disease. *J Am Coll Radiol*. Jun 2010;7(6):400-9. doi:10.1016/j.jacr.2010.02.015
38. McDonald MA, Kirsch CFE, Amin BY, et al. ACR Appropriateness Criteria(®) Cervical Neck Pain or Cervical Radiculopathy. *J Am Coll Radiol*. May 2019;16(5s):S57-s76. doi:10.1016/j.jacr.2019.02.023
39. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Central Nervous System Cancers Version 2.2022. National Comprehensive Cancer Network (NCCN). Updated September 29, 2022. Accessed January 23, 2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/cns.pdf](https://www.nccn.org/professionals/physician_gls/pdf/cns.pdf)
40. Alexandru D, So W. Evaluation and management of vertebral compression fractures. *Perm J*. Fall 2012;16(4):46-51. doi:10.7812/tpp/12-037
41. Lener S, Hartmann S, Barbagallo GMV, Certo F, Thomé C, Tschugg A. Management of spinal infection: a review of the literature. *Acta Neurochir (Wien)*. Mar 2018;160(3):487-496. doi:10.1007/s00701-018-3467-2
42. American College of Radiology. ACR Appropriateness Criteria® Suspected Spine Infection. American College of Radiology (ACR). Updated 2021. Accessed December 1, 2022. <https://acsearch.acr.org/docs/3148734/Narrative/>

43. Bond A, Manian FA. Spinal Epidural Abscess: A Review with Special Emphasis on Earlier Diagnosis. *Biomed Res Int*. 2016;2016:1614328. doi:10.1155/2016/1614328
44. Berbari EF, Kanj SS, Kowalski TJ, et al. 2015 Infectious Diseases Society of America (IDSA) Clinical Practice Guidelines for the Diagnosis and Treatment of Native Vertebral Osteomyelitis in Adults. *Clin Infect Dis*. Sep 15 2015;61(6):e26-46. doi:10.1093/cid/civ482
45. Düz B, Gocmen S, Secer HI, Basal S, Gönül E. Tethered cord syndrome in adulthood. *J Spinal Cord Med*. 2008;31(3):272-8. doi:10.1080/10790268.2008.11760722
46. Milhorat TH, Bolognese PA, Nishikawa M, et al. Association of Chiari malformation type I and tethered cord syndrome: preliminary results of sectioning filum terminale. *Surg Neurol*. Jul 2009;72(1):20-35. doi:10.1016/j.surneu.2009.03.008
47. Zalatio O. Tethered Spinal Cord Syndrome. American Association of Neurological Surgeons (AANS). Accessed December 1, 2022. <https://www.aans.org/Patients/Neurosurgical-Conditions-and-Treatments/Tethered-Spinal-Cord-Syndrome>
48. Whitson WJ, Lane JR, Bauer DF, Durham SR. A prospective natural history study of nonoperatively managed Chiari I malformation: does follow-up MRI surveillance alter surgical decision making? *J Neurosurg Pediatr*. Aug 2015;16(2):159-66. doi:10.3171/2014.12.Peds14301
49. Timpone VM, Patel SH. MRI of a syrinx: is contrast material always necessary? *AJR Am J Roentgenol*. May 2015;204(5):1082-5. doi:10.2214/ajr.14.13310
50. American College of Radiology. ACR Appropriateness Criteria® Headache. American College of Radiology. Updated 2022. Accessed January 23, 2023. <https://acsearch.acr.org/docs/69482/Narrative/>
51. American College of Radiology. ACR Appropriateness Criteria® Headache-Child. American College of Radiology. Updated 2017. Accessed December 1, 2022. <https://acsearch.acr.org/docs/69439/Narrative/>
52. Strahle J, Smith BW, Martinez M, et al. The association between Chiari malformation Type I, spinal syrinx, and scoliosis. *J Neurosurg Pediatr*. Jun 2015;15(6):607-11. doi:10.3171/2014.11.Peds14135
53. Juvenile Scoliosis. Scoliosis Research Society (SRS). Accessed December 1, 2022. <https://www.srs.org/professionals/online-education-and-resources/conditions-and-treatments/juvenile-scoliosis>
54. American College of Radiology. ACR Appropriateness Criteria® Scoliosis-Child. American College of Radiology. Updated 2018. Accessed December 1, 2022. <https://acsearch.acr.org/docs/3101564/Narrative/>
55. Trenga AP, Singla A, Feger MA, Abel MF. Patterns of congenital bony spinal deformity and associated neural anomalies on X-ray and magnetic resonance imaging. *J Child Orthop*. Aug 2016;10(4):343-52. doi:10.1007/s11832-016-0752-6
56. Ozturk C, Karadereler S, Ornek I, Enercan M, Ganiyusufoglu K, Hamzaoglu A. The role of routine magnetic resonance imaging in the preoperative evaluation of adolescent idiopathic scoliosis. *Int Orthop*. Apr 2010;34(4):543-6. doi:10.1007/s00264-009-0817-y
57. Strahle J, Muraszko KM, Kapurch J, Bapuraj JR, Garton HJ, Maher CO. Chiari malformation Type I and syrinx in children undergoing magnetic resonance imaging. *J Neurosurg Pediatr*. Aug 2011;8(2):205-13. doi:10.3171/2011.5.Peds1121

58. Radic JAE, Cochrane DD. Choosing Wisely Canada: Pediatric Neurosurgery Recommendations. *Paediatr Child Health*. Sep 2018;23(6):383-387. doi:10.1093/pch/pxy012
59. Hertzler DA, 2nd, DePowell JJ, Stevenson CB, Mangano FT. Tethered cord syndrome: a review of the literature from embryology to adult presentation. *Neurosurg Focus*. Jul 2010;29(1):E1. doi:10.3171/2010.3.Focus1079
60. Shah LM, Salzman KL. Imaging of spinal metastatic disease. *Int J Surg Oncol*. 2011;2011:769753. doi:10.1155/2011/769753
61. Last AR, Hulbert K. Chronic low back pain: evaluation and management. *Am Fam Physician*. Jun 15 2009;79(12):1067-74.
62. Stanford Medicine. Gait Abnormalities. Stanford University. Accessed January 23, 2023. <https://stanfordmedicine25.stanford.edu/the25/gait.html>
63. Haynes KB, Wimberly RL, VanPelt JM, Jo CH, Riccio AI, Delgado MR. Toe Walking: A Neurological Perspective After Referral From Pediatric Orthopaedic Surgeons. *J Pediatr Orthop*. Mar 2018;38(3):152-156. doi:10.1097/bpo.0000000000001115
64. Chhetri SK, Gow D, Shaunak S, Varma A. Clinical assessment of the sensory ataxias; diagnostic algorithm with illustrative cases. *Pract Neurol*. Aug 2014;14(4):242-51. doi:10.1136/practneurol-2013-000764
65. Foster H, Drummond P, Jandial S, Clinch J, Wood M, Driscoll S. Evaluation of gait disorders in children. BMJ Best Practice. Updated February 23, 2021. Accessed January 23, 2023. <https://bestpractice.bmj.com/topics/en-us/709>
66. Marshall FJ. Approach to the elderly patient with gait disturbance. *Neurol Clin Pract*. Jun 2012;2(2):103-111. doi:10.1212/CPJ.0b013e31825a7823
67. Pirker W, Katzenschlager R. Gait disorders in adults and the elderly : A clinical guide. *Wien Klin Wochenschr*. Feb 2017;129(3-4):81-95. doi:10.1007/s00508-016-1096-4
68. Vitzthum HE, Dalitz K. Analysis of five specific scores for cervical spondylogenic myelopathy. *Eur Spine J*. Dec 2007;16(12):2096-103. doi:10.1007/s00586-007-0512-x
69. Ziu E, Viswanathan VK, Mesfin FB. Spinal Metastasis. StatPearls Publishing. Updated August 22, 2022. Accessed December 1, 2022. <https://www.ncbi.nlm.nih.gov/books/NBK441950/>
70. Borofsky S, Levy LM. Neurofibromatosis: types 1 and 2. *AJNR Am J Neuroradiol*. Dec 2013;34(12):2250-1. doi:10.3174/ajnr.A3534
71. Evans DGR, Salvador H, Chang VY, et al. Cancer and Central Nervous System Tumor Surveillance in Pediatric Neurofibromatosis 2 and Related Disorders. *Clin Cancer Res*. Jun 15 2017;23(12):e54-e61. doi:10.1158/1078-0432.Ccr-17-0590
72. Krueger DA, Northrup H. Tuberous sclerosis complex surveillance and management: recommendations of the 2012 International Tuberous Sclerosis Complex Consensus Conference. *Pediatr Neurol*. Oct 2013;49(4):255-65. doi:10.1016/j.pediatrneurol.2013.08.002
73. Varshney N, Kebede AA, Owusu-Dapaah H, Lather J, Kaushik M, Bhullar JS. A Review of Von Hippel-Lindau Syndrome. *J Kidney Cancer VHL*. 2017;4(3):20-29. doi:10.15586/jkcvhl.2017.88
74. Comi AM. Presentation, diagnosis, pathophysiology, and treatment of the neurological features of Sturge-Weber syndrome. *Neurologist*. Jul 2011;17(4):179-84. doi:10.1097/NRL.0b013e318220c5b6



75. Ahmed A. MRI features of disseminated 'drop metastases'. *S Afr Med J*. Jul 2008;98(7):522-3.
76. Batool A, Kasi A. Leptomeningeal Carcinomatosis. StatPearls Publishing  
Copyright © 2022, StatPearls Publishing LLC. Updated April 5, 2022. Accessed December 1, 2022. <https://www.ncbi.nlm.nih.gov/books/NBK499862/>

## ADDITIONAL RESOURCES

1. Cohen SP, Gupta A, Strassels SA, et al. Effect of MRI on treatment results or decision making in patients with lumbosacral radiculopathy referred for epidural steroid injections: a multicenter, randomized controlled trial. *Arch Intern Med*. Jan 23 2012;172(2):134-42. doi:10.1001/archinternmed.2011.593
2. D'Alessandro DM. Does This Sacral Dimple Need to be Evaluated? *PediatricEducation.org™*. Updated July 20, 2009. Accessed December 1, 2022 November 10, 2021. <https://pediatriceducation.org/2009/07/20/does-this-sacral-dimple-need-to-be-evaluated/>
3. Goh C, Phal PM, Desmond PM. Neuroimaging in acute transverse myelitis. *Neuroimaging Clin N Am*. Nov 2011;21(4):951-73. x. doi:10.1016/j.nic.2011.07.010
4. Hou X, Sun C, Liu X, et al. Clinical Features of Thoracic Spinal Stenosis associated Myelopathy: A Retrospective Analysis of 427 Cases. *Clin Spine Surg*. Mar 2016;29(2):86-9. doi:10.1097/bsd.0000000000000081
5. Magge SN, Smyth MD, Governale LS, et al. Idiopathic syrinx in the pediatric population: a combined center experience. *J Neurosurg-Pediatr*. Jan 2011;7(1):30-6. doi:10.3171/2010.10.Peds1057
6. North American Spine Society. Five things physicians and patients should question. Choosing Wisely Initiative ABIM Foundation. Updated 201921. Accessed December 1 November 9, 20221. <https://www.choosingwisely.org/societies/north-american-spine-society/>
7. Papanastassiou ID, Gerochristou M, Aghayev K, Vrionis FD. Defining the indications, types and biomaterials of corpectomy cages in the thoracolumbar spine. *Expert Rev Med Devices*. Mar 2013;10(2):269-79. doi:10.1586/erd.12.79

## POLICY HISTORY

<u>Date</u>	<u>Summary</u>
<u>May 2023</u>	<ul style="list-style-type: none"> <li>• <u>Updated references</u></li> <li>• <u>Updated background section</u></li> <li>• <u>Clarified pathological reflexes</u></li> <li>• <u>General Information moved to beginning of guideline with added statement on clinical indications not addressed in this guideline</u></li> <li>• <u>Added statement regarding further evaluation of indeterminate findings on prior imaging</u></li> <li>• <u>Clarified cerebellar ataxia in gait table</u></li> <li>• <u>Removed radicular pain and malaise from Isolated Back Pain in the Pediatric population: Red flags</u></li> <li>• <u>Removed Additional Resources</u></li> </ul>
<u>March 2022</u>	<p><u>Added</u></p> <ul style="list-style-type: none"> <li>• <u>Combination request for overlapping body part statement</u></li> <li>• <u>Clarified muscle weakness not related to plexopathy or peripheral neuropathy</u></li> <li>• <u>Clarified bowel and bladder dysfunction – not related to an inherent bowel or bladder problem</u></li> <li>• <u>Clarified combination MS for cervical and/or thoracic spine combination requests</u></li> <li>• <u>Descriptions for tethered cord</u></li> <li>• <u>Background section of Drop Metastases</u></li> <li>• <u>Background section of Leptomeningeal Carcinomatosis</u></li> <li>• <u>Clarified toe walking in pediatric patient with myelopathy for thoracic spine</u></li> </ul> <p><u>Removed</u></p> <ul style="list-style-type: none"> <li>• <u>Removed from combination section syrinx and syringomyelia and added subsection for cervical and thoracic spine section</u></li> <li>• <u>Removed pediatric back pain from the total spine combination section</u></li> </ul>
<u>November 2021</u>	<p><del>— Added +0698T</del></p>
<u>April 2021</u>	<p><del>— Added/modified</del></p> <ul style="list-style-type: none"> <li><del>— <u>Modified section on neurological deficits</u></del></li> <li><del>— <u>Back pain in a child added/modified red flags</u></del></li> <li><del>— <u>Gait table in background</u></del></li> <li><del>— <u>Post-surgical modified/clarified surgical criteria for combination exams</u></del></li> <li><del>— <u>Removed myelopathy combination studies</u></del></li> <li><del>— <u>Updated/added MS Criteria</u></del></li> <li><del>— <u>Combination section for initial imaging and follow-up</u></del></li> </ul>



	<ul style="list-style-type: none"> <li>— <del>Added pediatric MS</del></li> <li>— <del>Modified known tumor imaging into primary and metastatic disease</del></li> <li>— <del>Added toe walking for pediatric patients</del></li> <li>— <del>Modified Combination exam wording</del></li> </ul>
<u>May 2020</u>	<ul style="list-style-type: none"> <li>— <del>Added</del> <ul style="list-style-type: none"> <li>— <del>For evaluation of neurologic deficits when new deficits are present</del></li> <li>— <del>Removed pars defect section</del></li> <li>— <del>Added ankylosing spondylitis for evaluation of trauma/acute injury</del></li> <li>— <del>Added follow up of osteoporotic fracture from metastatic disease</del></li> <li>— <del>Added transverse myelitis</del></li> <li>— <del>Modified Initial imaging of new or increasing non-traumatic back pain or radiculopathy or back pain that occurs at night and wakes the patient from sleep with known active cancer and a tumor that tends to metastasize to the spine</del></li> <li>— <del>Added Imaging of Ossification of the Posterior Longitudinal Ligament (OPPL)</del></li> <li>— <del>Added Osteopathic Manipulative medicine to conservative care therapy</del></li> </ul> </li> </ul>
<u>June 2019</u>	<ul style="list-style-type: none"> <li>— <del>Added:</del> <ul style="list-style-type: none"> <li>— <del>new or worsening objective neuro deficits for chronic and acute back pain</del></li> <li>— <del>CSF leak</del></li> <li>— <del>last 6 months for allowable post op f/u period and removed EMG comment</del></li> <li>— <del>red flags specifically for peds back pain and pain related to malignancy, infection, inflammation</del></li> <li>— <del>new sections: pars defect; compression fractures; congenital abnormalities including section on scoliosis and vertebral anomalies in children w/back pain;</del></li> <li>— <del>For combination studies cervical/thoracic/lumbar added drop metastasis, tumor evaluation for neurocutaneous syndromes, and abnormalities associated w/Arnold Chiari, as well as separate indication for tethered cord or spinal dysraphism</del></li> <li>— <del>Myelopathy</del></li> </ul> </li> </ul>

	<ul style="list-style-type: none"> <li><del>— <u>Pre op for spinal cord stimulator</u></del></li> <li><del>— <u>Evaluation of MS including NMO disorders and recurrent transverse myelitis</u></del></li> <li><del>— <u>Back pain in cancer patients with known active cancer in tumors that tend to metastasize</u></del></li> <li><del>— <u>Expanded on tethered cord in Other Indications for Imaging and added content on sacral dimple</u></del></li> <li><del>— </del></li> </ul>
--	---

## Reviewed / Approved by NIA Clinical Guideline Committee

**Disclaimer:** National Imaging Associates, Inc. (NIA) authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Evolent Health LLC subsidiaries including, but not limited to, National Imaging Associates (“NIA”). The policies constitute only the reimbursement and coverage guidelines of NIA. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. NIA reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.



### **GENERAL INFORMATION**

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

**Disclaimer:** Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates ("Magellan"). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.