

AmeriHealth Caritas Louisiana

National Imaging Associates, Inc.	
Clinical guidelines HEART (Cardiac) PET	Original Date: July 1999
CPT Codes: 78459, 78491, 78492, +78434	Last Revised Date: February 2022 May 2023
Guideline Number: NIA_CG_072	Implementation Date: January 202324

GENERAL INFORMATION

- *It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.*
- ***Where a specific clinical indication is not directly addressed in this guideline, medical necessity determination will be made based on widely accepted standard of care criteria. These criteria are supported by evidence-based or peer-reviewed sources such as medical literature, societal guidelines and state/national recommendations.***

This guideline is for stress imaging, specifically Heart (Cardiac) PET imaging, with appropriate preference for suitable alternatives, such as stress echocardiography (SE) or myocardial perfusion imaging (MPI), when more suitable, unless otherwise stated (refer to [Background section](#)).

INDICATIONS FOR HEART PET¹⁻⁴

SUSPECTED CAD (When neither SE nor MPI have provided or are expected to provide optimal imaging)

- **Symptomatic patients without known CAD (use [Diamond Forrester Table](#))**
 - Low or intermediate pretest probability and unable to exercise (SE diversion not required)
 - High pretest probability (SE diversion not required)
 - Repeat testing in a patient with new or worsening symptoms and negative result at least one year ago **AND** meets one of the criteria above
- **Asymptomatic patients without known CAD (SE diversion not required)**

*National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.

- Previously unevaluated ECG evidence of possible myocardial ischemia including substantial ischemic ST segment or T wave abnormalities (~~(see section in Background)~~)
- Previously unevaluated pathologic Q waves (~~(see section in Background)~~)
- Unevaluated complete left bundle branch block

ABNORMAL CALCIUM SCORES (CAC)¹⁻⁵ ~~3, 5-8~~ (When neither SE nor MPI have provided, or are expected to provide, optimal imaging)

- ASYMPTOMATIC patient with a calcium score >400, not previously evaluated
- SYMPTOMATIC patient with prior CAC ≥100

INCONCLUSIVE CAD EVALUATION ~~WITHIN THE PAST 2 YEARS~~ AND OBSTRUCTIVE CAD REMAINS A CONCERN (When neither SE nor MPI have provided, or are expected to provide, optimal imaging)

- Exercise stress ECG with low-risk Duke treadmill score (≥5) ([see section in Background](#)) but patient's current symptoms indicate an intermediate or high pretest probability (SE diversion not required for high pretest probability)
- Exercise stress ECG with an intermediate Duke treadmill score (SE diversion not required for symptoms consistent with high pretest probability)
- Inconclusive/borderline coronary computed tomography angiography (CCTA) (e.g., 40 - 70% lesions)
- Non-diagnostic exercise stress test with physical inability to achieve target heart rate (THR) (SE diversion not required)
- An intermediate evaluation by prior stress imaging (~~within the past 2 years~~) (SE diversion not required)
- Coronary stenosis of unclear significance on previous coronary angiography³

FOLLOW-UP OF PATIENT'S POST CORONARY REVASCULARIZATION (PCI or CABG) (When neither SE nor MPI have provided, or are expected to provide, optimal imaging)³

- **Asymptomatic, follow-up stress imaging** at a minimum of 2 years post coronary artery bypass grafting (CABG), or percutaneous coronary intervention (PCI), (whichever is later), is appropriate only for patients with a history of silent ischemia or a history of a prior left main stent

OR

- For patients with high occupational risk (e.g., associated with public safety, airline and boat pilots, bus and train drivers, bridge and tunnel workers/toll collectors, police officers, and firefighters)

New, recurrent, or worsening symptoms post coronary revascularization are an indication for stress imaging, if it will alter management (SE diversion not required for typical anginal symptoms or symptoms documented to be similar to those prior to revascularization)

FOLLOW-UP OF KNOWN³ CAD (When neither SE nor MPI have provided, or are expected to provide, optimal imaging)

- **Follow-up of asymptomatic or stable symptoms** when last invasive or non-invasive assessment of coronary disease showed hemodynamically significant CAD (ischemia on stress test or FFR ≤ 0.80 or **significant** stenosis ~~greater than or equal to 70% of in~~ a major vessel), ($\geq 50\%$ left main coronary artery or $\geq 70\%$ LAD, LCX or RCA), over two years ago, without intervening coronary revascularization is an appropriate indication for stress imaging in patients if it will alter management

SPECIAL DIAGNOSTIC CONDITIONS REQUIRING CORONARY EVALUATION (When neither SE nor MPI have provided, or are expected to provide, optimal imaging)

- Prior acute coronary syndrome (as documented in MD notes), without subsequent invasive or non-invasive coronary evaluation
- Newly diagnosed systolic heart failure or diastolic heart failure, *with reasonable suspicion of cardiac ischemia (prior events, risk factors)*, unless invasive coronary angiography is immediately planned^{6-8, 9, 10}
- Reduced LVEF $\leq 50\%$ requiring myocardial viability assessment to assist with decisions regarding coronary revascularization. (Diversion from PET not required when LVEF less than or equal to 40%)^{6, 7, 9, 9-11}
- Ventricular arrhythmias
 - Sustained ventricular tachycardia (VT) > 100 bpm, ventricular fibrillation (VF), or exercise-induced VT, when invasive coronary arteriography is not the immediately planned test^{10, 12}
 - Nonsustained VT, multiple episodes, each ≥ 3 beats at ≥ 100 bpm, frequent PVC's (defined as greater than or equal to 30/hour on remote monitoring) without known cause or associated cardiac pathology, when an exercise ECG cannot be performed
- Prior to Class IC antiarrhythmic drug initiation (Propafenone or Flecainide), as well as annually in intermediate and high global risk patients (SE diversion not required)^{14, 3}
- Assessment of hemodynamic significance of one of the following documented conditions^{12, 14}:
 - Anomalous coronary arteries^{13, 5}
 - Muscle bridging of coronary artery^{3, 14, 3, 16}
- Coronary aneurysms in Kawasaki's disease^{15, 7} or due to atherosclerosis
- Following radiation therapy to the anterior or left chest, at 5 years post initiation and every 5 years thereafter^{16, 18}

- ~~Cardiac Sarcoidosis~~¹⁷⁻¹⁹
- To diagnose microvascular dysfunction in patients with persistent stable anginal chest pain with suspected ischemia and nonobstructive coronary artery disease (INOCA), as documented in provider notes (no MPI diversion required).
- Cardiac Sarcoidosis¹⁹⁻²¹ (may be approved as a combination study with MPI for the evaluation and treatment of sarcoidosis)²²
 - Evaluation and therapy monitoring in patients with sarcoidosis, after documentation of suspected cardiac involvement by echo or ECG, when CMR has not been performed
 - Evaluation of suspected cardiac sarcoid, after CMR has shown equivocal or negative findings in the setting of a high clinical suspicion²¹⁹
 - Evaluation of CMR findings showing highly probable cardiac sarcoidosis, when PET could serve to identify inflammation and the consequent potential role for immunosuppressive therapy²¹⁹
 - Initial and follow-up PET in monitoring therapy for cardiac sarcoid with immunosuppressive therapy, typically about 4 times over 2 years
- **Infective Endocarditis**
 - In suspected infective endocarditis with moderate to high probability (i.e., staph bacteremia, fungemia, prosthetic heart valve, or intracardiac device), when TTE and TEE have been inconclusive with respect to diagnosis of infective endocarditis or characterization of paravalvular invasive complications^{20, 2423, 24}
- **Aortitis**
 - For diagnosis and surveillance of Aortitis, PET/CT or PET/MRI[†] hybrid imaging²²²⁵
[†]**NOTE:** If PET/MR study is requested, there is no specific CPT Code for this imaging study and a Health Plan review will be required.

PRIOR TO ELECTIVE NON-CARDIAC SURGERY (When neither SE nor MPI have provided or are expected to provide optimal imaging)

- An intermediate or high risk surgery with of one or more risk factors (see below), AND documentation of an inability to walk (or <4 METs) AND there has not been an imaging stress test within 1 year²³⁻²⁵²⁶⁻²⁸
 - **Risk factors:** history of ischemic heart disease, history of congestive heart failure, history of cerebrovascular disease, preoperative treatment with insulin, and preoperative serum creatinine >2.0 mg/dL.
 - **Surgical Risk:**
 - **High risk surgery:** Aortic and other major vascular surgery, peripheral vascular surgery, anticipated prolonged surgical procedures associated with large fluid shifts and/or blood loss
 - **Intermediate risk surgery:** Carotid endarterectomy, head and neck surgery, intraperitoneal and intrathoracic surgery, orthopedic surgery, prostate surgery

- **Low risk surgery:** Endoscopic procedures, superficial procedure, cataract surgery, breast surgery
- Planning for any organ or stem cell transplantation is an indication for preoperative stress imaging, if there has not been a conclusive stress evaluation, CTA, or heart catheterization within the past year, at the discretion of the transplant service^{26,29}

POST CARDIAC TRANSPLANT (SE diversion not required)^{27,30}

- Annually, for the first five years post cardiac transplantation, in a patient not undergoing invasive coronary arteriography
- After the first five years post cardiac transplantation, patients with documented transplant coronary vasculopathy can be screened annually if invasive coronary arteriography is not planned

BACKGROUND^{28, 29, 31, 32}

PET Scan

- Indicated when all the criteria for MPI are met **AND** there is likely to be equivocal imaging results because of BMI ~~or~~, large breasts or implants, mastectomy, chest wall deformity, pleural or pericardial effusion, or prior thoracic surgery or results of a prior MPI
- Can identify regions of myocardial viability with hibernating myocardium (viable, with poor flow and contractility) by imaging with fluorine-18 (F-18) fluorodeoxyglucose (FDG or 18-FDG) for this purpose.
- Useful in the evaluation of inflammation: e.g., evaluation and therapy monitoring in patients with sarcoidosis, after documentation of cardiac involvement by echo or electrocardiography (ECG), in place of, or subsequent to CMR if needed to help with an uncertain diagnosis

Coronary application of PET includes evaluation of **stable patients without known CAD**, who fall into two categories^{3, 8, 30, 4}

- **Asymptomatic**, for whom global risk of CAD events can be determined from coronary risk factors, using calculators available online (see Websites for [Global Cardiovascular Risk Calculators](#) section).
- **Symptomatic**, for whom we estimate the pretest probability that their chest-related symptoms are due to clinically significant ($\geq 50\%$) CAD (below):

The 3 Types of Chest Pain or Discomfort

- **Typical Angina (Definite)** is defined as including **all 3** characteristics:
 - Substernal chest pain or discomfort with characteristic quality and duration
 - Provoked by exertion or emotional stress

- Relieved by rest and/or nitroglycerine
- **Atypical Angina (Probable)** has only **2** of the above characteristics
- **Nonanginal Chest Pain/Discomfort** has only **0 - 1** of the above characteristics

The medical record should provide enough detail to establish the type of chest pain. From those details, The Pretest Probability of obstructive CAD is estimated from the **Diamond Forrester Table** below, recognizing that in some cases multiple additional coronary risk factors could increase pretest probability^{3,82,3}:

Diamond Forrester Table

Age (Years)	Gender	Typical/Definite Angina Pectoris	Atypical/Probable Angina Pectoris	Nonanginal Chest Pain
≤ 39	Men	Intermediate	Intermediate	Low
	Women	Intermediate	Very low	Very low
40 – 49	Men	High	Intermediate	Intermediate
	Women	Intermediate	Low	Very low
50 – 59	Men	High	Intermediate	Intermediate
	Women	Intermediate	Intermediate	Low
≥ 60	Men	High	Intermediate	Intermediate
	Women	High	Intermediate	Intermediate

- **Very Low:** < 5% pretest probability, usually not requiring stress evaluation
- **Low:** 5 - 10% pretest probability of CAD
- **Intermediate:** 10% - 90% pretest probability of CAD
- **High:** > 90% pretest probability of CAD

OVERVIEW

ECG Stress Test Alone versus Stress Testing with Imaging

Prominent scenarios suitable for an ECG stress test WITHOUT imaging (i.e., exercise treadmill ECG test) require that the patient can exercise for at least 3 minutes of Bruce protocol with achievement of near maximal heart rate AND has an interpretable ECG for ischemia during exercise³:

- The (symptomatic) low or intermediate pretest probability patient who is able to exercise and has an interpretable ECG³
- The patient who is under evaluation for exercise-induced arrhythmia
- The patient who requires an entrance stress test ECG for a cardiac rehab program or for an exercise prescription
- For the evaluation of syncope or presyncope during exertion³⁴³³

~~Duke Exercise ECG Treadmill Score³²~~ Duke Exercise ECG Treadmill Score³⁴

Calculates risk from ECG treadmill alone:

- The equation for calculating the Duke treadmill score (DTS) is: $DTS = \text{exercise time in minutes} - (5 \times \text{ST deviation in mm or 0.1 mV increments}) - (4 \times \text{exercise angina score})$, with angina score being 0 = none, 1 = non-limiting, and 2 = exercise-limiting.
- The score typically ranges from - 25 to + 15. These values correspond to low-risk (with a score of $\geq + 5$), intermediate risk (with scores ranging from - 10 to + 4), and high-risk (with a score of $\leq - 11$) categories.

An uninterpretable baseline ECG includes³²:

- ST segment depression 1 mm or more (not for non-specific ST- T wave changes)
- Ischemic looking T waves; at least 2.5 mm inversions (excluding V1 and V2)
- LVH with repolarization abnormalities, pre-excitation pattern such as WPW, ventricular paced rhythm, or left bundle branch block
- Digitalis use with associated ST segment abnormalities

Previously unevaluated pathologic Q waves (in two contiguous leads) defined as the following:

- > 40 ms (1 mm) wide
- > 2 mm deep
- > 25% of depth of QRS complex

Global Risk of Cardiovascular Disease

Global risk of CAD is defined as the probability of manifesting cardiovascular disease over the next 10 years and refers to **asymptomatic** patients without known cardiovascular disease. It should be determined using one of the risk calculators below. A high risk is considered greater than a 20% risk of a cardiovascular event over the ensuing 10 years. **High global risk by itself generally lacks scientific support as an indication for stress imaging.** There are rare exceptions, such as patients requiring IC antiarrhythmic drugs who might require coronary risk stratification prior to initiation of the drug.

- **CAD Risk—Low**
10-year absolute coronary or cardiovascular risk less than 10%
- **CAD Risk—Moderate**
10-year absolute coronary or cardiovascular risk between 10% and 20%
- **CAD Risk—High**
10-year absolute coronary or cardiovascular risk of greater than 20%

Websites for Global Cardiovascular Risk Calculators*~~33-37~~³⁵⁻³⁹

~~*Patients who have already manifested cardiovascular disease are already at high global risk and are not applicable to the calculators.~~

Risk Calculator	Websites for Online Calculator
Framingham Cardiovascular Risk	https://reference.medscape.com/calculator/framingham-cardiovascular-disease-risk
Reynolds Risk Score Can use if no diabetes Unique for use of family history	http://www.reynoldsriskscore.org/
Pooled Cohort Equation	http://clincalc.com/Cardiology/ASCVD/PooledCohort.aspx?example
ACC/AHA Risk Calculator	http://tools.acc.org/ASCVD-Risk-Estimator/
MESA Risk Calculator With addition of Coronary Artery Calcium Score, for CAD-only risk	https://www.mesa-nhlbi.org/MESACHDRisk/MesaRiskScore/RiskScore.aspx

*Patients who have already manifested cardiovascular disease are already at high global risk and are not applicable to the calculators.

Definitions of Coronary Artery Disease^{2, 8, 392, 4, 6}

Percentage stenosis refers to the reduction in diameter stenosis when angiography is the method and can be estimated or measured using angiography or more accurately measured with intravascular ultrasound (IVUS).

- Coronary artery calcification is a marker of risk, as measured by Agatston score on coronary artery calcium imaging. Its incorporation into global risk can be achieved by using the MESA risk calculator.
- Ischemia-producing disease (also called hemodynamically or functionally significant disease, for which revascularization might be appropriate) generally implies at least one of the following:
 - Suggested by percentage diameter stenosis $\geq 70\%$ by angiography; intermediate lesions are 50 – 69%³⁸⁴⁰
 - For a left main artery, suggested by a percentage stenosis $\geq 50\%$ or minimum lumen cross-sectional area on IVUS ≤ 6 square mm^{8, 392, 41}
 - FFR (fractional flow reserve) ≤ 0.80 for a major vessel³⁹⁴¹
 - Demonstrable ischemic findings on stress testing (ECG or stress imaging), that are at least mild in degree

- A major vessel would be a coronary vessel that would be amenable to revascularization if indicated. This assessment is made based on the diameter of the vessel and/or the extent of myocardial territory served by the vessel.
- FFR (fractional flow reserve) is the distal to proximal pressure ratio across a coronary lesion during maximal hyperemia induced by either intravenous or intracoronary adenosine. Less than or equal to 0.80 is considered a significant reduction in coronary flow.
- Newer technology that estimates FFR from CCTA image is covered under the separate NIA Guideline for FFR-CT.

Anginal Equivalent^{8, 312, 33}

Development of an anginal equivalent (e.g., shortness of breath, fatigue, or weakness) either with or without prior coronary revascularization should be based upon the documentation of reasons to suspect that symptoms other than chest discomfort are not due to other organ systems (e.g., dyspnea due to lung disease, fatigue due to anemia), by presentation of clinical data, such as respiratory rate, oximetry, lung exam, etc. (as well as d-dimer, chest CT(A), and/or PFTs, when appropriate), and then incorporated into the evaluation of coronary artery disease as would chest discomfort. Most syncope per se is not an anginal equivalent.

Abbreviations

ADLs	Activities of daily living
BMI	Body mass index
CABG	Coronary artery bypass grafting
CAC	Coronary artery calcium
CAD	Coronary artery disease
CCTA	Coronary computed tomography angiography
CMR	Cardiac magnetic resonance imaging
CT(A)	Computed tomography (angiography)
DTS	Duke Treadmill Score
ECG	Electrocardiogram
FFR	Fractional flow reserve
IVUS	Intravascular ultrasound
LBBB	Left bundle-branch block
LVEF	Left ventricular ejection fraction
LVH	Left ventricular hypertrophy
MESA	Multi-Ethnic Study of Atherosclerosis
MET	Estimated metabolic equivalent of exercise
MI	Myocardial infarction
MPI	Myocardial perfusion imaging
MR(I)	Magnetic resonance (imaging)
PCI	Percutaneous coronary intervention
PET	Positron emission tomography
PFT	Pulmonary function test
PVCs	Premature ventricular contractions
SE	Stress echocardiography
TEE	Transesophageal echocardiography
THR	Target heart rate
TTE	Transthoracic echocardiography
VF	Ventricular fibrillation
VT	Ventricular tachycardia
WPW	Wolff-Parkinson-White

Policy History

Date	Summary
<u>May 2023</u>	<ul style="list-style-type: none"> • <u>Removed time limitation “within past two years” for further evaluation inconclusive prior CAD evaluation</u> • <u>Added coronary stenosis of unclear significance on previous coronary angiography.</u> • <u>Added indication for evaluation of ischemia and nonobstructive coronary artery disease (INOCA)</u> • <u>Clarified indication for PET/MPI combination study for evaluation of cardiac sarcoidosis</u> • <u>Added statement on clinical indications not addressed in this guideline</u>
February 2022	<ul style="list-style-type: none"> • Moved the sentence regarding utilization of suitable alternatives such as Stress Echocardiography to the General Information section • Clarified “intermediate lesions are 50-69%” for ischemia-producing disease • Clarified evaluation of possible ischemia in newly diagnosed heart failure by stating “<i>with reasonable suspicion of cardiac ischemia (prior events, risk factors, or symptoms and signs)</i>” • Placed Link to Overview Section in General Information • Added stress imaging approval for calcium score > 100 with low to intermediate probability symptoms • Deleted the requirement for diabetes when calcium score > 400 for stress imaging • Added Calcium score section: <ul style="list-style-type: none"> ○ Added stress imaging approval for calcium score > 100 with symptoms consistent with low to intermediate pretest probability • Added reminder <u>(SE diversion not required for CABG)</u> • Changed preoperative guideline to include intermediate risk surgery with one or more risk factors AND documentation of an inability to walk (or <4 METs) AND there has not been an imaging stress test within 1 year • Changed solid organ transplant guideline to include stem cell transplant and “any” organ transplant • Added definition of surgical risk to preop guidelines • In Background section clarified the requirement for description of chest pain by adding sentence “The medical record should provide enough detail to establish the type of chest pain. “ • Added definition of Q waves

	<ul style="list-style-type: none"> Deleted sentence regarding calcium scoring within the Global Risk Section Deleted sentence regarding using calcium score solely for risk stratification Deleted redundant statement on viability Deleted IFR references
March 2021	<ul style="list-style-type: none"> Added annual indication for IC antiarrhythmics Added History of diabetes mellitus, > 40 years old, with calcium score >400
March 2020	<ul style="list-style-type: none"> Added general information section as Introduction which outlines requirements for documentation of pertinent office notes by a licensed clinician, and inclusion of laboratory testing and relevant imaging results for case review Added clarification of repeat testing in a patient with new or worsening symptoms and negative result at least one year prior to include the statement “AND meets one of the criteria above” Added clarification of frequent PVCs under ventricular arrhythmias which states defined as greater than or equal to 30/hour to include “on remote monitoring” Edited indication of planning for solid organ transplantation to remove the requirement of limited functional capacity but maintaining requirement of ≥ 3 listed risk factors Edits to the Background section include the following: <ul style="list-style-type: none"> Indication changed to read as follows: PET is indicated when all the criteria for MPI are met AND There is likely to be equivocal imaging results because of BMI or large breasts or implants or prior thoracic surgery or results of a prior MPI Removed the statement regarding radiation burden Added edits to the Coronary Artery disease definition section Updated and added new references
November 2019	<ul style="list-style-type: none"> Removed CPT code +0482T and replaced with code +78434
August 2019	<ul style="list-style-type: none"> Changes in CAD indications in line with MPI/SE Added infective endocarditis and aortitis indications Removed cardiac neoplasms and masses indication section Added myocardial viability indications Expanded indications for cardiac sarcoidosis as the initial and follow up study

REFERENCES

1. Hendel RC, Berman DS, Di Carli MF, et al. ACCF/ASNC/ACR/AHA/ASE/SCCT/SCMR/SNM 2009 Appropriate Use Criteria for Cardiac Radionuclide Imaging: A Report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, the American Society of Nuclear Cardiology, the American College of Radiology, the American Heart Association, the American Society of Echocardiography, the Society of Cardiovascular Computed Tomography, the Society for Cardiovascular Magnetic Resonance, and the Society of Nuclear Medicine. *J Am Coll Cardiol*. Jun 9 2009;53(23):2201-29. doi:10.1016/j.jacc.2009.02.013
2. Fihn SD, Gardin JM, Abrams J, et al. 2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology Foundation/American Heart Association task force on practice guidelines, and the American College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *Circulation*. Dec 18 2012;126(25):e354-471. doi:10.1161/CIR.0b013e318277d6a0
3. Wolk MJ, Bailey SR, Doherty JU, et al. ACCF/AHA/ASE/ASNC/HFSA/HRS/SCAI/SCCT/SCMR/STS 2013 multimodality appropriate use criteria for the detection and risk assessment of stable ischemic heart disease: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. Feb 4 2014;63(4):380-406. doi:10.1016/j.jacc.2013.11.009
4. Montalescot G, Sechtem U, Achenbach S, et al. 2013 ESC guidelines on the management of stable coronary artery disease: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *Eur Heart J*. Oct 2013;34(38):2949-3003. doi:10.1093/eurheartj/eh296
5. Gulati M, Levy PD, Mukherjee D, et al. 2021 AHA/ACC/ASE/CHEST/SAEM/SCCT/SCMR Guideline for the Evaluation and Diagnosis of Chest Pain: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Cardiovasc Comput Tomogr*. Dec 01 2021;doi:10.1016/j.jcct.2021.11.009
26. Patel MR, Calhoon JH, Dehmer GJ, et al. ACC/AATS/AHA/ASE/ASNC/SCAI/SCCT/STS 2017 Appropriate Use Criteria for Coronary Revascularization in Patients With Stable Ischemic Heart Disease: A Report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. May 2 2017;69(17):2212-2241. doi:10.1016/j.jacc.2017.02.001

- ~~7. 3. Wolk MJ, Bailey SR, Doherty JU, et al. ACCF/AHA/ASE/ASNC/HFSA/HRS/SCAI/SCCT/SCMR/STS 2013 multimodality appropriate use criteria for the detection and risk assessment of stable ischemic heart disease: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. Feb 4 2014;63(4):380-406. doi:10.1016/j.jacc.2013.11.009~~
4. Budoff MJ, Raggi P, Beller GA, et al. Noninvasive Cardiovascular Risk Assessment of the Asymptomatic Diabetic Patient: The Imaging Council of the American College of Cardiology. *JACC Cardiovasc Imaging*. Feb 2016;9(2):176-92. doi:10.1016/j.jcmg.2015.11.011
58. Aguilar-Salinas CA, Rojas R, Gomez-Perez FJ, et al. Analysis of the agreement between the World Health Organization criteria and the National Cholesterol Education Program-III definition of the metabolic syndrome: results from a population-based survey. *Diabetes Care*. May 2003;26(5):1635.
69. Patel MR, White RD, Abbara S, et al. 2013 ACCF/ACR/ASE/ASNC/SCCT/SCMR appropriate utilization of cardiovascular imaging in heart failure: a joint report of the American College of Radiology Appropriateness Criteria Committee and the American College of Cardiology Foundation Appropriate Use Criteria Task Force. *J Am Coll Cardiol*. May 28 2013;61(21):2207-31. doi:10.1016/j.jacc.2013.02.005
710. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. Oct 15 2013;62(16):e147-239. doi:10.1016/j.jacc.2013.05.019
- ~~8. Fihn SD, Gardin JM, Abrams J, et al. 2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology Foundation/American Heart Association task force on practice guidelines, and the American College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *Circulation*. Dec 18 2012;126(25):e354-471. doi:10.1161/CIR.0b013e318277d6a0~~
911. Tsai JP, Yun CH, Wu TH, et al. A meta-analysis comparing SPECT with PET for the assessment of myocardial viability in patients with coronary artery disease. *Nucl Med Commun*. Sep 2014;35(9):947-54. doi:10.1097/mnm.0000000000000140
1012. Al-Khatib SM, Stevenson WG, Ackerman MJ, et al. 2017 AHA/ACC/HRS Guideline for Management of Patients With Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol*. Oct 2 2018;72(14):e91-e220. doi:10.1016/j.jacc.2017.10.054

- 1113.** Reiffel JA, Camm AJ, Belardinelli L, et al. The HARMONY Trial: Combined Ranolazine and Dronedarone in the Management of Paroxysmal Atrial Fibrillation: Mechanistic and Therapeutic Synergism. *Circ Arrhythm Electrophysiol*. Oct 2015;8(5):1048-56. doi:10.1161/circep.115.002856
- 1214.** Anagnostopoulos C, Harbinson M, Kelion A, et al. Procedure guidelines for radionuclide myocardial perfusion imaging. *Heart*. 2004;90 Suppl 1(Suppl 1):i1-i10. doi:10.1136/heart.90.suppl_1.i1
- 1315.** Gräni C, Buechel RR, Kaufmann PA, Kwong RY. Multimodality Imaging in Individuals With Anomalous Coronary Arteries. *JACC Cardiovasc Imaging*. Apr 2017;10(4):471-481. doi:10.1016/j.jcmg.2017.02.004
- 1416.** Tang K, Wang L, Shi R, et al. The role of myocardial perfusion imaging in evaluating patients with myocardial bridging. *J Nucl Cardiol*. Feb 2011;18(1):117-22. doi:10.1007/s12350-010-9303-6
- 1517.** McCrindle BW, Rowley AH, Newburger JW, et al. Diagnosis, Treatment, and Long-Term Management of Kawasaki Disease: A Scientific Statement for Health Professionals From the American Heart Association. *Circulation*. Apr 25 2017;135(17):e927-e999. doi:10.1161/cir.0000000000000484
- 1618.** Lancellotti P, Nkomo VT, Badano LP, et al. Expert consensus for multi-modality imaging evaluation of cardiovascular complications of radiotherapy in adults: a report from the European Association of Cardiovascular Imaging and the American Society of Echocardiography. *Eur Heart J Cardiovasc Imaging*. Aug 2013;14(8):721-40. doi:10.1093/ehjci/jet123
- 1719.** Birnie DH, Nery PB, Ha AC, Beanlands RS. Cardiac Sarcoidosis. *J Am Coll Cardiol*. Jul 26 2016;68(4):411-21. doi:10.1016/j.jacc.2016.03.605
- 1820.** Blankstein R, Waller AH. Evaluation of Known or Suspected Cardiac Sarcoidosis. *Circ Cardiovasc Imaging*. Mar 2016;9(3):e000867. doi:10.1161/circimaging.113.000867
- 1921.** Vita T, Okada DR, Veillet-Chowdhury M, et al. Complementary Value of Cardiac Magnetic Resonance Imaging and Positron Emission Tomography/Computed Tomography in the Assessment of Cardiac Sarcoidosis. *Circ Cardiovasc Imaging*. Jan 2018;11(1):e007030. doi:10.1161/circimaging.117.007030
- 2022.** Skali H, Schulman AR, Dorbala S. 18F-FDG PET/CT for the assessment of myocardial sarcoidosis. *Curr Cardiol Rep*. Apr 2013;15(4):352.
- 23.** Habib G, Lancellotti P, Antunes MJ, et al. 2015 ESC Guidelines for the management of infective endocarditis: The Task Force for the Management of Infective Endocarditis of the European Society of Cardiology (ESC). Endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the European Association of Nuclear Medicine (EANM). *Eur Heart J*. Nov 21 2015;36(44):3075-3128. doi:10.1093/eurheartj/ehv319
- 2124.** Doherty JU, Kort S, Mehran R, Schoenhagen P, Soman P. ACC/AATS/AHA/ASE/ASNC/HRS/SCAI/SCCT/SCMR/STS 2017 Appropriate Use Criteria for Multimodality Imaging in Valvular Heart Disease: A Report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular

Magnetic Resonance, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. Sep 26 2017;70(13):1647-1672. doi:10.1016/j.jacc.2017.07.732

~~22. Bhawe NM, Nienaber CA, Clough RE, Eagle KA. Multimodality Imaging of Thoracic Aortic Diseases in Adults. *JACC Cardiovasc Imaging*. Jun 2018;11(6):902-919. doi:10.1016/j.jcmg.2018.03.009~~

2325. Isselbacher EM, Preventza O, Hamilton Black J, 3rd, et al. 2022 ACC/AHA Guideline for the Diagnosis and Management of Aortic Disease: A Report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines. *Circulation*. Dec 13 2022;146(24):e334-e482. doi:10.1161/cir.0000000000001106

26. Kristensen SD, Knuuti J, Saraste A, et al. 2014 ESC/ESA Guidelines on non-cardiac surgery: cardiovascular assessment and management: The Joint Task Force on non-cardiac surgery: cardiovascular assessment and management of the European Society of Cardiology (ESC) and the European Society of Anaesthesiology (ESA). *Eur Heart J*. Sep 14 2014;35(35):2383-431. doi:10.1093/eurheartj/ehu282

2427. Fleisher LA, Fleischmann KE, Auerbach AD, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *J Am Coll Cardiol*. Dec 09 2014;64(22):e77-137. doi:10.1016/j.jacc.2014.07.944

2528. Velasco A, Reyes E, Hage FG. Guidelines in review: Comparison of the 2014 ACC/AHA guidelines on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery and the 2014 ESC/ESA guidelines on noncardiac surgery: Cardiovascular assessment and management. *J Nucl Cardiol*. 02 2017;24(1):165-170. doi:10.1007/s12350-016-0643-8

2629. Lentine KL, Costa SP, Weir MR, et al. Cardiac disease evaluation and management among kidney and liver transplantation candidates: a scientific statement from the American Heart Association and the American College of Cardiology Foundation. *J Am Coll Cardiol*. Jul 31 2012;60(5):434-80. doi:10.1016/j.jacc.2012.05.008

2730. Mc Ardle BA, Dowsley TF, deKemp RA, Wells GA, Beanlands RS. Does rubidium-82 PET have superior accuracy to SPECT perfusion imaging for the diagnosis of obstructive coronary disease?: A systematic review and meta-analysis. *J Am Coll Cardiol*. Oct 30 2012;60(18):1828-37. doi:10.1016/j.jacc.2012.07.038

2831. Bateman TM, Dilsizian V, Beanlands RS, DePuey EG, Heller GV, Wolinsky DA. American Society of Nuclear Cardiology and Society of Nuclear Medicine and Molecular Imaging Joint Position Statement on the Clinical Indications for Myocardial Perfusion PET. *J Nucl Med*. Oct 2016;57(10):1654-1656. doi:10.2967/jnumed.116.180448

2932. Fazel R, Dilsizian V, Einstein AJ, Ficaro EP, Henzlova M, Shaw LJ. Strategies for defining an optimal risk-benefit ratio for stress myocardial perfusion SPECT. *J Nucl Cardiol*. May 2011;18(3):385-92. doi:10.1007/s12350-011-9353-4

~~30. Montalescot G, Sechtem U, Achenbach S, et al. 2013 ESC guidelines on the management of stable coronary artery disease: the Task Force on the management of stable coronary artery~~

~~disease of the European Society of Cardiology. *Eur Heart J*. Oct 2013;34(38):2949-2003.
doi:10.1093/eurheartj/ehz296~~

3433. Shen WK, Sheldon RS, Benditt DG, et al. 2017 ACC/AHA/HRS Guideline for the Evaluation and Management of Patients With Syncope: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol*. Aug 1 2017;70(5):620-663.
doi:10.1016/j.jacc.2017.03.002

3234. Mark DB, Hlatky MA, Harrell FE, Jr., Lee KL, Califf RM, Pryor DB. Exercise treadmill score for predicting prognosis in coronary artery disease. *Ann Intern Med*. Jun 1987;106(6):793-800.
doi:10.7326/0003-4819-106-6-793

3335. Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. Sep 10 2019;74(10):e177-e232. doi:10.1016/j.jacc.2019.03.010

3436. D'Agostino RB, Sr., Vasan RS, Pencina MJ, et al. General cardiovascular risk profile for use in primary care: the Framingham Heart Study. *Circulation*. Feb 12 2008;117(6):743-53.
doi:10.1161/circulationaha.107.699579

3537. Goff DC, Jr., Lloyd-Jones DM, Bennett G, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. Jul 1 2014;63(25 Pt B):2935-2959. doi:10.1016/j.jacc.2013.11.005

3638. McClelland RL, Jorgensen NW, Budoff M, et al. 10-Year Coronary Heart Disease Risk Prediction Using Coronary Artery Calcium and Traditional Risk Factors: Derivation in the MESA (Multi-Ethnic Study of Atherosclerosis) With Validation in the HNR (Heinz Nixdorf Recall) Study and the DHS (Dallas Heart Study). *J Am Coll Cardiol*. Oct 13 2015;66(15):1643-53.
doi:10.1016/j.jacc.2015.08.035

3739. Ridker PM, Buring JE, Rifai N, Cook NR. Development and validation of improved algorithms for the assessment of global cardiovascular risk in women: the Reynolds Risk Score. *Jama*. Feb 14 2007;297(6):611-9. doi:10.1001/jama.297.6.611

3840. Patel MR, Bailey SR, Bonow RO, et al.
ACCF/SCAI/AATS/AHA/ASE/ASNC/HFSA/HRS/SCCM/SCCT/SCMR/STS 2012 appropriate use criteria for diagnostic catheterization: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, Society for Cardiovascular Angiography and Interventions, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, Society of Thoracic Surgeons. *J Thorac Cardiovasc Surg*. Jul 2012;144(1):39-71.
doi:10.1016/j.jtcvs.2012.04.013

3941. Lotfi A, Davies JE, Fearon WF, Grines CL, Kern MJ, Klein LW. Focused update of expert consensus statement: Use of invasive assessments of coronary physiology and structure: A

position statement of the society of cardiac angiography and interventions. *Catheter Cardiovasc Interv.* Aug 1 2018;92(2):336-347. doi:10.1002/ccd.27672

ADDITIONAL RESOURCES

1. Hendel RC, Berman DS, Di Carli MF, et al. ACCF/ASNC/ACR/AHA/ASE/SCCT/SCMR/SNM 2009 Appropriate Use Criteria for Cardiac Radionuclide Imaging: A Report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, the American Society of Nuclear Cardiology, the American College of Radiology, the American Heart Association, the American Society of Echocardiography, the Society of Cardiovascular Computed Tomography, the Society for Cardiovascular Magnetic Resonance, and the Society of Nuclear Medicine. *J Am Coll Cardiol.* Jun 9 2009;53(23):2201-29. doi:10.1016/j.jacc.2009.02.013
2. Hirshfeld JW, Jr., Ferrari VA, Bengel FM, et al. 2018 ACC/HRS/NASCI/SCAI/SCCT Expert Consensus Document on Optimal Use of Ionizing Radiation in Cardiovascular Imaging—Best Practices for Safety and Effectiveness, Part 1: Radiation Physics and Radiation Biology: A Report of the American College of Cardiology Task Force on Expert Consensus Decision Pathways Developed in Collaboration With Mended Hearts. *Catheter Cardiovasc Interv.* Aug 1 2018;92(2):203-221. doi:10.1002/ccd.27660
3. Jaarsma C, Leiner T, Bekkers SC, et al. Diagnostic performance of noninvasive myocardial perfusion imaging using single-photon emission computed tomography, cardiac magnetic resonance, and positron emission tomography imaging for the detection of obstructive coronary artery disease: a meta-analysis. *J Am Coll Cardiol.* May 8 2012;59(19):1719-28. doi:10.1016/j.jacc.2011.12.040
4. Osborne MT, Hultén EA, Singh A, et al. Reduction in ^{18}F -fluorodeoxyglucose uptake on serial cardiac positron emission tomography is associated with improved left ventricular ejection fraction in patients with cardiac sarcoidosis. *J Nucl Cardiol.* Feb 2014;21(1):166-74. doi:10.1007/s12350-013-9828-6
5. Parker MW, Iskandar A, Limone B, et al. Diagnostic accuracy of cardiac positron emission tomography versus single-photon emission computed tomography for coronary artery disease: a bivariate meta-analysis. *Circ Cardiovasc Imaging.* Nov 2012;5(6):700-7. doi:10.1161/circimaging.112.978270
6. Patel MR, Bailey SR, Bonow RO, et al. ACCF/SCAI/AATS/AHA/ASE/ASNC/HFSA/HRS/SCCM/SCCT/SCMR/STS 2012 appropriate use criteria for diagnostic catheterization: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, Society for Cardiovascular Angiography and Interventions, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *J Am Coll Cardiol.* May 29 2012;59(22):1995-2027. doi:10.1016/j.jacc.2012.03.003

7. ~~Rahbar K, Seifarth H, Schäfers M, et al. Differentiation of malignant and benign cardiac tumors using 18F-FDG PET/CT. *J Nucl Med*. Jun 2012;53(6):856-63. doi:10.2967/jnumed.111.095364~~
8. ~~Salaun E, Habib G. Beyond Standard Echocardiography in Infective Endocarditis: Computed Tomography, 3-Dimensional Imaging, and Multi-Imaging. *Circ Cardiovasc Imaging*. Mar 2018;11(3):e007626. doi:10.1161/circimaging.118.007626~~
9. ~~Schindler TH, Schelbert HR, Quercioli A, Dilsizian V. Cardiac PET imaging for the detection and monitoring of coronary artery disease and microvascular health. *JACC Cardiovasc Imaging*. Jun 2010;3(6):623-40. doi:10.1016/j.jcmg.2010.04.007~~

~~**Disclaimer:** Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.~~

Disclaimer: National Imaging Associates, Inc. (NIA) authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Evolent Health LLC subsidiaries including, but not limited to, National Imaging Associates (“NIA”). The policies constitute only the reimbursement and coverage guidelines of NIA. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. NIA reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.