

# **AmeriHealth Caritas Louisiana**

*National Imaging Associates, Inc.*	
Clinical guidelines	Original Date: June 2007
FUNCTIONAL BRAIN MRI	
CPT Codes: 70554, 70555	Last Revised Date: May 20223
Guideline Number: NIA_CG_013	Implementation Date: January 20234

### **GENERAL INFORMATION**

- <u>It is an expectation that all patients receive care/services from a licensed clinician. All</u> <u>appropriate supporting documentation, including recent pertinent office visit notes, laboratory</u> <u>data, and results of any special testing must be provided. If applicable: All prior relevant imaging</u> <u>results and the reason that alternative imaging cannot be performed must be included in the</u> <u>documentation submitted.</u>
- Where a specific clinical indication is not directly addressed in this guideline, medical necessity determination will be made based on widely accepted standard of care criteria. These criteria are supported by evidence-based or peer-reviewed sources such as medical literature, societal guidelines and state/national recommendations.

## **INDICATIONS FOR FUNCTIONAL BRAIN MRIPre-operative/procedural Evaluation<sup>1</sup>**

In the following where fMRI may have a significant role in the mapping of a lesion in relation to eloquent cortex (i.e., <sup>1, 2</sup>

## Pre-operative/procedural Evaluation<sup>1</sup>

In the following where fMRI may have a significant role in the mapping of a lesion in relation to eloquent cortex (i.e., language, motor, sensory and visual centers) to determine the appropriateness of surgical intervention

- Focal brain lesion (i.e., tumor or vascular malformation) for presurgical planning<sup>2-53-6</sup>
- Pre-operative evaluation for epilepsy surgery<sup>6,77,8</sup>
- Brain tumor for radiation treatment planning<sup>8,99,10</sup>

#### Post-operative/procedural Evaluation

• Therapeutic follow-up. A documented medical reason must clearly explain the medical necessity for follow up (i.e., evaluation of post-treatment eloquent cortex).

\*National Imaging Associates, Inc. (NIA) is a subsidiary of Evolent Health LLC. © 2007-2024 National Imaging Associates, Inc., All Rights Reserved.

<sup>\*</sup> National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.

# BACKGROUND

Functional MRI (fMRI) of the brain is a non-invasive imaging technique, using radio waves and a strong magnetic field, to image the brain activity of a patient prior to undergoing brain surgery for tumors or epilepsy. It is based on the increase in blood flow to the local vasculature when parts of the brain are activated and helps to determine the location of vital areas of brain function. fMRI images capture blood oxygen levels in parts of the brain that are responsible for perception, cognition, and movement allowing neurosurgeons to operate with less possibility of harming areas that are critical to the patient's quality of life. fMRI is primarily used for presurgical planning, operative risk assessment and therapeutic follow-up.

# Task vs Resting-state fMRI

During resting-state fMRI (rs-fMRI), unlike task-based functional MRI, the individual is not required<sup>11-13</sup> to perform any specific task. This is beneficial for patients who have difficulty performing tasks, such as pediatric and certain neurologic or psychiatric patients. This technique has been well-utilized in research, and its clinical use is increasing considerably, especially in presurgical planning (e.g., mapping epileptic foci) and neuropsychiatric diseases. For the above indications, non-tasked based fMRI such as resting state fMRI can also be performed.

**fMRI** as an Alternative to the Invasive Wada test and Direct Electrical Stimulation – fMRI is considered an alternative to the Wada test and direct electrical stimulation as it is a non-invasive method for location of vital brain areas. The Wada test is used for the pre-operative evaluations of patients with brain tumors and seizures to determine which side of the brain is responsible for vital cognitive functions, e.g., speech and memory. It can assess the surgical risk of damaging the vital areas of the brain. The Wada test is invasive, involving an angiography procedure to guide a catheter to the internal carotid where a barbiturate is injected, putting one hemisphere of the brain to sleep. Direct electrical stimulation mapping is invasive requiring the placement of electrodes in the brain. The electrodes are used to stimulate multiple cortical sites in the planned area of resection to allow the surgeons to identify and mark which areas can be safely resected.<sup>40,1114,15</sup>

**fMRI and Brain Tumors** – fMRI may significantly affect therapeutic planning in patients who have potentially resectable brain tumors. Due to its non-invasiveness, its relatively high spatial resolution, and its pre-operative results, fMRI is used before surgery in the evaluation of patients with brain tumors. fMRI may have a significant role in mapping lesions that are located in close proximity to vital areas of brain function (language, sensory motor, and visual). It can determine the precise spatial relationship between the lesion and adjacent functionally essential parenchyma, allowing removal of as much pathological tissue as possible during

Page **2** of **9** Functional Brain MRI



resection of brain tumors without compromising essential brain functions. fMRI provides an alternative to other invasive tests, such as the Wada test and direct electrical stimulation.<sup>4216</sup>

**fMRI and Seizures** – Brain fMRI can influence the diagnostic and therapeutic decisions of the seizure team, thereby affecting the surgical approach and outcomes. Brain surgery is often the treatment for patients with refractory epilepsy, especially patients with a single seizure focus. fMRI can be used to image and localize abnormal brain function in patients with seizures. fMRI can help determine brain functions (language, sensory motor, and visual) of areas bordering the lesion, resulting in better outcomes with less neurologic deficit.<sup>78</sup>

fMRI is increasingly being used to evaluate candidates for surgical treatment of intractable epilepsy (Phase 1 evaluation) and can aid in surgical decision-making. It can 1) help to improve functional outcome by enabling surgery that spares functional cortex, 2) guide surgical intervention by revealing when reorganization of function has occurred, and 3) show when abnormal cortex is also functionally active, and hence that surgery may not be the best option<sup>17, 18</sup>.



# **REFERENCES**

1. American College of Radiology, American Society of Neuroradiology, Society for Pediatric Radiology. ACR–ASNR–SPR practice parameter for the performance of functional magnetic resonance imaging (fMRI) of the brain. American College of Radiology. Updated 2022. Accessed January 29, 2023. https://www.acr.org/-/media/ACR/Files/Practice-Parameters/fmr-brain.pdf

2. Kumar VA, Heiba IM, Prabhu SS, et al. The role of resting-state functional MRI for clinical preoperative language mapping. *Cancer Imaging*. Jul 11 2020;20(1):47. doi:10.1186/s40644-020-00327-w

3. Jiao Y, Lin F, Wu J, et al. Brain Arteriovenous Malformations Located in Language Area: Surgical Outcomes and Risk Factors for Postoperative Language Deficits. *World Neurosurg*. Sep 2017;105:478-491. doi:10.1016/j.wneu.2017.05.159

4. Silva MA, See AP, Essayed WI, Golby AJ, Tie Y. Challenges and techniques for presurgical brain mapping with functional MRI. *Neuroimage Clin*. 2018;17:794-803.

doi:10.1016/j.nicl.2017.12.008

5. Stancanello J, Cavedon C, Francescon P, et al. BOLD fMRI integration into radiosurgery treatment planning of cerebral vascular malformations. *Med Phys.* Apr 2007;34(4):1176-84. doi:10.1118/1.2710326

6. Vysotski S, Madura C, Swan B, et al. Preoperative FMRI Associated with Decreased Mortality and Morbidity in Brain Tumor Patients. *Interdiscip Neurosurg*. Sep 2018;13:40-45. doi:10.1016/j.inat.2018.02.001

7. Benjamin CFA, Dhingra I, Li AX, et al. Presurgical language fMRI: Technical practices in epilepsy surgical planning. *Hum Brain Mapp*. Oct 2018;39(10):4032-4042. doi:10.1002/hbm.24229

8. Janecek JK, Swanson SJ, Sabsevitz DS, et al. Language lateralization by fMRI and Wada testing in 229 patients with epilepsy: rates and predictors of discordance. *Epilepsia*. Feb 2013;54(2):314-22. doi:10.1111/epi.12068

9. Kovács A, Tóth L, Glavák C, et al. Integrating functional MRI information into conventional 3D radiotherapy planning of CNS tumors. Is it worth it? *J Neurooncol*. Dec 2011;105(3):629-37. doi:10.1007/s11060-011-0633-2

<u>10. Wang M, Ma H, Wang X, et al. Integration of BOLD-fMRI and DTI into radiation treatment</u> planning for high-grade gliomas located near the primary motor cortexes and corticospinal tracts. *Radiat Oncol*. Mar 8 2015;10:64. doi:10.1186/s13014-015-0364-1

11. Lv H, Wang Z, Tong E, et al. Resting-State Functional MRI: Everything That Nonexperts Have Always Wanted to Know. *AJNR Am J Neuroradiol*. Aug 2018;39(8):1390-1399. doi:10.3174/ajnr.A5527

<u>12. J. K, A. G, L. T, K. T, S. F, R. G. National medicare trends in the utilization of fMRI.</u> *Neuroscience Informatics*. March 2022 2022;2(1):100031.

13. Lee MH, Smyser CD, Shimony JS. Resting-state fMRI: a review of methods and clinical applications. *AJNR Am J Neuroradiol*. Oct 2013;34(10):1866-72. doi:10.3174/ajnr.A3263



14. Binder JR. Functional MRI is a valid noninvasive alternative to Wada testing. *Epilepsy Behav.* Feb 2011;20(2):214-22. doi:10.1016/j.yebeh.2010.08.004

15. Bizzi A, Blasi V, Falini A, et al. Presurgical functional MR imaging of language and motor functions: validation with intraoperative electrocortical mapping. *Radiology*. Aug 2008;248(2):579-89. doi:10.1148/radiol.2482071214

16. Petrella JR, Shah LM, Harris KM, et al. Preoperative functional MR imaging localization of language and motor areas: effect on therapeutic decision making in patients with potentially resectable brain tumors. *Radiology*. Sep 2006;240(3):793-802. doi:10.1148/radiol.2403051153

17. Liégeois F, Cross JH, Gadian DG, Connelly A. Role of fMRI in the decision-making process: epilepsy surgery for children. *J Magn Reson Imaging*. Jun 2006;23(6):933-40. doi:10.1002/jmri.20586

18. Vakharia VN, Duncan JS, Witt J-A, Elger CE, Staba R, Engel Jr J. Getting the best outcomes from epilepsy surgery. *Annals of Neurology*. 2018;83(4):676-690.

doi:https://doi.org/10.1002/ana.25205



# **POLICY HISTORY**

Date	Summary
<u>May 2023</u>	Updated background and references
	<ul> <li>Added - to determine the appropriateness of surgical</li> </ul>
	intervention
	<ul> <li>Background section regarding non-task-based fMRI</li> </ul>
	General Information moved to beginning of guideline with added
	statement on clinical indications not addressed in this guideline
May 2022	Updated Background and references
February 2021	Updated References
May 2020	Updated References
	Re-ordered indications
August 2019	<ul> <li>Modified pre-operative/procedural evaluation section to include</li> </ul>
	focal brain lesion for pre-surgical planning, brain tumor for
	radiation treatment planning AND epilepsy surgery pre-operative
	evaluation.

#### REFERENCES

1. American College of Radiology, American Society of Neuroradiology, Society for Pediatric Radiology. ACR-ASNR-SPR practice parameter for the performance of functional magnetic resonance imaging (fMRI) of the brain. American College of Radiology. Updated 2017. Accessed December 21, 2021. <u>https://www.acr.org/-/media/ACR/Files/Practice-Parameters/fmr-</u> <u>brain.pdf</u>

2. Jiao Y, Lin F, Wu J, et al. Brain Arteriovenous Malformations Located in Language Area: Surgical Outcomes and Risk Factors for Postoperative Language Deficits. *World Neurosurg*. Sep 2017;105:478-491. doi:10.1016/j.wneu.2017.05.159

3. Silva MA, See AP, Essayed WI, Golby AJ, Tie Y. Challenges and techniques for presurgical brain mapping with functional MRI. *Neuroimage Clin.* 2018;17:794-803.

doi:10.1016/j.nicl.2017.12.008

4. Stancanello J, Cavedon C, Francescon P, et al. BOLD fMRI integration into radiosurgery treatment planning of cerebral vascular malformations. *Med Phys.* Apr 2007;34(4):1176-84. doi:10.1118/1.2710326

5. Vysotski S, Madura C, Swan B, et al. Preoperative FMRI Associated with Decreased Mortality and Morbidity in Brain Tumor Patients. *Interdiscip Neurosurg*. Sep 2018;13:40-45. doi:10.1016/j.inat.2018.02.001

6. Benjamin CFA, Dhingra I, Li AX, et al. Presurgical language fMRI: Technical practices in epilepsy surgical planning. *Hum Brain Mapp*. Oct 2018;39(10):4032-4042. doi:10.1002/hbm.24229

7. Janecek JK, Swanson SJ, Sabsevitz DS, et al. Language lateralization by fMRI and Wada testing in 229 patients with epilepsy: rates and predictors of discordance. *Epilepsia*. Feb 2013;54(2):314-22. doi:10.1111/epi.12068

8. Kovács A, Tóth L, Glavák C, et al. Integrating functional MRI information into conventional 3D radiotherapy planning of CNS tumors. Is it worth it? *J Neurooncol*. Dec 2011;105(3):629-37. doi:10.1007/s11060-011-0633-2

9. Wang M, Ma H, Wang X, et al. Integration of BOLD-fMRI and DTI into radiation treatment planning for high-grade gliomas located near the primary motor cortexes and corticospinal tracts. *Radiat Oncol*. Mar 8 2015;10:64. doi:10.1186/s13014-015-0364-1

10. Binder JR. Functional MRI is a valid noninvasive alternative to Wada testing. *Epilepsy Behav*. Feb 2011;20(2):214-22. doi:10.1016/j.yebeh.2010.08.004

11. Bizzi A, Blasi V, Falini A, et al. Presurgical functional MR imaging of language and motor functions: validation with intraoperative electrocortical mapping. *Radiology*. Aug 2008:248(2):579-89. doi:10.1148/radiol.2482071214

12. Petrella JR, Shah LM, Harris KM, et al. Preoperative functional MR imaging localization of language and motor areas: effect on therapeutic decision making in patients with potentially resectable brain tumors. *Radiology*. Sep 2006;240(3):793-802. doi:10.1148/radiol.2403051153

#### **ADDITIONAL RESOURCES**

Page **7** of **9** Functional Brain MRI



1. American College of Radiology, American Society of Neuroradiology, Society for Pediatric Radiology. ACR-ASNR-SPR practice parameter for the performance of functional magnetic resonance imaging (fMRI) of the brain. American College of Radiology. Updated 2017. Accessed December 21, 2021. <u>https://www.acr.org/\_/media/ACR/Files/Practice\_Parameters/fmr-</u> <u>brain.pdf</u>

2. Jiao Y, Lin F, Wu J, et al. Brain Arteriovenous Malformations Located in Language Area: Surgical Outcomes and Risk Factors for Postoperative Language Deficits. *World Neurosurg*. Sep 2017;105:478-491. doi:10.1016/j.wneu.2017.05.159

3. Silva MA, See AP, Essayed WI, Golby AJ, Tie Y. Challenges and techniques for presurgical brain mapping with functional MRI. *Neuroimage Clin.* 2018;17:794-803. doi:10.1016/j.nicl.2017.12.008

4. Stancanello J, Cavedon C, Francescon P, et al. BOLD fMRI integration into radiosurgery treatment planning of cerebral vascular malformations. *Med Phys.* Apr 2007;34(4):1176-84. doi:10.1118/1.2710326

5. Vysotski S, Madura C, Swan B, et al. Preoperative FMRI Associated with Decreased Mortality and Morbidity in Brain Tumor Patients. *Interdiscip Neurosurg*. Sep 2018;13:40-45. doi:10.1016/j.inat.2018.02.001

6. Benjamin CFA, Dhingra I, Li AX, et al. Presurgical language fMRI: Technical practices in epilepsy surgical planning. *Hum Brain Mapp*. Oct 2018;39(10):4032–4042. doi:10.1002/hbm.24229

7. Janecek JK, Swanson SJ, Sabsevitz DS, et al. Language lateralization by fMRI and Wada testing in 229 patients with epilepsy: rates and predictors of discordance. *Epilepsia*. Feb 2013;54(2):314-22. doi:10.1111/epi.12068

8. Kovács A, Tóth L, Glavák C, et al. Integrating functional MRI information into conventional 3D radiotherapy planning of CNS tumors. Is it worth it? *J Neurooncol*. Dec 2011;105(3):629-37. doi:10.1007/s11060-011-0633-2

9. Wang M, Ma H, Wang X, et al. Integration of BOLD-fMRI and DTI into radiation treatment planning for high-grade gliomas located near the primary motor cortexes and corticospinal tracts. *Radiat Oncol*. Mar 8 2015;10:64. doi:10.1186/s13014-015-0364-1

10. Binder JR. Functional MRI is a valid noninvasive alternative to Wada testing. *Epilepsy Behav*. Feb 2011;20(2):214-22. doi:10.1016/j.yebeh.2010.08.004

11. Bizzi A, Blasi V, Falini A, et al. Presurgical functional MR imaging of language and motor functions: validation with intraoperative electrocortical mapping. *Radiology*. Aug 2008;248(2):579–89. doi:10.1148/radiol.2482071214

12. Petrella JR, Shah LM, Harris KM, et al. Preoperative functional MR imaging localization of language and motor areas: effect on therapeutic decision making in patients with potentially resectable brain tumors. *Radiology*. Sep 2006;240(3):793-802. doi:10.1148/radiol.2403051153





## **Reviewed / Approved by NIA Clinical Guideline Committee**

#### GENERAL INFORMATION

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

**Disclaimer:** Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates ("Magellan"). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.

Disclaimer: National Imaging Associates, Inc. (NIA) authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Evolent Health LLC subsidiaries including, but not limited to, National Imaging Associates ("NIA"). The policies constitute only the reimbursement and coverage guidelines of NIA. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. NIA reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.

Page **9** of **9** Functional Brain MRI

