

## Clinical Policy: Ferric Gluconate (Ferrlecit)

Reference Number: LA.PHAR.166

Effective Date:

Last Review Date: 01.21

Line of Business: Medicaid

Coding Implications

Revision Log

See Important Reminder at the end of this policy for important regulatory and legal information.

### Description

Sodium ferric gluconate complex in sucrose (Ferrlecit®) injection is an iron replacement product.

### FDA Approved Indication(s)

Ferrlecit is indicated for the treatment of iron deficiency anemia (IDA) in adult patients and in pediatric patients age 6 years and older with chronic kidney disease (CKD) receiving hemodialysis who are receiving supplemental epoetin therapy.

### Policy/Criteria

Prior authorization is required. Provider must submit documentation (including such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that Ferrlecit is medically necessary when the following criteria are met:

#### **I. Initial Approval Criteria**

- A. Iron Deficiency Anemia associated with Chronic Kidney Disease (must meet all):**
  - 1. Diagnosis of IDA and CKD;**
  - 2. IDA is confirmed by either of the following:**
    - a. Transferrin saturation (TSAT) ≤ 30%;**
    - b. Serum ferritin ≤ 500 ng/mL;**
  - 3. If CKD does not require hemodialysis or peritoneal dialysis, oral iron therapy is not optimal due to any of the following:**
    - a. TSAT < 12%;**
    - b. Hgb < 7 g/dL;**
    - c. Symptomatic anemia;**
    - d. Severe or ongoing blood loss;**
    - e. Oral iron intolerance;**
    - f. Unable to achieve therapeutic targets with oral iron;**
    - g. Co-existing condition that may be refractory to oral iron therapy;**
  - 4. Dose does not exceed 125 mg of elemental iron per infusion/injection.**

Approval duration: 3 months

- B. Iron Deficiency Anemia without Chronic Kidney Disease (off-label) (must meet all):**
  - 1. Diagnosis of IDA confirmed by any of the following:**

- a. Serum ferritin < 15 ng/mL or < 30 ng/mL if pregnant;
- b. Serum ferritin ≤ 41 ng/mL and Hgb < 12 g/dL (women)/< 13 g/dL (men);
- c. TSAT < 20%;
- d. Absence of stainable iron in bone marrow;
- e. Increased soluble transferring receptor (sTfR) or sTfR-ferritin index;
- f. Increased erythrocyte protoporphyrin level;
2. Oral iron therapy is not optimal due to any of the following:
  - a. TSAT < 12%;
  - b. Hgb < 7 g/dL;
  - c. Symptomatic anemia;
  - d. Severe or ongoing blood loss;
  - e. Oral iron intolerance;
  - f. Unable to achieve therapeutic targets with oral iron;
  - g. Co-existing condition that may be refractory to oral iron therapy;
3. At the time of the request, member does not have CKD;
4. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

Approval duration 3 months

**C. Other diagnoses/indications:**

Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.

**II. Continued Approval Criteria**

- A. Iron Deficiency Anemia with Chronic Kidney Disease (must meet all):
  1. Currently receiving the medication via Louisiana Healthcare Connections benefit or member has previously met all initial approval criteria;
  2. Documentation of one of the following laboratory results measured since the last IV iron administration:
    - a. TSAT < 30%;
    - b. Serum ferritin ≤ 500 ng/mL;
  3. If request is for a dose increase, new dose does not exceed 125 mg elemental iron per infusion/injection.

Approval duration 3 months

**B. Iron Deficiency Anemia without Chronic Kidney Disease (off-label) (must meet all):**

1. Currently receiving the medication via Louisiana Healthcare Connections benefit or member has previously met all initial approval criteria;
2. Documentation of one of the following laboratory results measured since the last IV iron administration:
  - a. Serum ferritin < 15 ng/mL or < 30 ng/mL if pregnant;
  - b. Serum ferritin ≤ 41 ng/mL and Hb < 12 g/dL (women)/< 13 g/dL (men);
  - c. TSAT < 20%;
  - d. Absence of stainable iron in bone marrow;

# CLINICAL POLICY

## Ferric Gluconate



- e. Increased sTfR or sTfR-ferritin index;
- f. Increased erythrocyte protoporphyrin level;
3. At the time of the request, member does not have CKD;
4. If request is for a dose increase, new dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

Approval duration 3 months

**C. Other diagnoses/indications (must meet 1 or 2):**

1. Currently receiving medication via Louisiana Healthcare Connections benefit and documentation supports positive response to therapy.  
Approval duration: Duration of request or 6 months (whichever is less); or
2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – LA.PMN.53 for Medicaid, or evidence of coverage documents.

**IV. Appendices/General Information**

Appendix A: Abbreviation/Acronym Key

CKD: chronic kidney disease

IDA: iron deficiency anemia

ESA: erythropoiesis stimulating agent

TSAT: transferrin saturation

Hb: hemoglobin

sTfR: soluble transferring receptor

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may require prior authorization.

<u>Drug Name</u>	<u>Dosing Regimen</u>	<u>Dose Limit/Maximum Dose</u>
<b>Examples of OTC Oral Iron Formulations*</b>		
<u>Ferrous fumarate (Ferretts, Ferrimin 150, Hemocyte)</u>	<u>Varies</u>	
<u>Ferrous gluconate (Ferate)</u>		
<u>Ferrous sulfate (BProtected Pedia Iron, Fer-In-Sol, FeroSul, FerrouSul, Iron Supplement, Iron Supplement Childrens, Slow Fe, Slow Iron)</u>		
<u>Polysaccharide-iron complex (EZFE 200, Ferrex 150, Ferrix x-150, Myferon 150, NovaFerrum 125, NovaFerrum 50, NovaFerrum Pediatric Drops, Nu-Iron, Poly-Iron 150)</u>		

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

## CLINICAL POLICY

### Ferric Gluconate

\*Oral formulations include elixirs, liquids, solutions, syrups, capsules, and tablets - including delayed/extended-release tablets.

#### Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): Known hypersensitivity to sodium ferric gluconate or any of its inactive components.
- Boxed warning(s): None reported.

#### V. Dosage and Administration

<u>Indication</u>	<u>Dosing Regimen</u>	<u>Maximum Dose</u>
<u>IDA with CKD with Hemodialysis: Iron Repletion</u>		
<u>Adults</u>	<u>125 mg by IV infusion or injection per dialysis session. - May require a cumulative dose of 1000 mg over 8 dialysis sessions.</u>	<u>125 mg of elemental iron per dose</u>
<u>Children age ≥ 6 years</u>	<u>1.5 mg/kg administered by IV infusion per dialysis session.</u>	<u>125 mg of elemental iron per dose</u>

#### VI. Product Availability

Intravenous solution single-dose vial: 12.5 mg/mL (5 mL)

#### VII. References

1. Ferrlecit prescribing information. Bridgewater, NJ: Sanofi-Aventis, Inc.; March 2020. Available from <http://products.sanofi.us/ferrlecit/ferrlecit.html>. Accessed November 18, 2020.
2. KDIGO 2012 clinical practice guideline for evaluation and management of chronic kidney disease. Kidney International Supplements. January 2013; 3(1): 1-136.
3. KDIGO 2012 clinical practice guideline for anemia in chronic kidney disease. Kidney International Supplements. August 2012; 2(4): 279-331.
4. Camaschella C. Iron-Deficiency Anemia. N Engl J Med. 2015; 372: 1832-43. DOI: 10.1056/NEJMra1401038.
5. Short MW, Domagalski JE. Iron Deficiency Anemia: Evaluation and Management. Am Fam Physician. 2013; 87(2): 98-104. <http://www.aafp.org/afp/2013/0115/p98.pdf>
6. Oral iron monographs. In: UpToDate (Lexicomp), Waltham, MA: Walters Kluwer Health; 2018. Available at [www.uptodate.com](http://www.uptodate.com). Accessed November 18, 2020.

#### Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

<u>HCPCS Codes</u>	<u>Description</u>
<u>J2916</u>	<u>Injection, sodium ferric gluconate complex in sucrose injection, 12.5 mg</u>

## CLINICAL POLICY

### Ferric Gluconate

<u>Reviews, Revisions, and Approvals</u>	<u>Date</u>
Converted corporate to local policy	01.21

#### Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information.  
LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results.  
Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

## CLINICAL POLICY

### Ferric Gluconate

**Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.**

**This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.**

**©2020 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.**