

AmeriHealth Caritas Louisiana

National Imaging Associates, Inc.*	
Clinical guidelines THORACIC SPINE CT	Original Date: September 1997
CPT Codes: 72128, 72129, 72130	Last Revised Date: May 2020
Guideline Number: NIA(CG)_043	Implementation Date: <u>January 2021 TBD</u>

GENERAL INFORMATION:

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. All prior relevant imaging results, and the reason that alternative imaging (gold standard, protocol, contrast, etc.) cannot be performed must be included in the documentation submitted.

INDICATIONS FOR THORACIC SPINE CT:

For evaluation of back pain with any of the following when Thoracic Spine MRI is contraindicated

(AANSCNS, 2014; Allegri, 2016; Jarvik, 2015; Last, 2009)

- With new or worsening objective neurologic deficits on exam**
- Failure of conservative treatment* for at least six (6) weeks within the last six (6) months (ACR, 2013; Eubanks, 2010).**
- With progression or worsening of symptoms during the course of conservative treatment*.**
- With an abnormal electromyography (EMG) or nerve conduction study (if performed) indicating a thoracic radiculopathy. (EMG is not recommended to determine the cause of axial lumbar, thoracic, or cervical spine pain (NASS, 2013)).**

For evaluation of chronic back pain with any of the following when Thoracic MRI is contraindicated

(Jarvik, 2015; Last, 2009; Allegri, 2016)

- With new or worsening objective neurologic deficits on exam**
- Failure of conservative treatment* for at least six (6) weeks (ACR, 2013; Eubanks, 2010) within the last six (6) months.**
- With progression or worsening of symptoms during the course of conservative treatment*.**

*** National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.**

With an abnormal electromyography (EMG) or nerve conduction study (if performed) indicating a thoracic radiculopathy. (EMG is not recommended to determine the cause of axial lumbar, thoracic, or cervical spine pain (NASS, 2013)).

For evaluation of new onset of back pain when Thoracic Spine MRI is contraindicated (AANSCNS, 2014)

With new or worsening objective neurologic deficits on exam
Failure of conservative treatment* for at least six (6) weeks within the last six (6) months.
With progression or worsening of symptoms during the course of conservative treatment*.
With an abnormal electromyography (EMG) or nerve conduction study (if performed) indicating a thoracic radiculopathy. (EMG is not recommended to determine the cause of axial lumbar, thoracic, or cervical spine pain (NASS, 2013)).

For evaluation of trauma or acute injury (ACR, 2012)8)

- Presents with any of the following neurological deficits: muscle weakness, abnormal reflexes, and/or sensory changes along a particular dermatome (nerve distribution).
- With progression or worsening of symptoms during the course of conservative treatment*.
- **History of underlying spinal abnormalities (i.e., ankylosing spondylitis or Diffuse idiopathic skeletal hyperostosis (DISH) (ACR, 2018; Koivikko, 2008; ACR 2018)**
- If the patient has had a recent CT scan of the chest, abdomen or pelvis, at the time of injury, those images should be reviewed prior to obtaining repeat imaging of the thoracic spine (ACR 2018)

For evaluation of known fracture

- To assess union of a fracture when physical examination, plain radiographs, or prior imaging suggest delayed or non-healing
- To determine the position of fracture fragments.

For evaluation of neurologic deficits when Thoracic Spine MRI is contraindicated or inappropriate

- With any of the following new neurological deficits: extremity muscular weakness; pathologic (e.g. Babinski, Hoffman's) or abnormal reflexes; or abnormal sensory changes along a particular dermatome (nerve distribution) as documented on physical exam; bowel or bladder dysfunction; spasticity, sensory, or motor level.
- With any of the following new neurological deficits: lower extremity weakness; abnormal reflexes; or abnormal sensory changes along a particular dermatome (nerve distribution) as documented on physical exam; evidence of Cauda Equina Syndrome; bowel or bladder dysfunction; spasticity, sensory or motor level (Stolper, 2017).

CT myelogram is indicated when signs and symptoms are incongruent with MRI findings or MRI cannot be performed/contraindicated (Grams, 2010; Morita, 2011; Naganawa, 2011; NASS, 2012; Ozdoba; 2011) (NASS, 2010)

For evaluation of suspected myelopathy when Thoracic Spine MRI is contraindicated (ACR, 2015; Ando, 2019; Behrbalk, 2013; ACR, 2015; Hou, 2016)

- Does not require conservative care
- Concurrent cervical/thoracic imaging not recommended
- Progressive symptoms including hand clumsiness, worsening handwriting, difficulty with
- With any of the following new neurological deficits: extremity muscular weakness; pathologic (e.g. Babinski, Hoffman's) or abnormal reflexes; or abnormal sensory changes along a particular dermatome (nerve distribution) as documented on physical exam; bowel or bladder dysfunction; spasticity, sensory, or motor level.
- Does not require conservative care
- Progressive symptoms including unsteadiness, broad-based gait, increased muscle tone, weakness and wasting of the upper and lower limbs; diminished sensation to light touch, temperature, proprioception, vibration; limb hyperreflexia and pathologic reflexes (upper or lower extremity); bowel and bladder dysfunction in more severe cases).
- Progressive symptoms including lower extremity weakness, numbness in the legs, increasing difficulty with balance and ambulation (Signs: unsteadiness, broad-based gait, increased muscle tone, pins and needles sensation, weakness and wasting of the lower limbs, and diminished sensation to light touch, temperature, proprioception, and vibration; limb hyperreflexia and pathologic reflexes; bowel and bladder dysfunction in more severe cases).

Ossification Posterior Longitudinal Ligament (OPPL) (Choi, 2011)

- Most common in cervical spine (rare but more severe in thoracic spine)
- CT to evaluate the calcification and MR for evaluation of cord.
 - Both CT and MRI would be approvable if surgery is planned as signal changes in the cord would suggest a poorer prognosis after surgery.

For evaluation of chronic back pain with any of the following when Thoracic MRI is contraindicated

(Harvik, 2015; Last, 2009; Allegri, 2016)

- With new or worsening objective neurologic deficits on exam
- Failure of conservative treatment* for at least six (6) weeks (ACR, 2013; Eubanks, 2010) within the last six (6) months.
- With progression or worsening of symptoms during the course of conservative treatment*.

- With an abnormal electromyography (EMG) or nerve conduction study (if performed) indicating a thoracic radiculopathy. (EMG is not recommended to determine the cause of axial lumbar, thoracic, or cervical spine pain (NASS, 2013)).

For evaluation of new onset of back pain when Thoracic Spine MRI is contraindicated (AANSCNS, 2014)

- With new or worsening objective neurologic deficits on exam
- Failure of conservative treatment* for at least six (6) weeks within the last six (6) months.
- With progression or worsening of symptoms during the course of conservative treatment*.
- With an abnormal electromyography (EMG) or nerve conduction study (if performed) indicating a thoracic radiculopathy. (EMG is not recommended to determine the cause of axial lumbar, thoracic, or cervical spine pain (NASS, 2013)).

~~Pars defect (spondylolysis) or spondylolisthesis~~

- ~~Pars defect (spondylolysis) or spondylolisthesis in adults when Flexion/Extension x rays show instability.~~
- ~~Pars defect (spondylolysis with spondylolisthesis) on plain films in pediatric population (<18 yr) (flexion extension instability not required) when MRI is contraindicated.~~

~~NOTE: Initial imaging (x ray, or planar bone scan without SPECT; Bone scan with SPECT is superior to MRI and CT in the detection of pars intrarticularis pathology including spondylolysis) (Matesan, 2016).~~

For evaluation of known or new compression fractures with worsening back pain (ACR, 2018)

- With history of malignancy (if MRI is contraindicated)
- With an associated new focal neurologic deficit
- Prior to a planned surgery/intervention or if the results of the ~~MRI-CT~~ will change management.

For evaluation of known tumor, cancer, or evidence of metastasis with any of the following
(MRI is usually the preferred study, but CT may help characterize solitary indeterminate bone lesions).

(Kim, 2012)

- For staging of known tumor.
- For follow-up evaluation of patient undergoing active cancer treatment.
- Presents with new signs or symptoms (e.g. physical, laboratory, and/or imaging findings) of new tumor or change in tumor.
- With evidence of metastasis on bone scan or previous imaging study.
-
- New or increasing non-traumatic thoracic back pain or radiculopathy or back that pain occurs at night, and wakes the patient from sleep** with known active cancer and a tumor that tends to metastasize to the spine and MRI is contraindicated (ACR, 2018; Ziu, 2019).

For evaluation of suspected tumor when Thoracic Spine MRI is contraindicated or inappropriate

(ACR, 2018)

- Prior abnormal or indeterminate imaging that requires further clarification.

Indication for combination studies for the initial pre-therapy staging of cancer, OR active monitoring for recurrence as clinically indicated, OR evaluation of suspected metastases:

- ≤ 5 concurrent studies to include CT or MRI of any of the following areas as appropriate depending on the cancer: Neck, Abdomen, Pelvis, Chest, Brain, Cervical Spine, Thoracic Spine, or Lumbar Spine.

For evaluation of known or suspected infection, abscess, or inflammatory disease when Thoracic MRI is contraindicated

(ACR, 2018; Lerner, 2018)

- Infection:
 - Infection:
 - Most common site is the lumbar spine (58%), followed by the thoracic spine (30%) and the cervical spine (11%)
 - As evidenced by signs and/or symptoms, laboratory (i.e., abnormal white blood cell count, ESR and/or CRP) or prior imaging findings (Bond, 2016).
 - High risk populations (indwelling hardware, history of endocarditis, IVDA, recent procedures) with appropriate signs/symptoms.
 - Follow up imaging of infection
 - With worsening symptoms/laboratory values (i.e., white blood cell count, ESR/CRP) or radiographic findings (Berbari, 2015)
- As evidenced by signs/symptoms, laboratory or prior imaging findings.

For evaluation of spine abnormalities related to immune system suppression, e.g., HIV, chemotherapy, leukemia, or lymphoma when Thoracic MRI is contraindicated

(ACR, 2018)

- As evidenced by signs/symptoms, laboratory, or prior imaging findings.

As part of initial post-operative / procedural evaluation (“CT best examination to assess for hardware complication, extent of fusion” (ACR, 2015; Rao, 2018) and MRI for cord, nerve root compression, disc pathology, or post-op infection):

- A follow-up study may be needed to help evaluate a patient’s progress after treatment, procedure, intervention, or surgery in the last 6 months. Documentation requires a medical reason that clearly indicates why additional imaging is needed for the type and area(s) requested.
- Changing neurologic status post-operatively.
- Surgical infection as evidenced by signs/symptoms, laboratory, or prior imaging findings.

- Residual or recurrent symptoms with any of the following neurological deficits: Lower extremity weakness, objective sensory loss, or abnormal reflexes (Rao, 2018).

Other Indications for a Thoracic Spine CT:

- For preoperative evaluation and Thoracic MRI is contraindicated
- Prior to spinal cord stimulator to exclude canal stenosis if no prior imaging of the thoracic spine has been done recently and MRI is contraindicated.
- CT discogram.
- Suspected cord compression with any of the following neurologic deficits, e.g., extremity weakness, sensory deficits, abnormal gait; abnormal reflexes; spinal level; bowel or bladder incontinence and Thoracic Spine MRI is contraindicated.
- Tethered cord, or spinal dysraphism (known or suspected) based on preliminary imaging, neurological exam, and/or high riskhigh-risk cutaneous stigmata (AANS; Duz, 2008; Milhorat, 2009; NIH) when Thoracic Spine MRI is contraindicated
- Ankylosing Spondylitis/Spondyloarthropathies with non-diagnostic or indeterminate x-ray and rheumatology workup (Akgul, 2011; Bennett, 2010; Ostergaard, 2012; Seiper, 2009)
- Known Arnold-Chiari syndrome and Thoracic MRI is contraindicated (Milhorat, 2009; Strahle, 2015).
- Congenital abnormalities when Thoracic Spine MRI is contraindicated or for characterization of bony detail (Trenga, 2016):
 - In the presence of neurologic deficit, progressive spinal deformity, or for preoperative planning (Trenga, 2016)
 - Back pain and vertebral anomalies (hemivertebrae, hypoplasia, agenesis, butterfly, segmentation defect, bars, or congenital wedging) in a child on preliminary imaging.
 - Scoliosis with any of the following:
 - Progressive spinal deformity;
 - Neurologic deficit;
 - Early onset;
 - Atypical curve (e.g., short segment, >30'-30 degee kyphosis, left thoracic curve, associated organ anomalies);
 - Pre-operative planning; OR
 - When office notes clearly document how imaging will change management.
- Syrinx or syringomyelia (known or suspected): and Thoracic Spine MRI is contraindicated:
 - With neurologic findings and/or predisposing conditions (e.g., Chiari malformation, prior trauma, neoplasm, arachnoiditis, severe spondylosis (Timpone, 2015)),
 - To further characterize a suspicious abnormality seen on prior imaging.
 - Known syrinx with worsening symptoms.
- CSF leak highly suspected and supported by patient history and/or physical exam findings (CT myelogram).
- For pediatric population (ACR, 2016)
 - Red flags that prompt imaging should include the presence of constant pain, night pain, and radicular pain lasting for 4 weeks or more and initial radiographs preformed (ACR, 2016)..

- Back pain associated with suspected inflammation, infection, or malignancy

COMBINATION STUDIES WITH THORACIC SPINE CT:

Cervical/Thoracic/Lumbar CTs:

- CT myelogram or discogram.
- Any combination of these for scoliosis survey in infant/child with congenital scoliosis or under the age of 10 ([ACR, 2018](#); Strahle, 2015; [ACR, 2018](#)).
- Any combination of these for spinal survey in patient with metastases.
- For evaluation of spinal abnormalities associated with Arnold-Chiari Malformation and Spine MRI is contraindicated. (C/T/L spine due to association with tethered cord and syringomyelia) (Milhorat, 2009; Strahle, 2015).
- Tethered cord, or spinal dysraphism (known or suspected) based on preliminary imaging, neurological exam, and/or cutaneous stigmata (AANS; Duz, 2008; Milhorat, 2009; [NIH](#)), when anesthesia required for imaging and **MRI is contraindicated**.
- Drop metastasis from brain or spine **when MRI contraindicated** (imaging also includes brain; CT spine imaging in this scenario is usually CT myelogram).
- **CSF leak highly suspected and supported by patient history and/or physical exam findings**

BACKGROUND:

Computed tomography is used for the evaluation, assessment of severity, and follow-up of diseases of the spine. Its use in the thoracic spine is limited, however, due to the lack of epidural fat in this part of the body. CT myelography improves the contrast severity of CT, but it is also invasive. CT may be used for conditions, e.g., degenerative changes, infection, and immune suppression, when magnetic resonance imaging (MRI) is contraindicated. It may also be used in the evaluation of tumors, cancer, or metastasis in the thoracic spine and it may be used for preoperative and post-surgical evaluations. CT obtains images from different angles and uses computer processing to show a cross-section of body tissues and organs. CT is fast and is often performed in acute settings. It provides good visualization of cortical bone.

OVERVIEW:

Ankylosing Spondylitis/Spondyloarthropathies is a cause of back or sacroiliac pain of insidious onset (usually > 3 month), associated with morning stiffness not relieved with rest (usually age at onset <40). It is associated with any of the following (Akgul, 2011; Bennett, 2010; Ostergaard, 2012; Seiper, 2014):

- Sedimentation rate and/or C-reactive protein (not an essential criteria).
- HLA B27 (not an essential criteria).
- Non-diagnostic or indeterminate x-ray
- Personal or family history of sacroilitis, peripheral inflammatory arthritis, and/or inflammatory bowel disease.

***Conservative Therapy:** (spine) should include a multimodality approach consisting of a **combination of active and inactive components**. Inactive components such as rest, ice, heat, modified activities, medical devices, acupuncture and/or stimulators, medications, injections (epidural, facet, bursal, and/or joint, not including trigger point), and diathermy can be utilized. Active modalities may consist of physical therapy, a physician supervised home exercise program**, regular Osteopathic Manipulative medicine treatments, and/or chiropractic care.

****Home Exercise Program - (HEP)/Therapy** – the following elements are required to meet guidelines for completion of conservative therapy (ACR, 2015; Last, 2009):

- Information provided on exercise prescription/plan AND
- Follow up with member with documentation provided regarding lack of improvement (failed) after completion of HEP (after suitable 6 week period), or inability to complete HEP due to physical reason- i.e. increased pain, inability to physically perform exercises. (Patient inconvenience or noncompliance without explanation does not constitute “inability to complete” HEP).
- Dates and duration of failed PT, physician supervised HEP, or chiropractic treatment should be documented in the original office notes or an addendum to the notes.

Myelopathy: Symptom severity varies and a high index of suspicion is essential for making the proper diagnosis in early cases. Symptoms of pain and radiculopathy may not be present. The natural history of myelopathy is characterized by neurological deterioration. The most frequently encountered symptom is gait abnormality (86%) followed by increased muscular reflexes (79.1%), pathological reflexes (65.1%), paresthesia of upper limb (69.8%), and pain (67.4%) (Vitzthum, 2007).

CT and Infection of the spine - Infection of the spine is not easy to differentiate from other spinal disorders, e.g., degenerative disease, spinal neoplasms, and non-infective inflammatory lesions. Infections may affect different parts of the spine, e.g., vertebrae, intervertebral discs, and paraspinal tissues. Imaging is important to obtain early diagnosis and treatment to avoid permanent neurology deficits. When MRI is contraindicated, CT may be used to evaluate infections of the spine.

CT and Degenerative Disc Disease – Degenerative disc disease is very common and CT may be indicated when MRI is contraindicated, when chronic degenerative changes are accompanied by conditions, e.g., new neurological deficits; onset of joint tenderness of a localized area of the spine; new abnormal nerve conduction studies; exacerbation of chronic back pain unresponsive to conservative treatment; and unsuccessful physical therapy/home exercise program.

Infection, Abscess, or Inflammatory disease

- Most common site is the lumbar spine (58%), followed by the thoracic spine (30%) and the cervical spine (11%) (Graeber, 2019)

- High risk populations (indwelling hardware, history of endocarditis, IVDA, recent procedures) with appropriate signs/symptoms

CT Myelogram

Myelography is the instillation of intrathecal contrast media under fluoroscopy. Patients are then imaged with CT to evaluate for spinal canal pathology. Although this technique has diminished greatly due to the advent of MRI due to its non-invasiveness and superior soft-tissue contrast, myelography is still a useful technique for conventional indications, such as spinal stenosis, when MRI is contraindicated or nondiagnostic, brachial plexus injury in neonates, radiation therapy treatment planning, and cerebrospinal fluid (CSF) leak (Pomerantz, 2016).

Cauda Equina Syndrome:

- Symptoms include severe back pain or sciatica along with one or more of the following:
 - Saddle anesthesia - loss of sensation restricted to the area of the buttocks, perineum, and inner surfaces of the thighs (areas that would sit on a saddle).
 - Recent bladder/bowel dysfunction (as listed above)
 - Achilles reflex absent on both sides
 - Sexual dysfunction that can come on suddenly
 - Absent anal reflex and bulbocavernosus reflex

MRI and Cutaneous Stigmata (Dias, 2015)

TABLE 1 Risk Stratification for Various Cutaneous Markers

High Risk	Intermediate Risk	Low Risk
Hypertrichosis	Capillary malformations (also referred to as NFS or salmon patch when pink and poorly defined, or PWS when darker red and well defined)	Coccygeal dimple
Infantile hemangioma		Light hair
Atretic meningocele		Isolated café au lait spots
DST		Mongolian spots
Subcutaneous lipoma		Hypo- and hypermelanotic macules or papules
Caudal appendage		Deviated or forked gluteal cleft
Segmental hemangiomas in association with LUMBAR syndrome		Nonmidline lesions

LUMBAR, lower body hemangioma and other cutaneous defects, urogenital abnormalities, ulcerations, myelopathy, bony defects, anorectal malformations, arterial anomalies, and renal anomalies.

Back Pain with Cancer History - Radiographic (x-ray) examination should be performed in cases of back pain when a patient has a cancer history, but without known active cancer or a tumor that tends to metastasize to the spine. This can make a diagnosis in many cases. This may occasionally allow for selection of bone scan in lieu of MRI in some cases. When radiographs do

not answer the clinical question, then MRI may be appropriate after a consideration of conservative care.

Neoplasms causing VCF (vertebral compression fractures) include: primary bone neoplasms, such as hemangioma or giant cell tumors, and tumor-like conditions causing bony and cellular remodeling, such as aneurysmal bone cysts, or Paget's disease (osteitis deformans); infiltrative neoplasms including and not limited to multiple myeloma and lymphoma, and metastatic neoplasms (ACR, 2018).

Most common spine metastasis involving primary metastasis originate from the following tumors in descending order: breast (21%), lung (19%), prostate (7.5%), renal (5%), gastrointestinal (4.5%), and thyroid (2.5%). While all tumor can seed to the spine, the cancers mentioned above metastasize to the spinal column early in the disease process (Ziu, 2019).

POLICY HISTORY:

Review Date: June 2019

Review Summary:

- **Added:**
 - new or worsening objective neuro deficits for chronic and acute back pain; CSF leak
 - last 6 months for allowable post op f/u period and removed EMG comment
 - red flags specifically for peds back pain and pain related to malignancy, infection, inflammation
 - new sections: pars defect; compression fractures; congenital abnormalities including section on scoliosis and vertebral anomalies in children w/back pain;
 - For combination studies cervical/thoracic/lumbar added drop metastasis, tumor evaluation for neurocutaneous syndromes, and abnormalities associated w/Arnold Chiari, as well as separate indication for tethered cord or spinal dysraphism
 - Spinal cord stimulator
 - New or increasing back pain in cancer patients with high suspicion of mets

Review Date: May 2020

Review Summary:

- For evaluation of neurologic deficits when new deficits are present
- Removed pars defect section
- Added ankylosing spondylitis for evaluation of trauma/acute injury
- Modified Initial imaging of new or increasing non-traumatic back pain or radiculopathy or back pain that occurs at night and wakes the patient from sleep with known active cancer and a tumor that tends to metastasize to the spine
- Added Imaging of Ossification of the Posterior Longitudinal Ligament (OPPL)
- Added Osteopathic Manipulative medicine to conservative care therapy

REFERENCES:

Akgul O, Ozgocmen S. Classification criteria for spondyloarthropathies. *World J Orthop.* Dec 18, 2011; 2(12):107-115. doi: 10.5312/wjo.v2.i12.07.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3302034/>. Retrieved March 29, 2018.

Allegri M, Montella S, Salici F, et al. Mechanisms of low back pain: A guide for diagnosis and therapy. *F1000Res.* 2016 Oct; [22\(F1000Res\)](#).

American Association of Neurological Surgeons and Congress of Neurological Surgeons (AANSCNS). Choosing Wisely®. <http://www.choosingwisely.org/clinician-lists/american-association-neurological-surgeons-imaging-for-nonspecific-acute-low-back-pain/>. Released June 24, 2014.

American College of Radiology (ACR). ACR Appropriateness Criteria®. Published 2015.

[Ando K, Imagama S, Kobayashi K, et al. Clinical features of thoracic myelopathy: A single-center study. *J Am Acad Orthop Surg Glob Res Rev.* 2019 Nov; 3\(11\):e10.5435.](#)

Behrbalk E, Salame K, Regev GJ, et al. Delayed diagnosis of cervical spondylotic myelopathy by primary care physicians. *Neurosurg Focus.* July 2013; 35(1):1-6.

Bennett AN, Marzo-Ortega H, Rehman A, et al. The evidence for whole-spine MRI in the assessment of axial spondyloarthropathy. *Rheumatology.* March 2010; 49(3):426-432. <https://doi.org/10.1093/rheumatology/kep427>.

<https://academic.oup.com/rheumatology/article/49/3/426/1787410>. Retrieved May 12, 2018.

[Bond A, Manian FA. Spinal epidural abscess: A review with special emphasis on earlier diagnosis. *Biomed Res Int.* 2016; \[2016:1614328\]\(#\).](#)

Budoff MJ, Khairallah W, Li D, et al. Trabecular bone mineral density measurement using thoracic and lumbar quantitative computed tomography. *Aca Radiology.* 2012; 19(2):179-183. doi: 10.1016/j.acra.2011.10.006.

[Choi BW, Song KJ, Chang H. Ossification of the posterior longitudinal ligament: A review of literature. *Asian Spine J.* 2011; 5\(4\):267–276. doi:10.4184/asj.2011.5.4.267](#)

Choosing Wisely®. Lists. <http://www.choosingwisely.org/clinician-lists/#keyword=spine&topic-area=Radiology&service=Imaging>. Retrieved March 27, 2018.

D' Alessandro. Does This Sacral Dimple Need to be Evaluated? PediatricEducation.org™. <https://pediatriceducation.org/2009/07/20/does-this-sacral-dimple-need-to-be-evaluated/>. Posted July 20, 2009. Retrieved March 29, 2018.

Eubanks JD. Cervical radiculopathy: Nonoperative management of neck pain and radicular symptoms. *Am Fam Physician*. 2010 Jan 1; 81(1):33-40.
<https://www.aafp.org/afp/2010/0101/p33.pdf>.

Graeber A, Cecava ND. Vertebral Osteomyelitis. [Updated 2019 Jun 3]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan.

Hou X, Sun C, Liu X, et al. Clinical features of thoracic spinal stenosis-associated myelopathy: A retrospective analysis of 427 cases. *Clin Spine Surg*. 2016 Mar; 29(2):86-9. doi: 10.1097/BSD.0000000000000081.

Jarvik JG, Gold LS, Comstock BA, et al. Association of early imaging for back pain with clinical outcomes in older adults. *JAMA*. 2015; 313(11):1143-1153. doi: 10.1001/jama.2015.1871.

Kim YS, Han IH, Lee IS, et al. Imaging findings of solitary spinal bony lesions and the differential diagnosis of benign and malignant lesions. *J Korean Neurosurg Soc*. August 2012; 52(2):126-132. doi: 10.3340/jkns.2012.52.2. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3467370/>. Retrieved March 29, 2018.

Koivikko MP, Koskinen SK. MRI of cervical spine injuries complicating ankylosing spondylitis. Skeletal Radiol. 2008 Sep; 37(9):813-9. doi:10.1007/s00256-008-0484-x. Epub 2008 Apr 18.

Last AR, Hulbert K. Chronic low back pain: Evaluation and management. *Am Fam Physician*. 2009; 79(12):1067-74.

Lerner S, Hartmann S, Barbagallo GMV, et al. Management of spinal infection: A review of the literature. *Acta Neurochir (Wien)*. 2018; 160(3):487-496. doi: 10.1007/s00701-018-3467-2.

Milhorat TH, Bolognese PA, Nishikawa M, et al. Association of Chiari malformation type I and tethered cord syndrome: Preliminary results of sectioning filum terminale. *Surg Neurol*. July 1, 2009; 72(1):20-35. <http://europepmc.org/abstract/med/19559924>. Retrieved March 29, 2018.

Muller D, Bauer JS, Zeile M, et al. Significance of sagittal reformations in routine thoracic and abdominal multislice CT studies for detecting osteoporotic fractures and other spine abnormalities. *Eur Radiol*. 2008; 18(8):1696-1702. doi: 10.10007/s00330-008-0920-2.

North American Spine Society (NASS). Clinician Lists. Choosing Wisely®.
<http://www.choosingwisely.org/clinician-lists/nass-emg-nerve-conduction-studies-to-determine-cause-of-spine-pain/>. Released October 9, 2013.

North American Spine Society (NASS). Diagnosis and treatment of cervical radiculopathy from degenerative disorders. 2010.

North American Spine Society (NASS). Five Things Physicians and Patients Should Question. Choosing Wisely®. <http://www.choosingwisely.org/doctor-patient-lists/north-american-spine-society/>. 2014.

Ostergaard M, Lambert RG. Imaging in ankylosing spondylitis. *Ther Adv Musculoskelet Dis*. August 2012; 4(4):301-311. doi: 10.1177/1759720X11436240. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3403247/>. Retrieved March 29, 2018.

Pomerantz SR. Myelography: Modern technique and indications. *Handb Clin Neurol*. 2016; 135:193-208.

Sieper J, Rudwaleit M, Baraliakos X, et al. The assessment of SpondyloArthritis international Society (ASAS) handbook: A guide to assess spondyloarthritis. *Ann Rheum Dis*. June 2009; 68 Suppl 2:ii1-44.

Stolper K, Haug JC, Christensen CT, et al. Prevalence of thoracic spine lesions masquerading as cauda equina syndrome: yield of a novel magnetic resonance imaging protocol. *Intern Emerg Med*. 2017 Dec; 12(8):1259-1264. doi: 10.1007/s11739-016-1565-9.

Strahle J, Smith BW, Martinez M, et al. The association between Chiari malformation Type I, spinal syrinx, and scoliosis. *J Neurosurg Pediatr*. 2015; 15:607-611. <http://thejns.org/doi/pdf/10.3171/2014.11.PEDS14135>. Retrieved March 29, 2018.

Trenga AP, Singla A, Feger MA, et al. Patterns of congenital bony spinal deformity and associated neural anomalies on X-ray and magnetic resonance imaging. *J Child Orthop*. August 2016; 10(4):343-352. doi: 10.1007/s11832-016-0752-6. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4940247/>. Retrieved March 29, 2018.

Vitzthum H, Dalitz K. Analysis of five specific scores of cervical spondylogenic myelopathy. *Eur Spine J*. December 2007; 16(12):2096-2103. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2140133/>.

Reviewed/Approved by *M. Atif Khalid MD* M. Atif Khalid, M.D., Medical Director, Radiology

Disclaimer: Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates ("Magellan"). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.