

<u>Subject:</u>	Enhanced External Counterpulsation in the Outpatient Setting	<u>Publish Date:</u>	12/18/2019
<u>Guideline #:</u>	CG-MED-86	<u>Last Review Date:</u>	11/07/2019
<u>Status:</u>	New		

Description

This document addresses the use of enhanced external counterpulsation (EECP) in the outpatient setting. EECP is commonly used to treat disabling, chronic, stable angina.

Clinical Indications**Medically Necessary:**

I. **A single* course of enhanced external counterpulsation (EECP) is considered medically necessary for individuals who have disabling, chronic, stable angina, (defined as Class III or Class IV Canadian Cardiovascular Society Classification [see definition section] angina or equivalent) when the criteria below are met:**

A. **Individuals refractive to optimal medical therapy and not readily amenable to surgical intervention such as percutaneous transluminal coronary angioplasty (PTCA) or cardiac bypass due to any of the following:**

- 1. Condition is inoperable; or**
- 2. High risk of operative complications or postoperative failure; or**
- 3. Coronary anatomy is not readily amenable to such procedures; or**
- 4. Has a comorbid state which creates excessive risk.**

Note: *A single course of treatment consists of a total of 35 hours of EECP; treatment is administered for one to two hours daily, 5 days a week, for approximately 3½ to 7 weeks.

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

II. A repeat course of EECP therapy is considered medically necessary in individuals who met the criteria in section (I) above, have chronic stable angina and who have objectively demonstrated a response to EECP. This would include those individuals who demonstrate one or more of the following:

- A. Early improvement in radionuclide stress perfusion imaging compared to a pre-EECP baseline;
or
- B. Reduction in antianginal medication use; or
- C. Improvement in exercise tolerance.

Not Medically Necessary:

The use of EECP is considered not medically necessary when the criteria above are not met and for all other indications, including, but not limited to:

- A. Aortic insufficiency (regurgitation might prevent diastolic augmentation);
- B. Arrhythmias such as atrial fibrillation, atrial flutter, ventricular tachycardia, and frequent premature ventricular beats (might interfere with the device's triggering mechanism);
- C. Bleeding diatheses;
- D. Cardiac catheterization within 2 weeks (may cause bleeding at the femoral puncture site);
- E. Congestive heart failure;
- F. Deep vein thrombosis, varicosities or stasis ulcers;
- G. Peripheral vascular disease or phlebitis (increased risk of thromboembolus);
- H. Pregnant women and women of childbearing potential who do not employ a reliable contraceptive method (possible danger to the fetus);
- I. Receiving Coumadin (warfarin) therapy with prothrombin time (PT) greater than 15 seconds or international normalized ratio (INR) greater than 2.0 (cuffs could cause bleeding in legs);
- J. Severe hypertension, greater than 180/110 mmHg, (treatment could produce diastolic blood pressure above acceptable limits);
- K. Stroke;
- L. Unstable angina.

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Coding

The following codes for treatments and procedures applicable to this guideline are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

CPT

92971

Cardioassist-method of circulatory assist; external [when specified as outpatient EECP]

HCPCS

G0166

External counterpulsation, per treatment session

ICD-10 Diagnosis

I20.0-I20.9

Angina pectoris

I25.110-I25.119

Atherosclerotic heart disease of native coronary artery with angina pectoris

I25.700-I25.709

Atherosclerosis of coronary artery bypass graft(s), unspecified, with angina pectoris

I25.710-I25.719

Atherosclerosis of autologous vein coronary artery bypass graft(s) with angina pectoris

I25.720-I25.729

Atherosclerosis of autologous artery coronary artery bypass graft(s) with angina pectoris

I25.730-I25.739

Atherosclerosis of nonautologous biological coronary artery bypass graft(s) with angina pectoris

I25.750-I25.759

Atherosclerosis of native coronary artery of transplanted heart with angina pectoris

I25.760-I25.769

Atherosclerosis of bypass graft of coronary artery of transplanted heart with angina pectoris

I25.790-I25.799

Atherosclerosis of other coronary artery bypass graft(s) with angina pectoris

Discussion/General Information

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Enhanced external counterpulsation (EECP) is a non-invasive outpatient treatment that uses a specialized pants-like device that incorporates three cuffs that contain air bladders in them. These pants are attached to an air compression system and electrocardiogram (ECG). The compressor works in conjunction with the ECG to trigger the compressor to inflate or deflate the air bladders in a sequential manner, coordinated with the diastolic (resting) phase of heart function. The sequential compression begins near the feet and progresses towards the torso in order to force blood out of the lower extremities and back into major blood vessels and the heart. This acts to increase blood pressure in the aorta during the diastole, increasing blood delivery to the coronary arteries and the heart muscle.

EECP is used to treat angina, a condition characterized by chest pain due to insufficient blood flow to the heart. Treatment with EECP is intended to provide symptomatic relief of angina, improve blood flow to blood-deprived areas of heart muscle, and to improve a person's level of physical activity. Additionally, EECP is believed to enhance the development of new blood vessels in the heart and increase flow through existing coronary blood vessels. A course of treatment typically includes 35 hours of EECP performed in 1- to 2-hour sessions at a time in the physician's office. EECP is not intended as a first-line therapy for angina, and is reserved for treating cases of angina which do not respond to medical and/or surgical therapy.

The use of EECP for the treatment of disabling, chronic, stable angina in people who are not suitable candidates for surgical intervention has been established in the medical evidence. Several large-scale prospective studies evaluating the efficacy of EECP in people with chronic stable angina demonstrate significant improvements in anginal symptoms, myocardial perfusion and output (Arora, 2002; Arora, 1999; Lawson, 2000; Loh, 2008; Soran, 2006; Urano, 2001; Weisfogel, 2001). One randomized, sham-controlled trial demonstrated significant improvement after 12 months of treatment for people who underwent a single 35-hour course of EEPC. In this study treatment-group, subjects reported significant improvements compared to sham treated individuals in all nine quality of life scales included on the Medical Outcomes Study SF-36 health survey, including the activities of daily living, ability to work, bodily pain, and others (Arora, 2002). Loh and colleagues reported on the results of a large case series study with 1427 participants who underwent a single course of 35 one-hour EECP treatments (2008). The authors reported that immediately following the treatment series, the proportion of subjects dropping at least one class on the

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Enhanced External Counterpulsation in the Outpatient Setting

Canadian Cardiovascular angina Classification system (CCS) was 77.9% and that 38.0% had dropped by 2 CCS classes. Based on participant reports, 76% of subjects experienced at least a 50% reduction in the frequency of angina symptoms. These improvements were durable and were noted to continue through to the three-year follow-up visits. There was also a significant decrease in calcium channel blocker and long-acting nitrate use at all follow-up points.

EECP has also been studied for the treatment of congestive heart failure. In 2002, Soran and colleagues reported on a feasibility study of EECP, as a treatment for congestive heart failure in 26 subjects. In this uncontrolled study, the subjects were treated with 35 daily, one-hour sessions and followed for 6 months after completion of the course of therapy. The study suggests that the treatment was safe and well tolerated.

Results from the PEECH trial (Prospective Evaluation of EECP in Congestive Heart Failure) have been published (Feldman, 2006). The article discusses the 6-month outcomes of the PEECH trial which was a randomized controlled trial of 187 subjects with mild to moderate symptoms of heart failure. Participants were selected to receive either EECP and pharmacologic therapy or pharmacologic therapy alone. At 6 months, the study identified significant improvement in the EECP group compared to the control group in exercise time (also 35% increased exercise time by at least 60 seconds compared to 25% of controls), and NYHA functional class. No significant differences were identified in peak volume oxygen uptake (VO₂), respiratory exchange rate, or measurements on the Minnesota Living with Heart Failure or Borg scales. While the results of this study are promising, small numbers of participants (71 subjects completed the EECP regime), a high dropout rate in the experimental group due to adverse experiences (24% vs. 14% in the control group), lack of blinding which the authors acknowledge may have allowed for a placebo effect and lack of a sham treatment group, hinder the value of the study results. Finally, a subgroup analysis of the study data found that significant improvement in exercise time was seen only in the subjects with ischemic cardiomyopathy (approximately 40 seconds compared to controls), and not in those with non-ischemic cardiomyopathy.

In a separate report, Abbottsmith and colleagues report on a subgroup analysis of the PEECH trial (2006). Their report focused on PEECH participants greater than 64 years of age. The study results indicated that at 6 months, exercise responder rate and peak VO₂ were significantly improved compared to the control

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Enhanced External Counterpulsation in the Outpatient Setting

group. NYHA classification was significantly improved compared to controls at 3 months follow-up but not at 6 months and Minnesota Living with Heart Failure scores were not found to be significantly different. Again, while these results are promising, the same limitations of the larger PEECH trial affect these subgroup results. The authors warn that, as a subgroup analysis, their data should be interpreted with caution, but that it may be of benefit in identifying subgroups in which additional research should be targeted. Data from a larger, blinded randomized controlled trial are needed to validate the PEECH trial findings.

In 2014, the American College of Cardiology/American Heart Association/American Association for Thoracic Surgery/Preventive Cardiovascular Nurses Association/Society for Cardiovascular Angiography and Interventions/Society of Thoracic Surgeons focused update of the Guideline for the Diagnosis and Management of Patients With Stable Ischemic Heart Disease (SIHD) issued recommendations for EECP for relief of refractory angina in patients with SIHD (Class IIb Level of Evidence B). A class IIb, level of evidence B recommendation indicates the procedure/treatment may be considered. The benefit is equal to or greater than the risk. Additional studies with broad objectives are needed, and additional registry data would be helpful. The usefulness/efficacy is less well established, and greater conflicting evidence from single randomized trials or nonrandomized studies exists (Fihn, 2014).

The 2015 ACC/AHA/Society for Cardiovascular Angiography and Interventions (SCAI) focused update to the 2011 practice guideline for percutaneous coronary intervention, does not does not address EECP procedure/treatment. (Levine, 2015).

The evidence regarding the use of EECP for other indications, including other anginal or cardiac conditions, such as non-disabling stable angina, unstable angina, CHF or stroke is currently insufficient to allow conclusions to be made.

Definitions

Angina: A condition characterized by chest pain, due to insufficient blood flow to the heart.

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Aortic Insufficiency: A condition caused by back flow of blood from the aorta into the left ventricle of the heart in between beats because of failure of an aortic valve to close properly.

Arrhythmia: A condition caused by an abnormal heartbeat; includes atrial fibrillation, atrial flutter, ventricular tachycardia, and frequent premature ventricular beats.

Bleeding diathesis: The occurrence of excessive bleeding.

Canadian Cardiovascular Society Score: This organization defines anginal classes as follows:

- **Class I - Ordinary physical activity does not cause angina;**
- **Class II - Slight limitation of ordinary activity;**
- **Class III - Marked limitation of ordinary physical activity;**
- **Class IV - Inability to carry on physical activity without discomfort.**

Cardiac Bypass: A surgical procedure that re-routes the blood flow supplying the heart to prevent heart attacks.

Cardiac Catheterization: A diagnostic procedure that involves passing a catheter (i.e., a thin flexible tube) through an artery or a vein to the heart, and into a coronary artery; dye that can be seen on x-rays is then injected into the heart to produce angiograms (i.e., x-ray images) of the coronary arteries and the left ventricle, the heart's main pumping chamber; it may also be used to measure pressures in the pulmonary artery and to monitor heart function, usually in critically ill patients.

Comorbid States: Diseases that exist at the same time as other diseases within the same individual; for instance, if an individual has both diabetes and high blood pressure, they are considered comorbid conditions.

Congestive Heart Failure: A condition caused by a weakening of the heart muscle where the heart is unable to adequately pump blood to the body.

Deep Vein thrombosis: A condition characterized by blood clots forming in the main veins of the legs when a vein is damaged or if the flow of blood slows down or stops; clots may become loosened and travel to the lungs causing a pulmonary embolus, a life-threatening condition.

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Dyspnea: Shortness of breath.

Enhanced external counterpulsation (EECP): A device that squeezes the leg, and in some circumstances the buttocks, to increase blood and fluid flow back to the heart.

Hypertension: High blood pressure.

Palpitation: An abnormally fast heartbeat.

Percutaneous Transluminal Coronary Angioplasty (PTCA): Also known as coronary artery balloon dilation or balloon angioplasty; a surgical procedure where a long thin surgical tool is passed through the blood vessels to the main arteries that supply blood to the heart; the tools are then used to widen the blood vessels to improve blood flow.

Peripheral Vascular Disease: A common circulation problem in which the arteries that carry blood to the legs or arms become narrowed or clogged causing cold hands and feet, nerve problems, and problems with healing.

Phlebitis: Inflammation of a vein resulting in decreased or obstructed blood flow through the vessel.

Stasis Ulcers: A condition characterized by the disintegration of skin due to the buildup of fluid in the tissues that do not have enough oxygen to support adequate tissue function.

Varicosities: Abnormally, markedly swollen or dilated veins, as occurs with poor circulation to the limbs.

References

Peer Reviewed Publications:

1. **Abbottsmith CW, Chung ES, Varricchione T, et al. Prospective Evaluation of EECP in Congestive Heart Failure Investigators. Enhanced external counterpulsation improves exercise duration and peak oxygen**

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

consumption in older patients with heart failure: a subgroup analysis of the PEECH trial. Congest Heart Fail. 2006; 12(6):307-311.

2. Arora RR, Chou TM, Jain D, et al. Effects of enhanced external counterpulsation on Health-Related Quality of Life continue 12 months after treatment: a substudy of the Multicenter Study of Enhanced External Counterpulsation. J Investig Med. 2002; 50(1):25-32.

3. Arora RR, Chou TM, Jain D, et al. The multicenter study of enhanced external counterpulsation (MUST-EECP): effect of EECP on exercise-induced myocardial ischemia and anginal episodes. J Am Coll Cardiol. 1999; 33(7):1833-1840.

4. Braith RW, Conti CR, Nichols WW, et al. Enhanced external counterpulsation improves peripheral artery flow-mediated dilation in patients with chronic angina: a randomized sham-controlled study. Circulation. 2010; 122(16):1612-1620.

5. Feldman AM, Silver MA, Francis GS, et al. PEECH Investigators. Enhanced external counterpulsation improves exercise tolerance in patients with chronic heart failure. J Am Coll Cardiol. 2006; 48(6):1198-1205.

6. Feldman AM, Silver MA, Francis GS, et al. Treating heart failure with enhanced external counterpulsation (EECP): design of the Prospective Evaluation of EECP in Heart Failure (PEECH) Trial. J Card Fail. 2005; 11(3):240-245.

7. Kumar A, Aronow WS, Vadnerkar A, et al. Effect of enhanced external counterpulsation on clinical symptoms, quality of life, 6-minute walking distance, and echocardiographic measurements of left ventricular systolic and diastolic function after 35 days of treatment and at 1-year follow up in 47 patients with chronic refractory angina pectoris. Am J Ther. 2009; 16(2):116-118.

8. Lawson WE, Hui JC, Cohn PF. Long-term prognosis of patients with angina treated with enhanced external counterpulsation: five-year follow-up study. Clin Cardiol. 2000; 23(4):254-258.

9. Lawson WE, Hui JC, Lang G. Treatment benefit in the enhanced external counterpulsation consortium. Cardiology. 2000; 94(1):31-35.

10. Lin S, Liu M, Wu B, et al. External counterpulsation for acute ischaemic stroke. Cochrane Database Syst Rev. 2012;1:CD009264.

11. Loh PH, Cleland JG, Louis AA, et al. Enhanced external counterpulsation in the treatment of chronic refractory angina: a long-term follow-up outcome from the International Enhanced External Counterpulsation Patient Registry. Clin Cardiol. 2008; 31(4):159-164.

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Enhanced External Counterpulsation in the Outpatient Setting

12. **Masuda D, Nohara R, Hiari T, et al. Enhanced external counterpulsation improved myocardial perfusion and coronary flow reserve in patients with chronic stable angina. Eur Heart J. 2001; 22(16):1451-1458.**
13. **McCullough PA, Henry TD, Kennard ED, et al.; IEPR Investigators. Residual high-grade angina after enhanced external counterpulsation therapy. Cardiovasc Revasc Med. 2007; 8(3):161-165.**
14. **Michaels AD, Accad M, Ports TA, Grossman W. Left ventricular systolic unloading and augmentation of intracoronary pressure and Doppler flow during enhanced external counterpulsation. Circulation. 2002; 106(10):1237-1242.**
15. **Michaels AD, Kennard ED, Kelsey SE, et al. Does higher diastolic augmentation predict clinical benefit from enhanced external counterpulsation?: Data from the international EECP patient registry (IEPR). Clin Cardiol. 2001; 24(6):453-458.**
16. **Michaels AD, Raisinghani A, Soran O, et al. The effects of enhanced external counterpulsation on myocardial perfusion in patients with stable angina: a multicenter radionuclide study. Am Heart J. 2005; 150(5):1066-1073.**
17. **Shah SA, Shapiro RJ, Mehta R, Snyder JA. Impact of enhanced external counterpulsation on Canadian Cardiovascular Society angina class in patients with chronic stable angina: a meta-analysis. Pharmacotherapy. 2010; 30(7):639-645.**
18. **Shakouri SK, Razavi Z, Eslamian F, et al. Effect of enhanced external counterpulsation and cardiac rehabilitation on quality of life, plasma nitric oxide, endothelin 1 and high sensitive CRP in patients with coronary artery disease: a pilot study. Ann Rehabil Med. 2015; 39(2):191-198.**
19. **Soran O, Crawford LE, Schneider VM, Feldman AM. Enhanced external counterpulsation in the management of patients with cardiovascular disease. Clin Cardiol. 1999; 22(3):173-178.**
20. **Soran O, Fleishman B, Demarco T., et al. Enhanced external counterpulsation in patients with heart failure: A multicenter feasibility study. Congestive Heart Failure. 2002; 8(6):204-208.**
21. **Soran O, Kennard ED, Kfouri AG, Kelsey SF; IEPR Investigators. Two-year clinical outcomes after enhanced external counterpulsation (EECP) therapy in patients with refractory angina pectoris and left ventricular dysfunction (report from The International EECP Patient Registry). Am J Cardiol. 2006; 97(1):17-20.**
22. **Springer S, Fife A, Lawson W, et al. Psychosocial effects of enhanced external counterpulsation in the angina patient: a second study. Psychosomatics. 2001; 42(2):124-132.**

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Enhanced External Counterpulsation in the Outpatient Setting

23. Tartaglia J, Stenerson J Jr, Charney R, et al. Exercise capability and myocardial perfusion in chronic angina patients treated with enhanced external counterpulsation. Clin Cardiol. 2003; 26(6):287-290.
24. Urano H, Ikeda H, Ueno T, et al. Enhanced external counterpulsation improves exercise tolerance, reduces exercise-induced myocardial ischemia and improved left ventricular diastolic filling in patients with coronary artery disease. J Am Coll Cardiol. 2001; 37(1):93-99.
25. Vijayaraghavan K, Santora L, Kahn J, et al. New graduated pressure regimen for external counterpulsation reduces mortality and improves outcomes in congestive heart failure: a report from the Cardiomedics External Counterpulsation Patient Registry. Congest Heart Fail. 2005; 11(3):147-152.
26. Zhang C1, Liu X, Wang X, Wang Q, Zhang Y, Ge Z. Efficacy of enhanced external counterpulsation in patients with chronic refractory angina on Canadian Cardiovascular Society (CCS) Angina Class: An updated meta-analysis.

Government Agency, Medical Society, and Other Authoritative Publications:

1. Amsterdam EA, Weger N, Peterson ED, et al. 2014 AHA/ACC Guideline for the management of patients with non-ST-elevation acute coronary syndromes. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. Developed in collaboration with the Society for Cardiovascular Angiography and Interventions and Society of Thoracic Surgeons. J Am Coll Cardiol. 2014; 64(24): e139-e228. Available at: http://www.onlinejacc.org/content/accj/64/24/e139.full.pdf?_ga=2.29302203.683828590.1568648102-1048116366.1542209605. Accessed on September 16, 2019.
2. Canadian Cardiovascular Society. Society grading of Anginal pectoris. Last updated 1976. Available at: http://www.ccs.ca/images/Guidelines/Guidelines_POS_Library/Ang_Gui_1976.pdf. Accessed on October 14, 2019.
3. Centers for Medicare and Medicaid Services National Coverage Determination for External Counterpulsation (ECP) for Severe Angina. NCD #20.20. Effective March 20, 2006. http://www.cms.hhs.gov/mcd/index_list.asp?list_type=ncd. Accessed on September 16, 2019.
4. Fihn SD, Blankenship JC, Naidu SS, et al. 2014 ACC/AHA/AATS/PCNA/SCAI/STS Focused update of the guideline for the diagnosis and management of patients with stable ischemic heart disease. A report of the American College of Cardiology/ American Heart Association Task Force on Practice Guidelines, and the American Association for Thoracic Surgery, Prevention Cardiovascular Nurses Association,

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Enhanced External Counterpulsation in the Outpatient Setting

Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. J Am Coll Cardiol. 2014; 64(18):1929-1949. Available at: http://www.onlinejacc.org/content/accj/64/18/1929.full.pdf?_ga=2.40777125.683828590.1568648102-1048116366.1542209605. Accessed on September 16, 2019.

5. Yancy CW, Jessup M, Masoudi FA, et al. 2017 ACC/AHA/HFSA Focused update of the 2013 ACCF/AHA guideline for the management of heart failure. A report of the American College/ American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. J Am Coll Cardiol. 2017; 70(6):776-803. Available at: http://www.onlinejacc.org/content/accj/70/6/776.full.pdf?_ga=2.37745831.683828590.1568648102-1048116366.1542209605. Accessed on September 16, 2019.

Websites for Additional Information

- American Heart Association. Angina pectoris. Reviewed July 31, 2015. Available at: <https://www.heart.org/en/health-topics/heart-attack/angina-chest-pain/angina-pectoris-stable-angina>. Accessed on September 16, 2019.**
- National Heart, Lung, and Blood Institute. Angina. Reviewed December 10, 2018. Available at: http://www.nhlbi.nih.gov/health/dci/Diseases/Angina/Angina_WhatIs.html. Accessed on September 16, 2019.**

Index

The use of specific product names is illustrative only. It is not intended to be a recommendation of one product over another, and is not intended to represent a complete listing of all products available.

History

Status	Date	Action
New	11/07/2019	Medical Policy & Technology Assessment Committee (MPTAC) review. Initial document development.

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

DRAFT

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.