

*National Imaging Associates, Inc.	
Clinical guidelines OUTPATIENT HABILITATIVE AND REHABILITATIVE SPEECH THERAPY	Original Date: November 2015
Physical Medicine – Clinical Decision Making	Last Revised Date: December December 2023 2
Guideline Number: NIA_CG_602	Implementation Date: July 2024 3

Table of Contents

GENERAL INFORMATION	2
STATEMENT	2
PURPOSE.....	2
SCOPE	2
MEDICAL NECESSITY	3
DOCUMENTATION.....	3
EVALUATION	6
TREATMENT GOALS.....	9
FREQUENCY AND DURATION	10
DISCONTINUATION OF TREATMENT	11
OTHER CONSIDERATIONS	12
BACKGROUND	14
DEFINITIONS	18
REFERENCES.....	23

General Information

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

Policy Statement

~~Habilitative speech therapy services may or may not be covered by all clients of this organization. If the service is covered, it may or may not require prior authorization. These guidelines apply to all markets and populations, including teletherapy, contracted with this organization through the corresponding state health plans unless a market-specific health plan has been developed. These services must be provided by a skilled and licensed therapy practitioner and in a manner that is in accordance with accepted standards of practice for discipline-specific therapies. It must also be clinically appropriate in amount, duration, and scope to achieve their purpose and considered effective treatment for the current injury, illness, or condition.~~

Habilitative/Rehabilitative speech therapy should meet the definitions below, be provided in a clinic, an office, at home, or in an outpatient setting, and be ordered by either a primary care practitioner or specialist.

Purpose

This guideline describes the documentation requirements of appropriate Habilitative/Rehabilitative Speech Therapy.

All recommendations in this guideline reflect practices that are evidence-based and/or supported by broadly accepted clinical specialty standards.

Scope

This guideline applies to all physical medicine practitioners, including Speech-Language Pathologists (SLP) and Speech-Language Pathology Assistants (SLP-A).

Definition

Habilitative Speech Therapy

~~Treatment provided by a state-regulated speech therapist to help a person attain, maintain, or prevent deterioration of a skill or function never learned or acquired. There must be measurable improvement and progress towards functional goals within an anticipated timeframe toward a patient's maximum potential. Treatment may also be appropriate in a child with a progressive disorder when it has the potential to prevent the loss of a functional skill or enhance the adaptation to such functional loss. The condition must be such that there is a~~

~~reasonable expectation that the services will bring about a significant improvement within a reasonable time frame, regardless of whether the individual has a coexisting disorder. Ongoing treatment is not appropriate when functioning is steady and treatment no longer yields measurable functional progress.~~

Rehabilitative Speech Therapy

~~Treatment provided by a state-regulated speech therapist designed to help a person recover from an acute injury or exacerbation of a chronic condition that has resulted in a decline in functional performance. The specific impact of injury or exacerbation on the patient's ability to perform in their everyday environment must be supported by appropriate tests and measures in addition to clinical observations. Services must be provided within a reasonable time frame (frequency/duration) to restore lost function or to teach compensatory techniques if full recovery of function is not possible.~~

Functional Skills

~~They are considered necessary communication activities of daily life. The initial plan of care documents baseline impairments as they relate to functional communication with specific goals developed that are measurable, sustainable and time-specific. Subsequent plans of care document progress toward attainment of these goals in perspective to the patients' potential ability. Discontinuation of therapy will be expected when the maximum therapeutic value of a treatment plan has been achieved, no additional functional improvement is apparent or expected to occur, and the provision of services for a condition ceases to be of therapeutic value.~~

National Imaging Associates will review all requests resulting in adverse determinations for Medicaid members for coverage under federal Early and Periodic Screening, Diagnostic and Treatment (EPSDT) guidelines. [1, 2]

Medical Necessity Requirements

The following criteria must be addressed to justify the medical necessity of the prescribed treatment.

Documentation

Progress notes or /updated plans of care that cover the patient's specific progress towards their goals with review by the primary care practitioner or other **non-physician practitioner (NPP)** will be required every 60-90 days or per state guidelines.

Documentation should include: [3]:

- Written referral from primary care practitioner or other non-physician practitioner (NPP) as required permitted by state guidelines.

- Patient's current level of function and any conditions that are impacting his/her ability to benefit from skilled intervention.
- Objective measures of the patient's overall functional progress relative to each treatment goal as well as a comparison to the previous progress report
- Skilled treatment techniques that are being utilized in therapy as well as the patient's response to therapy.
- If appropriate, documentation should provide a rationale for lack of progress or response to treatment ~~and why there may be a lack thereof~~
- Treatment goals that follow a hierarchy of complexity to achieve the target skills for a functional goal.
- Re-evaluation ~~or~~ annual testing (for habilitative therapy) using formal standardized assessment tools and formal assessment of progress must be performed to support progress, ongoing delays and medical necessity for continued services.
 - An explanation of any significant changes in the plan of care and clinical rationale for why the ongoing skills of a SLP are medically necessary.

When ~~s~~When skilled services are also being provided by other community service agencies and/or school systems the notes must show; ~~Frequency and duration of skilled services must also be in accordance with the following:~~

Intense frequencies (3x/week or more) will require additional documentation and testing supporting a medical need to achieve an identified new skill or recover function with specific, achievable goals within the requested intensive period^a. Details on why a higher frequency is more beneficial than a moderate or low frequency must be included. Higher frequencies may be considered when delays are classified as severe as indicated by corresponding testing guidelines used in the evaluation. More intensive frequencies may be necessary in the acute phase, however, progressive decline in frequency is expected within a reasonable time frame. Moderate frequency (2x/week) should be consistent with moderate delays as established in the general guidelines of formal assessments used in the evaluation. This frequency may be used for ongoing care when documentation supports this frequency as being clinically effective toward achieving the functional goals in the treatment plan within a reasonable time frame. Low frequency (1x/week or less) may be considered when testing guidelines indicate mild delays or when a higher frequency has not been clinically effective, and a similar outcome is likely with less treatment per week.

All requested frequencies must be supported by skilled treatment interventions regardless of level of severity of delay.

Additional factors may be considered on a case by case basis.

There must be evidence as to whether the services are considered reasonable, effective, and of such a complex nature that they require the technical knowledge and clinical decision-making skill of a therapist or whether they can be safely and effectively carried out by non-skilled personnel without the supervision of qualified professionals.

Treatment that requires the technical knowledge and clinical decision-making expertise to meet the skilled service needs of the individual. This includes analyzing medical/behavioral data and

selecting appropriate evaluation tools/protocols to determine communication/swallowing diagnosis and prognosis.

Progress notes/updated plans of care that cover the patient's specific progress towards their goals with review by the primary care practitioner or other NPP will be required every 60-90 days or per state guidelines. Documentation should include:

The patient's current level of function, any conditions that are impacting his/her ability to benefit from skilled intervention.

Objective measures of the patient's overall functional progress relative to each treatment goal as well as a comparison to the previous progress report.

Skilled treatment techniques that are being utilized in therapy as well as the patient's response to therapy and why there may be a lack thereof. Treatment goals that follow a hierarchy of complexity to achieve the target skills for a functional goal.

Re-evaluation/annual testing (for habilitative therapy) using formal standardized assessment tools and formal assessment of progress must be performed to support progress, ongoing delays and medical necessity for continued services.

An explanation of any significant changes in the plan of care and clinical rationale for why the ongoing skills of a SLP are medically necessary.

If the patient is not progressing, then documentation of a revised treatment plan is necessary. Discontinuation of therapy will be expected when the maximum therapeutic value of a treatment plan has been achieved, no additional functional improvement is apparent or expected to occur, and the provision of services for a condition cease to be of therapeutic value.

- It is expected that a specific discharge plan, with the expected treatment frequency and duration, must be included in the plan of care. The discharge plan must indicate the plan to wean services once the patient has attained their goals, if no measurable functional improvement has been demonstrated, or if the program can be carried out by caregivers or other non-skilled personnel.

It is expected that there be evidence of the development of age-appropriate home regimen to facilitate carry-over of target skills and strategies and education of patient, family, and caregiver in home practice exercises, self-monitoring as well as indication of compliance for maximum benefit of therapy.

For patients no longer showing functional improvement, a weaning process of one to two months should occur. Behaviors that interfere with the ability to progress with therapy qualify under the ASHA discharge criteria guidelines.¹¹ If the patient shows signs of regression in function, the need for skilled speech therapy can be re-evaluated at that time. Periodic episodes of care may be needed over a lifetime to address specific needs or changes in condition resulting in functional decline.

A maintenance level of therapy services may be considered when a member requires skilled therapy for ongoing periodic assessments and consultations and the member meets one of the following criteria:

Documentation shows the member and the responsible adult have a continuing need for education, or a periodic adjustment of the home program is needed to meet the member's needs.

Goals in the plan of care must be updated to reflect that care is focused on maintaining the current level of functioning and preventing regression, rather than progressing or improving function.

Clear documentation of the skilled interventions rendered and objective details of how these interventions are preventing deterioration or making the condition more tolerable must be provided. The notes must also clearly demonstrate that the specialized judgment, knowledge, and skills of a qualified therapist (as opposed to a non-skilled individual) are required for the safe and effective performance of services in a maintenance program.

- Documentation should include any applicable coordination of services with those agencies
- When services are not available, this should be indicated in the documentation

Evaluation

- Establishing a delay or deficit

Formal testing [4, 3]

- Age-appropriate, norm-referenced, standardized, and specific to the therapy
- Provided
- Different tests use different scoring methods and risk categories; any selected test must be interpreted in accordance with its scoring method.
- Test scores and interpretation should establish the presence of a significant delay
- While standardized testing is preferred, scores alone may not be used as the sole criteria for determining a patient's medical need for skilled intervention; test information must be linked to difficulty with or inability to perform everyday tasks [5, 6].
- In the absence of standardized testing or when test scores show skills within normal ranges, the documentation must include detailed clinical observations and objective data to document the degree and severity of the condition to support the medical need for skilled services; a caregiver interview/questionnaire can also support the request.
- Any time standardized testing cannot be completed, the documentation must clearly state the reason formal testing could not be done.

Evaluation for Services must be considered reasonable, effective, and of such a complex nature that they require the technical knowledge and clinical decision-making skill of a

therapist or can be safely and effectively carried out conducted by non-skilled personnel without the supervision of qualified professionals

- hHabilitative tTherapy should include:
 - Tthere is a reasonable expectation that the services will bring about a significant improvement within a reasonable time frame, regardless of whether the individual has a coexisting disorder.
 - Evidence that Ongoing treatment is not appropriate; note: ongoing treatment is not appropriate when patient -when-functioning is steady and treatment no longer yields measurable and -significant functional progress.
- Evaluation for rRehabilitative tTherapy should include:
 - The specific impact or exacerbation of injury on the patient's ability to perform in their everyday environment must be supported by appropriate tests and measures in addition to clinical observations.
- Functional Skills
 - The initial plan of care must document baseline impairments as they relate to functional communication and feeding/swallowing with specific goals developed that are measurable, sustainable and time-specific.

Establishing a delay or deficit There must be evidence as to whether the services are considered reasonable, effective, and of such a complex nature that they require the technical knowledge and clinical decision-making skill of a therapist or whether they can be safely and effectively carried out by non-skilled personnel without the supervision of qualified professionals.

Treatment that requires the technical knowledge and clinical decision-making expertise to meet the skilled service needs of the individual. This includes analyzing medical/behavioral data and selecting appropriate evaluation tools/protocols to determine communication/swallowing diagnosis and prognosis.

Progress notes/updated plans of care that cover the patient's specific progress towards their goals with review by the primary care practitioner or other NPP will be required every 60-90 days or per state guidelines. Documentation should include:

The patient's current level of function, any conditions that are impacting his/her ability to benefit from skilled intervention.

Objective measures of the patient's overall functional progress relative to each treatment goal as well as a comparison to the previous progress report.

Skilled treatment techniques that are being utilized in therapy as well as the patient's response to therapy and why there may be a lack thereof. Treatment goals that follow a hierarchy of complexity to achieve the target skills for a functional goal.

Re-evaluation/annual testing (for habilitative therapy) using formal standardized assessment tools and formal assessment of progress must be performed to support progress, ongoing delays and medical necessity for continued services.

An explanation of any significant changes in the plan of care and clinical rationale for why the ongoing skills of a SLP are medically necessary.

Formal testing [4, 3]:

must be a Age-appropriate, norm-referenced, standardized, and specific to the therapy provided Test scores should meet the following criteria to establish presence of a functional delay. Notes should document the following to establish the presence of delays or deficits:-

The following methods are generally accepted measures that may be considered to support a significant delay:

Standardized scores at or below the 10th percentile in at least one subtest area for the patient's age.3-

Standardized scores greater than or equal to 1.5 standard deviations below the mean in at least one subtest area for the patient's age.1-8

Functional delays may be established by 25% or greater deficit in age equivalency as indicated by established general guidelines of functional assessments or specific criterion-referenced tests or profiles.1-6 Different tests use different scoring methods and risk categories; a. Any selected test should be applied in accordance with its scoring method

While standardized testing is preferred, scores alone may not be used as the sole criteria for determining a patient's medical need for skilled intervention; t. Test information must be linked to difficulty with or inability to perform everyday tasks [5, 6].

In the absence of standardized testing or when test scores show skills within normal ranges despite functional deficits, the documentation must include detailed clinical observations and objective data to document the degree and severity of the condition, in order to support the medical need for skilled services; a. A caregiver interview/questionnaire can also support the request

Any time standardized testing cannot be completed, the documentation must clearly state the reason formal testing could not be done

Treatment Goals

- Treatment goals must be:
 - Realistic, measurable, and promote attainment of developmental milestones and functional communication abilities appropriate to the patient's age and circumstances [7, 4].
 - They should include the type, amount, duration, and frequency of therapy services.⁹
 - The amount, frequency, and duration of the services These must be consistent with accepted standards of practice and correspond be commensurate corresponding with the patient's medical and skilled therapy needs and level of disability.
 - Treatment goals must be individualized and measurable in order to identify the functional levels related to appropriate maintenance or maximum therapeutic benefit, targeted to identified functional deficits, and promote the attainment of; Goals of intervention should target the functional deficits identified by the skilled therapist during the assessment and promote attainment of:
 - Age-appropriate developmental milestones
 - Functional skills appropriate to the patient's age and circumstances Although identified as component parts of participation, underlying factors, performance skills, client factors or the environment should not be the targeted outcome of long-term goals. For sustained positive benefits from therapeutic interventions, activities can be practiced in the

child's environment and reinforced by the parents or other caregivers.
Practice in one's natural environment is essential for success⁻¹⁰

- Although identified as component parts of participation, underlying factors, performance skills, client factors and/or the environment should not be the targeted outcome of long-term goals.
- Services must be considered reasonable, effective, and of such a complex nature that they require the technical knowledge and clinical decision-making skill of a therapist or can be safely and effectively conducted by non-skilled personnel without the supervision of qualified professionals.
- For sustained positive benefits from therapeutic interventions, activities can be practiced in the child's environment and reinforced by the parents or other caregivers.

The plan of care must include goals detailing type, amount, duration, and frequency of therapy services required to achieve targeted outcomes. The frequency and duration must also be commensurate with the patient's level of disability, medical and skilled therapy needs, as well as accepted standards of practice while reflecting clinical reasoning and current evidence.⁹

Frequency and DurationFrequency and duration of skilled services must also be in accordance with the following:

- All requested frequencies must be supported by skilled treatment interventions regardless of level of severity of delay.
- Intense frequencies (i.e., 3x/week or more) will will require additional documentation and testing supporting a medical need to achieve an identified new skill or recover function with specific, achievable goals within the requested intensive period.⁻⁹
 - Details on why a higher frequency is more beneficial than a moderate or low frequency must be included. Higher frequencies may be considered when delays are classified as severe (as indicated by corresponding testing guidelines used in the evaluation) [4].
 - More intensive frequencies may be necessary in the acute phase, however, progressive decline in frequency is expected within a reasonable time frame [4].
- Moderate frequency (i.e., 2x/week) should be consistent with moderate delays as established in the general guidelines of formal assessments used in the evaluation.
 - This frequency may be used for ongoing care when documentation supports this frequency as being clinically effective toward achieving the functional goals in the treatment plan within a reasonable time frame.
- Low frequency (i.e., 1x/week or less) may be considered when testing guidelines indicate mild delays or when a higher frequency has not been clinically effective, and a similar outcome is likely with less treatment per week.
- All requested frequencies must be supported by skilled treatment interventions regardless of level of severity of delay.

- Additional factors may be considered on a case-by-case basis.
- If the patient is not progressing, documentation of a revised treatment plan is necessary.
- ~~MA~~ maintenance level of therapy services may be considered when a member requires skilled therapy for ongoing periodic assessments and consultations and the member meets one of the following criteria:
- Documentation shows the member and the responsible adult have a continuing need for education, or a periodic adjustment of the home program is needed to meet the member's needs.
- - Goals in the plan of care must be updated to reflect that care is focused on maintaining the current level of functioning and preventing regression, rather than progressing or improving function.
 - Clear documentation of the skilled interventions rendered and objective details of how these interventions are preventing deterioration or making the condition more tolerable must be provided. The notes must also clearly demonstrate that the specialized judgment, knowledge, and skills of a qualified therapist (as opposed to a non-skilled individual) are required for the safe and effective performance of services in a maintenance program.
 - It is expected that there be evidence of the development of age-appropriate home regimen to facilitate carry-over of target skills and strategies and education of patient, family, and caregiver in home practice exercises, self-monitoring as well as indication of compliance for maximum benefit of therapy.
 - Goals in the plan of care must be updated to reflect that care is focused on maintaining the current level of functioning and preventing regression, rather than progressing or improving function.
 - Clear documentation of the skilled interventions rendered and objective details of how these interventions are preventing deterioration or making the condition more tolerable must be provided. The notes must also clearly demonstrate that the specialized judgment, knowledge, and skills of a qualified therapist (as opposed to a non-skilled individual) are required for the safe and effective performance of services in a maintenance program.

Discontinuation of Treatment

- Alt is expected that a specific discharge plan, with the expected treatment frequency and duration is, must be included in the plan of care [7]. The discharge plan must indicate the plan to wean services once the patient has attained their goals, if [5].
- De ischarge may also be warranted if:
 - No measurable functional improvement has been demonstrated.
 - , ~~Wa weaning process of one to two months should occur~~

- Behaviors that interfere with the ability to progress with therapy qualify under the [American Speech-Language-Hearing Association \(ASHA\)](#) discharge criteria guidelines.
 - ~~or if the p~~rogram can be ~~carried out~~conducted by caregivers or other non-skilled personnel.
 - ~~Mthe m~~aximum therapeutic value of a treatment plan has been achieved.
 - ~~n~~No additional functional improvement is apparent or expected to occur.
 - ~~the p~~rovision of services for a condition cease to be of therapeutic value.
- ~~For patients no longer showing functional improvement, a weaning process of one to two months should occur. Behaviors that interfere with the ability to progress with therapy qualify under the ASHA discharge criteria guidelines.⁴¹~~If the patient shows signs of regression in function, the need for skilled speech therapy can be re-evaluated at that time.
 - Periodic episodes of care may be needed over a lifetime to address specific needs or changes in condition resulting in functional decline.

* A weaning process of one or two months should be implemented.

Other Considerations

- ~~When a For patient's whose~~language background differs from the rendering therapist and ~~in situations in which a clinician who has~~with native or near-native proficiency in the target language is not available, use of an interpreter is appropriate and should be documented accordingly.
 - If an interpreter is not present, ~~rationale for this should be documented as well as documentation that provides evidence along with evidence of a communication disorder,~~ and a treatment plan that supports linguistically appropriate services without the use of an interpreter.
- ~~If Further,~~if a patient is substantially exposed to more than one language, the assessment must evaluate both languages and contain appropriate tests and measures to clearly denote the presence that a communication disorder is present as opposed to normal linguistic variations related to second language learning [8].
- Swallowing disorders (dysphagia) and feeding disorders will need documentation of an oral, pharyngeal, and/or esophageal phase disorder, food intolerance or aversion [4, 5, 7].
 - -There must be evidence of ongoing progress and a consistent home regimen to facilitate carry-over of target feeding skills, strategies and education of patient, family, and caregiver.
 - Therapies for picky eaters who can eat and swallow normally, are meeting growth and developmental milestones, eat at least one food from all major food groups (protein, grains, fruits, etc.) and more than 20 different foods are ~~is not~~ medically necessary.
- Treatment that includes goals for reading/literacy must also have a primary diagnosis of a speech or language disorder.

- Documentation must support that the deficits in reading/literacy are affecting functional activities of daily living and are not the primary focus of treatment.—. They must show how the services for reading/literacy are of such a complex nature that they require the skills of a speech language pathologist.
- Treatment for voice disorders will need evidence of an instrumental assessment completed by an ENT or SLP to rule out a medical cause or structural deficit [9].
- Treatment for fluency disorders will need evidence that stuttering is a medical condition and is no longer developmental in nature [10].
- Treatment incorporating nonspeech oral motor exercises (NSOMEs) must be evidence based and paired with functional articulation and/or feeding/swallowing tasks [11]

Background

INDICATIONS

~~Must have written referral from primary care practitioner or other non-physician practitioner (NPP) as permitted by state guidelines.~~

~~When skilled services are also being provided by other community service agencies and/or school systems, the notes must show how the requested services are working in coordination with these agencies and not duplicating services. The extent or lack of these additional services must be indicated in the documentation.~~

~~Formal testing must be age appropriate, norm referenced, standardized, and specific to the therapy provided. Test scores should meet the following criteria to establish presence of a functional delay. Notes should document the following to establish the presence of delays or deficits:-~~

~~The following methods are generally accepted measures that may be considered to support a significant delay:~~

~~Standardized scores at or below the 10th percentile in at least one subtest area for the patient's age.³-~~

~~Standardized scores greater than or equal to 1.5 standard deviations below the mean in at least one subtest area for the patient's age.¹⁻⁸~~

~~Functional delays may be established by 25% or greater deficit in age equivalency as indicated by established general guidelines of functional assessments or specific criterion-referenced tests or profiles.¹⁻⁶-~~

~~While standardized testing is preferred, scores alone may not be used as the sole criteria for determining a patient's medical need for skilled intervention. Test information must be linked to difficulty with or inability to perform everyday tasks.-~~

~~In the absence of standardized testing or when test scores show skills within normal ranges despite functional deficits, the documentation must include detailed clinical observations and objective data to document the degree and severity of the condition, in order to support the medical need for skilled services. A caregiver interview/questionnaire can also support the request.~~

~~Any time standardized testing cannot be completed, the documentation must clearly state the reason formal testing could not be done.~~

Treatment goals must be realistic, measurable, and promote attainment of developmental milestones and functional communication abilities appropriate to the patient's age and circumstances. They should include the type, amount, duration, and frequency of therapy services.⁹ The amount, frequency, and duration of the services must be consistent with accepted standards of practice. Treatment goals must be individualized and measurable in order to identify the functional levels related to appropriate maintenance or maximum therapeutic benefit. Goals of intervention should target the functional deficits identified by the skilled therapist during the assessment and promote attainment of:-

Age-appropriate developmental milestones, functional skills appropriate to the patient's age and circumstances. Although identified as component parts of participation, underlying factors, performance skills, client factors or the environment should not be the targeted outcome of long-term goals. For sustained positive benefits from therapeutic interventions, activities can be practiced in the child's environment and reinforced by the parents or other caregivers. Practice in one's natural environment is essential for success¹⁰

The plan of care must include goals detailing type, amount, duration, and frequency of therapy services required to achieve targeted outcomes. The frequency and duration must also be commensurate with the patient's level of disability, medical and skilled therapy needs, as well as accepted standards of practice while reflecting clinical reasoning and current evidence.⁹

Frequency and duration of skilled services must also be in accordance with the following:

Intense frequencies (3x/week or more) will require additional documentation and testing supporting a medical need to achieve an identified new skill or recover function with specific, achievable goals within the requested intensive period.⁹ Details on why a higher frequency is more beneficial than a moderate or low frequency must be included. Higher frequencies may be considered when delays are classified as severe as indicated by corresponding testing guidelines used in the evaluation. More intensive frequencies may be necessary in the acute phase, however, progressive decline in frequency is expected within a reasonable time frame.

Moderate frequency (2x/week) should be consistent with moderate delays as established in the general guidelines of formal assessments used in the evaluation. This frequency may be used for ongoing care when documentation supports this frequency as being clinically effective toward achieving the functional goals in the treatment plan within a reasonable time frame.

Low frequency (1x/week or less) may be considered when testing guidelines indicate mild delays or when a higher frequency has not been clinically effective, and a similar outcome is likely with less treatment per week.

All requested frequencies must be supported by skilled treatment interventions regardless of level of severity of delay.

~~Additional factors may be considered on a case-by-case basis.~~

~~There must be evidence as to whether the services are considered reasonable, effective, and of such a complex nature that they require the technical knowledge and clinical decision-making skill of a therapist or whether they can be safely and effectively carried out by non-skilled personnel without the supervision of qualified professionals.~~

~~Treatment that requires the technical knowledge and clinical decision-making expertise to meet the skilled service needs of the individual. This includes analyzing medical/behavioral data and selecting appropriate evaluation tools/protocols to determine communication/swallowing diagnosis and prognosis.~~

~~Progress notes/updated plans of care that cover the patient's specific progress towards their goals with review by the primary care practitioner or other NPP will be required every 60-90 days or per state guidelines. Documentation should include:~~

~~The patient's current level of function, any conditions that are impacting his/her ability to benefit from skilled intervention.~~

~~Objective measures of the patient's overall functional progress relative to each treatment goal as well as a comparison to the previous progress report.~~

~~Skilled treatment techniques that are being utilized in therapy as well as the patient's response to therapy and why there may be a lack thereof. Treatment goals that follow a hierarchy of complexity to achieve the target skills for a functional goal.~~

~~Re-evaluation/annual testing (for habilitative therapy) using formal standardized assessment tools and formal assessment of progress must be performed to support progress, ongoing delays and medical necessity for continued services.~~

~~An explanation of any significant changes in the plan of care and clinical rationale for why the ongoing skills of a SLP are medically necessary.~~

~~If the patient is not progressing, then documentation of a revised treatment plan is necessary. Discontinuation of therapy will be expected when the maximum therapeutic value of a treatment plan has been achieved, no additional functional improvement is apparent or expected to occur, and the provision of services for a condition cease to be of therapeutic value.~~

~~It is expected that a specific discharge plan, with the expected treatment frequency and duration, must be included in the plan of care. The discharge plan must indicate the plan to wean services once the patient has attained their goals, if no measurable functional improvement has been demonstrated, or if the program can be carried out by caregivers or other non-skilled personnel.~~

~~It is expected that there be evidence of the development of age-appropriate home regimen to facilitate carry-over of target skills and strategies and education of patient, family, and caregiver in home practice exercises, self-monitoring as well as indication of compliance for maximum benefit of therapy.~~

~~For patients no longer showing functional improvement, a weaning process of one to two months should occur. Behaviors that interfere with the ability to progress with therapy qualify under the ASHA discharge criteria guidelines.¹¹ If the patient shows signs of regression in function, the need for skilled speech therapy can be re-evaluated at that time. Periodic episodes of care may be needed over a lifetime to address specific needs or changes in condition resulting in functional decline.~~

~~A maintenance level of therapy services may be considered when a member requires skilled therapy for ongoing periodic assessments and consultations and the member meets one of the following criteria:~~

~~Documentation shows the member and the responsible adult have a continuing need for education, or a periodic adjustment of the home program is needed to meet the member's needs.~~

~~Goals in the plan of care must be updated to reflect that care is focused on maintaining the current level of functioning and preventing regression, rather than progressing or improving function.~~

~~Clear documentation of the skilled interventions rendered and objective details of how these interventions are preventing deterioration or making the condition more tolerable must be provided. The notes must also clearly demonstrate that the specialized judgment, knowledge, and skills of a qualified therapist (as opposed to a non-skilled individual) are required for the safe and effective performance of services in a maintenance program.~~

~~For patients whose language background differs from the rendering therapist and in situations in which a clinician who has native or near-native proficiency in the target language is not available, use of an interpreter is appropriate and should be documented accordingly. If an interpreter is not present, rationale for this should be documented as well as documentation that provides evidence of a communication disorder, and a treatment plan that supports linguistically appropriate services without the use of an interpreter. Further, if a patient is substantially exposed to more than one language, the assessment must evaluate both languages and contain appropriate tests and measures to clearly denote the presence that a communication disorder is present as opposed to normal linguistic variations related to second language learning.^{12,13}~~

~~Swallowing disorders (dysphagia) and feeding disorders will need documentation of an oral, pharyngeal, and/or esophageal phase disorder, food intolerance or aversion. There must be evidence of ongoing progress and a consistent home regimen to facilitate carry-over of target feeding skills, strategies and education of patient, family, and caregiver. Therapies for picky eaters who can eat and swallow normally meeting growth and developmental milestones, eat at least one food from all major food groups (protein, grains, fruits, etc.) and more than 20 different foods is not medically necessary.~~

~~Documentation should include any applicable coordination of services with other community service agencies and/or school systems. If services are not available, then this should be indicated in the documentation.~~

~~Treatment that includes goals for reading/literacy must also have a primary diagnosis of a speech or language disorder. Documentation must support that the deficits in reading/literacy are affecting functional activities of daily living and are not the primary focus of treatment. They must show how the services for reading/literacy are of such a complex nature that they require the skills of a speech language pathologist.~~

Definitions

Habilitative Speech Therapy

Treatment provided by a state-regulated speech therapist to help a person attain, maintain, or prevent deterioration of a skill or function never learned or acquired. There must be measurable improvement and progress towards functional goals within an anticipated timeframe toward a patient's maximum potential. Treatment may also be appropriate in a child with a progressive disorder when it has the potential to prevent the loss of a functional skill or enhance the adaptation to such functional loss. The condition must be such that there is a reasonable expectation that the services will bring about a significant improvement within a reasonable time frame, regardless of whether the individual has a coexisting disorder. Ongoing treatment is not appropriate when functioning is steady and treatment no longer yields measurable functional progress.

Rehabilitative Speech Therapy

Treatment provided by a state-regulated speech therapist designed to help a person recover from an acute injury or exacerbation of a chronic condition that has resulted in a decline in functional performance. The specific impact of injury or exacerbation on the patient's ability to perform in their everyday environment must be supported by appropriate tests and measures in addition to clinical observations. Services must be provided within a reasonable time frame (frequency/duration) to restore lost function or to teach compensatory techniques if full recovery of function is not possible.

Functional Skills

They are considered necessary communication and feeding/swallowing activities of daily life. The initial plan of care documents baseline impairments as they relate to functional

communication with specific goals developed that are measurable, sustainable and time-specific. Subsequent plans of care document progress toward attainment of these goals in perspective to the patients' potential ability. Discontinuation of therapy will be expected when the maximum therapeutic value of a treatment plan has been achieved, no additional functional improvement is apparent or expected to occur, and the provision of services for a condition ceases to be of therapeutic value.

POLICY HISTORY

Date	Summary
December 2023 <u>August 2023</u>	<ul style="list-style-type: none"> Reorganized information for improved readability Added Table of Contents Updated Statement and Purpose to better match Magellan standards <u>Required test score cut-offs removed, replaced with requirement that any testing method be interpreted in accordance with its scoring method</u> <u>Distinction made between high frequency and intense frequency of treatments.</u> <u>Additional guidance on treatment for fluency disorders and nonspeech oral exercises added</u> <u>Updated References</u>
December 2022	<ul style="list-style-type: none"> Updated indications – revised criteria for standardized testing Revised language for maintenance programs Revised language for patients with a language background different than rendering therapist and for patients exposed to more than one language Clarified formal testing section and added references to support accepted measures for a significant delay Updated references
December 2021	<ul style="list-style-type: none"> Added “General Information” statement Added “resulting in adverse determinations” to EPSDT statement Reworded for clarity indication regarding bilingual patients (patients whose language background differs from rendering therapist) Added criteria stating that treatment including goals for reading/literacy must have primary diagnosis of speech or language disorder with documentation support showing how services for reading/literacy require skills of a speech language pathologist
August 2020	<ul style="list-style-type: none"> Changed guideline name to include ‘rehabilitative’: Outpatient Habilitative and <i>Rehabilitative</i> Speech Therapy Added to definition of Habilitative and Rehabilitative Therapy

	<ul style="list-style-type: none"> • Criteria for delay was revised to include clearer and more detailed specifications for functional delays, preferred scoring, and what is required in the absence of standardized testing. • Additional specifications included for linking testing to the treatment goals, inclusion of functional treatment goals, utilizing appropriate dosing of therapy and specifying skilled interventions. • Moved coordination with school program to end of guideline. • Added EPSDT language in policy statement section • Added indication of home program compliance for max benefit of therapy as part of updated POC • Added ASHA guideline for discharge qualification due to behavior • Added teletherapy to the policy statement • Formatted and adjusted language to match the PT/OT habilitative guideline where applicable
January 2020	<ul style="list-style-type: none"> • Added the <i>italicized</i> clauses as follows: For bilingual patients whose primary language differs from the rendering therapist and in situations in which a clinician who has native or near native proficiency in the target language is not available, use of an interpreter is appropriate and should be documented accordingly. If an interpreter is not present, rationale for this should be documented. Further, the assessment must contain appropriate tests and measures to clearly denote the presence that a communication disorder is present in both languages, as opposed to normal linguistic variations or a language learning problem for the non-dominant language.
July 2019	<ul style="list-style-type: none"> • Added the following definition for rehabilitative speech therapy: Rehabilitative Speech Therapy Treatments provided by a state-regulated speech therapist designed to improve, maintain, and prevent the deterioration of skills and functioning for daily living that have been lost or impaired. • Added the following to the definition of functional skills: Discontinuation of therapy will be expected when the maximum therapeutic value of a treatment plan has been

	<p>achieved, no additional functional improvement is apparent or expected to occur, and the provision of services for a condition ceases to be of therapeutic value.</p> <ul style="list-style-type: none"> • Speech therapy initial evaluation revised to require developmental delay or condition that has a standard/composite score that is ≥ 1.5 standard deviations below the mean • Clarified “picky eater” to state that for those who can eat and swallow normally meeting growth and developmental milestones, eat at least one food from all major food groups (protein, grains, fruits, etc.) and more than 20 different foods outpatient habilitative ST is not medically necessary
--	--

References

- [1] Centers for Medicare and Medicaid Services, "EPSDT - A Guide for States: Coverage in the Medicaid Benefit for Children and Adolescents," Medicaid CHIP Program, 2014.
- [2] Centers for Medicare and Medicaid Services, "Early and Periodic Screening, Diagnostic, and Treatment," June 2022. [Online]. Available: <https://www.medicaid.gov/medicaid/benefits/early-and-periodic-screening-diagnostic-and-treatment/index.html>. [Accessed August 2023].
- [3] P. H. Lipkin and M. M. Macias, "Promoting optimal development: Identifying infants and young children with developmental disorders through developmental surveillance and screening," *Pediatrics*, vol. 145, no. 1, 2020.
- [4] A. Houtrow and N. Murphy, "Prescribing physical, occupational, and speech therapy services for children with disabilities," *Pediatrics*, vol. 143, no. 4, 2019.
- [5] American Speech-Language-Hearing Association, "Admission/Discharge Criteria in Speech-Language Pathology," 2023. [Online]. [Accessed <https://www.asha.org/policy/gl2004-00046/#sec1.3> August 2023].
- [6] R. G. Voigt, "Clinical judgment and child development, revisited," *Pediatrics*, vol. 149, no. 3, 2022.
- [7] American Speech-Language-Hearing Association, "Scope of practice in speech-language pathology," 2016. [Online]. Available: <https://www.asha.org/policy/SP2016-00343/#Domains>. [Accessed August 2023].
- [8] American Speech-Language-Hearing Association, "Bilingual Service Delivery," 2023. [Online]. Available: https://www.asha.org/practice-portal/professional-issues/bilingual-service-delivery/#collapse_1. [Accessed August 2023].
- [9] American Speech-Language-Hearing Association, "Voice Disorders (Practice Portal)," [Online]. Available: www.asha.org/Practice-Portal/Clinical-Topics/Voice-Disorders/. [Accessed 28 November 2023].
- [10] B. Guitar and E. G. Conture, Eds., *The child who stutters: To the pediatrician*, Fourth ed., Memphis: Stuttering Foundation of America, 2006.
- [11] A. Alhaidary, "Treatment of speech sound disorders in children: Nonspeech oral exercises," *International Journal of Pediatrics and Adolescent Medicine*, vol. 8, pp. 1-4, 2021.

REFERENCES

1. Early and Periodic Screening, Diagnostic and Treatment (EPSDT)—A Guide for States. Coverage in the Medicaid Benefit for Children and Adolescents (2014). Centers for Medicare and Medicaid Services. December 2, 2022. https://www.medicaid.gov/sites/default/files/2019-12/epsdt_coverage_guide.pdf
2. Early and Periodic Screening, Diagnostic, and Treatment. Centers for Medicare and Medicaid Services. Updated June 29, 2022. Accessed December 2, 2022. <https://www.medicaid.gov/medicaid/benefits/early-and-periodic-screening-diagnostic-and-treatment/index.html>
3. Maine Department of Education. Severity Rating Scales/Guidelines for Speech/Language Communication Services—Language Severity Rating Scale. Updated August 1, 2020. Accessed December 2, 2022. <https://www.maine.gov/doe/sites/maine.gov.doe/files/2022-09/PROCEDURAL%20MANUAL%20Update%208-1-2020.pdf>
4. Georgetown University Center for Child and Human Development. Contemporary Practices in Early Intervention: Developmental Delay and IDEA Primer (2011). Accessed December 2, 2022. https://www.teachingei.org/disabilities/primers/Developmental_Delay.pdf
5. Maryland State Department of Education (MSDE), Division of Special Education/Early Intervention Services. Guidelines for the Use of the Developmental Delay (DD) Eligibility Category. Children Ages Birth through Seven Years. Updated March 2012. Accessed December 2, 2022.
6. Voigt RG. Clinical Judgment and Child Development, Revisited. *Pediatrics*. Mar 1 2022;149(3):doi:10.1542/peds.2021-054835
7. Mithyantha R, Kneen R, McCann E, Gladstone M. Current evidence-based recommendations on investigating children with global developmental delay. *Arch Dis Child*. Nov 2017;102(11):1071-1076. doi:10.1136/archdischild-2016-311271
8. The Speech Pathology Group (SPG). Professional Development Speech Resources—Bell Curve. Updated 2022. Accessed December 2, 2022. [SPG-Bell Curve-EnglishSpanish.pdf](https://www.spgtherapy.com/bell-curve/) (spgtherapy.com)
9. Bailes AF, Reder R, Burch C. Development of guidelines for determining frequency of therapy services in a pediatric medical setting. *Pediatr Phys Ther*. Summer 2008;20(2):194-8. doi:10.1097/PEP.0b013e3181728a7b
10. Houtrow A, Murphy N. Prescribing Physical, Occupational, and Speech Therapy Services for Children With Disabilities. *Pediatrics*. Apr 2019;143(4):doi:10.1542/peds.2019-0285
11. American Speech Language Hearing Association. Admission/discharge criteria in speech-language pathology [Guidelines]. American Speech Language Hearing Association (ASHA). Updated 2004. Accessed August 2, 2022. <https://www.asha.org/policy/gl2004-00046/>
12. American Speech Language Hearing Association. Bilingual Service Delivery. American Speech Language Hearing Association (ASHA). Accessed August 2, 2022. <https://www.asha.org/practice-portal/professional-issues/bilingual-service-delivery/>
13. Kester ES. *Difference or disorder?: understanding speech and language patterns in culturally and linguistically diverse students*. Bilingualistics Speech and Language Services; 2014:122.



ADDITIONAL RESOURCES

1. Biddle A, Watson L, Hooper C, Lohr K, Sutton S. 52 Criteria for Determining Disability in Speech Language Disorders: Summary. Agency for Healthcare Research and Quality (AHRQ). Updated January 2002. Accessed August 3, 2022. <https://www.ncbi.nlm.nih.gov/books/NBK11866/>
2. American Speech Language Hearing Association. Guidelines for Speech Language Pathologists Providing Swallowing and Feeding Services in Schools. American Speech Language Hearing Association (ASHA). Updated 2007. Accessed August 2, 2022. <https://www.psha.org/pdfs/asha-feeding-ga.pdf>
3. Kummer A. Speech pathology for the child with disability. 21st ed. McGraw-Hill; 2003:545.
4. Law J, Garrett Z, Nye C. Speech and language therapy interventions for children with primary speech and language delay or disorder. The Cochrane database of systematic reviews. 2003;2003(3):CD004110-CD004110. doi:10.1002/14651858.CD004110
5. Leung AK, Kao CP. Evaluation and management of the child with speech delay. Am Fam Physician. Jun 1999;59(11):3121-8, 3135.
6. Ong C, Phuah KY, Salazar E, How CH. Managing the 'picky eater' dilemma. Singapore Med J. 2014;55(4):184-190. doi:10.11622/smedj.2014049
7. Uher R, Rutter M. Classification of feeding and eating disorders: review of evidence and proposals for ICD-11. World Psychiatry. 2012;11(2):80-92. doi:10.1016/j.wpsyc.2012.05.005
8. Wood E, Bhalloo I, McCaig B, Feraru C, Molnar M. Towards development of guidelines for virtual administration of paediatric standardized language and literacy assessments: Considerations for clinicians and researchers. SAGE Open Med. 2021;9:20503121211050510. doi:10.1177/20503121211050510
9. McLeod S, Crowe K. Children's Consonant Acquisition in 27 Languages: A Cross-Linguistic Review. Am J Speech Lang Pathol. 2018 Nov 21;27(4):1546-1571. doi: 10.1044/2018_AJSLP-17-0100. PMID: 30177993.

Reviewed/Approved by NIA Clinical Guideline Committee

Disclaimer: National Imaging Associates, Inc. (NIA) authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Evolent Health LLC subsidiaries including, but not limited to, National Imaging Associates ("NIA"). The policies constitute only the reimbursement and coverage guidelines of NIA. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. NIA reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.

GENERAL INFORMATION

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

Disclaimer: Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates ("Magellan"). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.