

| *National Imaging Associates, Inc. | |
|--|---------------------------------------|
| Clinical guideline | Original Date: September 1997 |
| CERVICAL SPINE MRI | |
| CPT Codes: 72141, 72142, 72156, +0698T | Last Revised Date: DecemberMay 2023 |
| Guideline Number: NIA_CG_040 | Implementation Date: JulyJanuary 2024 |

GENERAL INFORMATION

- It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.
- Where a specific clinical indication is not directly addressed in this guideline, medical necessity determination will be made based on widely accepted standard of care criteria. These criteria are supported by evidence-based or peer-reviewed sources such as medical literature, societal guidelines and state/national recommendations.

INDICATIONS FOR CERVICAL SPINE MRI

⁺ If there is a combination request* for an overlapping body part, either requested at the same time or sequentially (within the past 3 months) the results of the prior study should be:

- Inconclusive or show a need for additional or follow up imaging evaluation OR
- The office notes should clearly document an indication why overlapping imaging is needed and how it will change management for the patient.

(*Unless approvable in the combination section as noted in the guidelines)

For evaluation of neurologic deficits¹⁻⁶

- With any of the following new neurological deficits documented on physical exam
 - Extremity muscular weakness (and not likely caused by plexopathy, or peripheral neuropathy)
 - Pathologic (e.g., Babinski, Lhermitte's sign, Chaddock Sign, Hoffman's and other upper motor neuron signs); OR abnormal deep tendon reflexes (and not likely caused by plexopathy, or peripheral neuropathy)
 - Absent/decreased sensory changes along a particular cervical dermatome (nerve distribution): pin prick, touch, vibration, proprioception, or temperature (and not likely caused by plexopathy, or peripheral neuropathy)

Page **1** of **19** Cervical Spine MRI

- Upper or lower extremity increase muscle tone/spasticity
- New onset bowel or bladder dysfunction (e.g., retention or incontinence)- not related to an inherent bowel or bladder process
- Gait abnormalities (see <u>Table 1</u> for more details)
- Suspected cervical cord compression with any neurological deficits as listed above

For evaluation of neck pain with any of the following⁷⁻⁹

- With new or worsening objective <u>neurologic deficits</u> (as listed above) on exam
- Failure of conservative treatment* for at<u>minimum of least</u> six (6) weeks within the last six (6) months;¹⁰

NOTE - Failure of conservative treatment is defined as one of the following:

- o Lack of meaningful improvement after a full course of treatment; OR
- Progression or worsening of symptoms during treatment; OR

<u>Documentation of a medical reason the member is unable to participate in treatment</u>
 <u>Closure of medical or therapy offices, patient inconvenience, or noncompliance without</u>
 <u>explanation does not constitute "inability to complete" treatment.</u>

- With progression or worsening of symptoms during the course of conservative treatment*
- With an abnormal electromyography (EMG) or nerve conduction study (if performed) indicating a cervical radiculopathy. (EMG is not recommended to determine the cause of axial lumbar, thoracic, or cervical spine pain.)¹¹
- Isolated back pain in pediatric population^{12, 13} conservative care not required if red flags present.

Red flags that prompt imaging include any of the following:

- Age 5 or younger OR
- o Constant pain, OR
- Pain lasting > 4 weeks, OR
- Abnormal neurologic examination, OR
- Early morning stiffness and/or gelling; OR
- Night pain that prevents or disrupts sleep; **OR**
- Radicular pain; OR
- Fever or weight loss or malaise **OR**^{14, 15}
- Postural changes (e.g., kyphosis or scoliosis) OR
- Limp (or refusal to walk in a younger child)

As part of initial pre-operative / post-operative / procedural evaluation ("CT best examination to assess for hardware complication, extent of fusion and pseudoarthrosis"^{12, 16} and MRI for cord, nerve root compression, disc pathology or post-op infection)

- For preoperative evaluation/planning
- CSF leak highly suspected and supported by patient history and/or physical exam findings (leak (known or suspected spontaneous (idiopathic) intracranial hypotension (SIH), post lumbar



puncture headache, post spinal surgery headache, orthostatic headache, rhinorrhea or otorrhea, or cerebrospinal-venous fistula))

- A follow-up study may be needed to help evaluate a patient's progress after treatment, procedure, intervention, or surgery in the last 6 months. Documentation requires a medical reason that clearly indicates why additional imaging is needed for the type and area(s) requested (routine surveillance post-op not indicated without symptoms)
- Surgical infection as evidenced by signs/symptoms, laboratory, or prior imaging findings
- New or changing neurological deficits or symptoms post-operatively^{16, 17} see <u>neurological</u> <u>deficit</u> section above
- When combo requests (see <u>above statement</u>⁺) are submitted (e.g., MRI and CT of the spine), the office notes should clearly document the need for both studies to be done simultaneously (e.g., the need for both soft tissue and bony anatomy is required)¹⁸
 - Combination requests where both cervical spine CT and MRI cervical spine are both approvable (not an all-inclusive list):
 - OPLL (Ossification of posterior longitudinal ligament)¹⁹
 - Pathologic or complex fractures
 - Malignant process of spine with both bony and soft tissue involvement
 - Unstable craniocervical junction
 - Clearly documented indication for bony and soft tissue abnormality where assessment will change management (i.e., surgical approach) for the patient

For evaluation of suspected myelopathy²⁰⁻²⁴

- Does **NOT** require conservative care
- Progressive symptoms including hand clumsiness, worsening handwriting, difficulty with grasping and holding objects, diffuse numbness in the hands, pins and needles sensation, increasing difficulty with balance and ambulation
- Any of the <u>neurological deficits</u> as noted above

For evaluation of known or suspected multiple sclerosis (MS)^{20, 25-27}

- Evidence of MS on recent baseline Brain MRI
- Suspected or known MS with new or changing symptoms consistent with cervical spinal cord disease (focal <u>neurologic deficit</u> or clinical sign, e.g., Lhermitte sign)
- Suspected or known pediatric demyelinating diseases (MS/ADEM)

Combination studies MS²⁸

• These body regions might be evaluated separately or in combination as guided by physical examination findings (e.g., localization to a particular segment of the spinal cord), patient history (e.g., symptom(s), time course, and where in the CNS the likely localization(s) is/are), and other available information, including prior imaging.



- Cervical and/or Thoracic MRI for evaluation of highly suspected multiple sclerosis (MS) when Brain MRI has indeterminate findings and/or does not fulfill the McDonald criteria for the diagnosis of MS²⁶
- Cervical and/or Thoracic MRI with suspected transverse myelitis with appropriate clinical symptoms (e.g., bilateral weakness, sensory disturbance, and autonomic dysfunction which typically evolve over hours or days)
- Brain MRI with Cervical and/or Thoracic MRI for evaluation of neuromyelitis optica spectrum disorders (recurrent or bilateral optic neuritis; recurrent transverse myelitis)²⁹
- Known MS, entire CNS axis (Brain, and/or Cervical and/or Thoracic spine) is approvable prior to the initiation or change of disease modification treatments and assess disease burden (to establish a new baseline)
- Known MS- Follow-up scans, including brain and spine imaging, if patients have known spine disease:
 - 6-12 months after starting/changing treatment
 - Every 1-2 years while on disease-modifying therapy to assess for subclinical disease activity, less frequently when stable for 2-3 years

For evaluation of trauma or acute injury^{12, 30}

- Presents with any of the following <u>neurological deficits</u> noted above
- With progression or worsening of symptoms during the course of <u>conservative treatment</u>*
- History of underlying spinal abnormalities (i.e., ankylosing spondylitis, diffuse idiopathic skeletal hyperostosis) (Both MRI and CT would be approvable)³¹⁻³³
- When the patient is clinically unevaluable or there are preliminary imaging findings (x-ray or CT) needing further evaluation
- When office notes specify the patient meets NEXUS (National Emergency X-Radiography Utilization Study) or CCR (Canadian Cervical Rules) criteria for imaging:
 - CT for initial imaging
 - MRI when suspect spinal cord or nerve root injury or when patient is obtunded, and CT is negative
 - CT or MRI for treatment planning of unstable spine

("MRI and CT provide complementary information. When indicated it is appropriate to perform both examinations")³¹

For evaluation of known or new compression fractures with worsening neck pain¹²

- With history of malignancy
 - \circ $\;$ To aid in differentiation of benign osteoporotic fractures from metastatic disease
 - A follow-up MRI in 6-8 weeks after initial MRI when initial imaging cannot decipher (indeterminate) benign osteoporotic fracture from metastatic disease (Kumar, 2016)
- With an associated new focal <u>neurologic deficit</u> as above³⁴
- Prior to a planned surgery/intervention or if the results of the MRI will change management



Page 4 of 19

For evaluation of tumor, cancer, or metastasis with any of the following:

(MRI is usually the preferred study, but CT may be needed to further characterize solitary indeterminate lesions seen on MRI)^{12, 35-37}

- Primary tumor
 - Initial staging primary spinal tumor³⁸
 - Follow-up of known primary cancer of patient undergoing active treatment within the past year or as per surveillance imaging guidance for that cancer
 - Known spinal tumor with new signs or symptoms (e.g., new or increasing nontraumatic pain, physical, laboratory, and/or imaging findings)
 - With an associated new focal <u>neurologic deficit</u> as above³⁴
- Metastatic tumor
 - With evidence of metastasis on bone scan needing further clarification OR inconclusive findings on a prior imaging exam
 - With an associated new focal neurologic deficit³⁴
 - Known malignancy with new signs or symptoms (e.g., new or increasing nontraumatic pain, radiculopathy or neck pain that occurs at night and wakes the patient from sleep with known active cancer, physical, laboratory, and/or imaging findings) in a tumor that tends to metastasize to the spine^{12, 39}

Further evaluation of indeterminate findings on prior imaging (unless follow up is otherwise specified within the guideline):

- For initial evaluation of an inconclusive finding on a prior imaging report that requires further clarification
- One follow-up exam of a prior indeterminate MR/CT finding to ensure no suspicious interval change has occurred. (No further surveillance unless specified as highly suspicious or change was found on last follow-up exam.)

Indication for combination studies for the initial pre-therapy staging of cancer, OR active monitoring for recurrence as clinically indicated, OR evaluation of suspected metastases

 ≤ 5 concurrent studies to include CT or MRI of any of the following areas as appropriate depending on the cancer: Neck, Abdomen, Pelvis, Chest, Brain, Cervical Spine, Thoracic Spine or Lumbar Spine

For evaluation of known or suspected infection (osteomyelitis)/abscess^{12,40}

• Infection

Page **5** of **19** Cervical Spine MRI

- As evidenced by signs and/or symptoms, laboratory (i.e., abnormal white blood cell count, ESR and/or CRP) or prior imaging findings⁴¹
- Follow-up imaging of infection



 With worsening symptoms/laboratory values (i.e., white blood cell count, ESR/CRP) or radiographic findings⁴²

For evaluation of known or suspected inflammatory disease or atlantoaxial instability

- In rheumatoid arthritis with neurologic signs/symptoms, or evidence of subluxation on radiographs (lateral radiograph in flexion and neutral should be the initial study)^{43, 44}
 - Patients with negative radiographs but symptoms suggestive of cervical instability or in patients with neurologic deficits MRI is indicated⁴⁵
- High-risk disorders affecting the atlantoaxial articulation, such as Down syndrome, Marfan syndrome with neurological signs/symptoms, abnormal neurological exam, or evidence of abnormal or inconclusive radiographs of the cervical spine⁴⁶
- Spondyloarthropathies, known or suspected
 - Ankylosing Spondylitis/Spondyloarthropathies with non-diagnostic or indeterminate x-ray and appropriate rheumatology workup

For evaluation of spine abnormalities related to immune system suppression, e.g., HIV, chemotherapy, leukemia, or lymphoma^{47, 48}

• As evidenced by signs/symptoms, laboratory, or prior imaging findings

Other Indications for a Cervical Spine MRI

(Note- See <u>combination requests</u>, below, for initial advanced imaging assessment and pre-operatively)

- Tethered cord or spinal dysraphism (known or suspected), based on preliminary imaging, neurological exam, and/or high-risk cutaneous stigmata⁴⁹⁻⁵¹
 - Known Arnold-Chiari syndrome (For <u>initial imaging</u> (one-time initial MRI-modality assessment) see combination below)
 - Known Chiari I malformation without syrinx or hydrocephalus, follow-up imaging after initial diagnosis with new or changing signs/symptoms or exam findings consistent with spinal cord pathology⁵²
 - o Known Chiari II (Arnold-Chiari syndrome), III, or IV malformation
- Achondroplasia (one Cervical Spine MRI to assess the craniocervical junction, as early as possible, even in asymptomatic cases)^{53, 54}
- Syrinx or syringomyelia (known or suspected)
 - With neurologic findings and/or predisposing conditions (e.g., Chiari malformation, prior trauma, neoplasm, arachnoiditis, severe spondylosis⁵⁵)
 - \circ $\,$ To further characterize a suspicious abnormality seen on prior imaging
 - Known syrinx with new/worsening symptoms
- Toe walking in a child with signs/symptoms of myelopathy localized to the Cervical Spine
- Suspected neuroinflammatory Conditions/Diseases (e.g., sarcoidosis, Behcet's)
 - \circ $\;$ After detailed neurological exam and appropriate initial work up
- Initial evaluation of trigeminal neuralgia⁵⁶ not explained on recent Brain imaging

Page 6 of 19 Cervical Spine MRI

COMBINATION OF STUDIES WITH CERVICAL SPINE MR

Brain MRI/Cervical MRI

• For evaluation of known Arnold-Chiari Malformation

Cervical and Thoracic MRI

- Initial evaluation of known or suspected syrinx or syringomyelia
 - With neurologic findings and/or predisposing conditions (e.g., Chiari malformation, prior trauma, neoplasm, arachnoiditis, severe spondylosis⁵⁵)
 - o To further characterize a suspicious abnormality seen on prior imaging
 - Known syrinx with new/worsening symptom

Any combination of Cervical and/or Thoracic and/or Lumbar MRIs

Note: These body regions might be evaluated separately or in combination as documented in the clinical notes by physical examination findings (e.g., localization to a particular segment of the spinal cord), patient history, and other available information, including prior imaging.

Exception- Indications for combination studies^{57, 58}: Are approved indications as noted below and being performed in children who will need anesthesia for the procedure

- Any combination of these studies for:
 - Survey/complete initial assessment of infant/child with congenital scoliosis or juvenile idiopathic scoliosis under the age of 10⁵⁹⁻⁶¹ (e.g., congenital scoliosis, idiopathic scoliosis, scoliosis with vertebral anomalies)
 - In the presence of neurological deficit, progressive spinal deformity, or for preoperative planning⁶²
 - Back pain with known vertebral anomalies (hemivertebrae, hypoplasia, agenesis, butterfly, segmentation defect, bars, or congenital wedging) in a child on preliminary imaging
 - Scoliosis with any of the following⁶³:
 - Progressive spinal deformity
 - Neurologic deficit (new or unexplained)
 - Early onset
 - Atypical curve (e.g., short segment, > 30' kyphosis, left thoracic curve, associated organ anomalies)
 - Pre-operative planning; OR
 - When office notes clearly document how imaging will change management
- Arnold-Chiari malformations^{64, 65}
 - o Arnold-Chiari I



- For evaluation of spinal abnormalities associated with initial diagnosis of Arnold-Chiari Malformation. (C/T/L spine due to association with tethered cord and syringomyelia), and initial imaging has not been completed^{51, 59}
- o Arnold-Chiari II-IV For initial evaluation and follow-up as appropriate
 - Usually associated with open and closed spinal dysraphism, particularly meningomyelocele
- Tethered cord, or spinal dysraphism (known or suspected) based on preliminary imaging, neurological exam, and/or high-risk cutaneous stigmata,⁴⁹⁻⁵¹ when anesthesia required for imaging⁶⁶ (e.g., meningomyelocele, lipomeningomyelocele, diastematomyelia, fatty/thickened filum terminale, and other spinal cord malformations)
- Oncological applications (e.g., primary nervous system, metastatic)
 - Drop metastasis from brain or spine (imaging also includes brain)- see Overview section
 - Suspected leptomeningeal carcinomatosis (LC)⁶⁷ -see Overview section
 - Any combination of these for spinal survey in patient with metastases
 - o Tumor evaluation and monitoring in neurocutaneous syndromes See Overview section
- CSF leak highly suspected and supported by patient history and/or physical exam findings (leak (known or suspected spontaneous (idiopathic) intracranial hypotension (SIH), post lumbar puncture headache, post spinal surgery headache, orthostatic headache, rhinorrhea or otorrhea, or cerebrospinal-venous fistula))

BACKGROUND

Magnetic resonance imaging (MRI) produces high quality multiplanar images of organs and structures within the body without radiation. It is the preferred modality for evaluating the internal structure of the spinal cord, providing assessment of conditions such as degenerative disc pathology, osteomyelitis, and discitis.

OVERVIEW

*Conservative Treatmentherapy - (Spine)

<u>Non-operative conservative treatment</u> should include a multimodality approach consisting of a<u>t least</u> <u>one (1)</u> combination of active and <u>one (1)</u> inactive component <u>targeting the affected region.</u>s.

Active Modalities

- Physical therapy
- Physician-supervised home exercise program**
- Chiropractic care

Inactive Modalities

- Medications (e.g., NSAIDs, steroids, analgesics)
- Injections (e.g., epidural injection, selective nerve root block)
- Medical Devices (e.g., TENS unit, bracing)



Inactive components, such as rest, ice, heat, modified activities, medical devices, acupuncture and/or stimulators, medications, injections (epidural, facet, bursal, and/or joint, not including trigger point), and diathermy can be utilized. Active modalities may consist of physical therapy, a physician-supervised home exercise program**, and/or osteopathic manipulative medicine (OMT) or chiropractic care when considered safe and appropriate.

**Home Exercise Program— (HEP)/ Therapy:

The following <u>two</u> elements are required to meet <u>guidelines for completion of</u> conservative therapy <u>guidelines for HEP:</u>^{68, 69}

- <u>Documentation of an Information provided on exercise prescription/plan provided by a</u> <u>physician, physical therapist, or chiropractor;</u> **AND**
- Follow-up with member with documentation provided regarding lack of improvement (failed) after completion of HEP (after the required suitable 6-week timeframe period), or inability to complete HEP due to a documented medical reason (e.g., physical reason i.e., increased pain, or inability to physically perform exercises).
- (Patient inconvenience or noncompliance without explanation does not constitute "inability to complete" HEP).

Dates and duration of failed PT, physician supervised HEP, or chiropractic treatment should be documented in the original office notes or an addendum to the notes.

Cervical myelopathy – Symptom severity varies, and a high index of suspicion is essential for making the proper diagnosis in early cases. Symptoms of pain and radiculopathy may not be present. The natural history of myelopathy is characterized by neurological deterioration. The most frequently encountered symptom is gait abnormality (86%) followed by increased muscular reflexes (79.1%), pathological reflexes (65.1%), paresthesia of upper limb (69.8%), and pain (67.4%).²⁴

| Gait | Characteristic | Work up/Imaging |
|----------------------|--|--|
| Hemiparetic | Spastic unilateral, circumduction | Brain and/or, Cervical spine imaging based on associated symptoms |
| Diplegic | Spastic bilateral, circumduction | Brain, Cervical and Thoracic Spine imaging |
| Myelopathic | Wide based, stiff, unsteady | Cervical and/or Thoracic spine MRI based on associated symptoms |
| Cerebellar Ataxic | Broad based, clumsy, staggering, lack of coordination, usually also with limb ataxia | Brain imaging - see Brain MRI Guideline |
| Apraxic | Magnetic, shuffling, difficulty initiating | Brain imaging - see Brain MRI Guideline |
| Parkinsonian | Stooped, small steps, rigid, turning en bloc, decreased arm swing | Brain Imaging - see Brain MRI Guideline |

Table 1: Gait and spine imaging⁷⁰⁻⁷⁵



| Choreiform | Irregular, jerky, involuntary movements | Medication review, consider brain imaging as per movement disorder Brain MR guidelines |
|----------------|--|--|
| Sensory ataxic | Cautious, stomping, worsening without visual input (ie + Romberg) | EMG, blood work, consider spinal (cervical or thoracic cord imaging) imaging based on EMG |
| Neurogenic | Steppage, dragging of toes | EMG initial testing; BUT if there is a foot drop, lumbar spine MRI is appropriate without EMG Pelvis MR if there is evidence of plexopathy |
| Vestibular | Insecure, veer to one side, worse when eyes closed, vertigo | Consider Brain/IAC MR- see Brain MRI Guideline |

Table 2: MRI and Cutaneous Stigmata

| Risk Stratification for Various Cutaneous Markers | | |
|--|--|---|
| <u>High Risk</u> | Intermediate Risk | Low Risk |
| Hypertrichosis Infantile hemangioma Atretic meningocele DST Subcutaneous lipoma Caudal appendage Segmental hemangiomas in association with LUMBAR[‡] syndrome | Capillary malformations (also referred to as NFS or salmon patch when pink and poorly defined or PWS when darker red and well-defined) | Coccygeal dimple Light hair Isolated café au lait spots Mongolian spots Hypo- and hypermelanotic macules or papules Deviated or forked gluteal cleft Nonmidline lesions |
| ⁺ LUMBAR, lower body hemangioma and other cutaneous defects, urogenital abnormalities, ulcerations, myelonathy, hony defects, anorectal malformations, arterial anomalies, and renal anomalies | | |
| | | |

Ossification Posterior Longitudinal Ligament (OPLL)¹⁹ – Most common in cervical spine (rare but more severe in thoracic spine)

Neck and Back Pain with Cancer History – Bone is the third most common site of metastases after the liver and the lungs, and approximately two-thirds of all osseous metastases occur in the spine. Approximately 60–70% of patients with systemic cancer will have spinal metastasis. Radiographic (x-



ray) examination should be performed in cases of back pain when a patient has a cancer history, but without known active cancer or a tumor that tends to metastasize to the spine. This can make a diagnosis in many cases. This may occasionally allow for selection of bone scan in lieu of MRI in some cases. When radiographs do not answer the clinical question, then MRI may be appropriate after a consideration of conservative care.

Most common spine metastasis involving primary metastasis originate from the following tumors in descending order: breast (21%), lung (19%), prostate (7.5%), renal (5%), gastrointestinal (4.5%), and thyroid (2.5%). While all tumors can seed to the spine, the cancers mentioned above metastasize to the spinal column early in the disease process. Spinal metastasis is more commonly found in the thoracic region, followed by the lumbar region, while the cervical region is the least likely site of metastasis.³⁹

MRI and Neurocutaneous Syndromes

- In NF-1, clinical evaluation appears to be more useful to detect complications than is screening imaging in asymptomatic patients. Imaging is indicated in evaluation of suspected tumors based on clinical evaluation and for follow-up of known intracranial and intraspinal tumors.⁷⁶
- Conversely in NF-2, routine MR imaging screening is always indicated, given the high prevalence
 of CNS tumors, especially vestibular schwannomas. In patients with NF-2, routine screening
 brain/IAC imaging is indicated annually starting from age 10, if asymptomatic, or earlier with
 clinical signs/symptoms. Most individuals with NF2 eventually develop a spinal tumor, mostly
 commonly schwannomas, but meningioma and ependymomas are also seen. Spinal imaging at
 baseline and every 2 to 3 years is also advised with more frequent imaging, if warranted, based
 on sites of tumor involvement.⁷⁷
- In patients with Tuberous Sclerosis, Brain MRI should be obtained every 1-3 years up until age 25 for surveillance for CNS abnormalities.⁷⁸
- In Von Hippel Lindau Syndrome, imaging of the brain and spinal cord for hemangioblastomas is recommended every 2 years.⁷⁹
- In Sturge Weber Syndrome, Brain MRI can rule out intracranial involvement after only age 1 and is recommended in patients <1 year old only if symptomatic.⁸⁰

Drop Metastases⁸¹ – Drop metastases are intradural extramedullary spinal metastases that arise from intracranial lesions. Common examples of intracranial neoplasms that result in drop metastases include pineal tumors, ependymomas, medulloblastomas, germinomas, primitive neuroectodermal tumors (PNET), glioblastomas multiform, anaplastic astrocytomas, oligodendrogliomas, and less commonly choroid plexus neoplasms and teratomas.

Leptomeningeal Carcinomatosis⁸² – Leptomeningeal carcinomatosis is complication of cancer in which cancerous cells spread to the membranes (meninges) that covers the brain and spinal cord. The most common solid tumors that involve the leptomeninges are breast, lung, melanoma, gastrointestinal, and primary central nervous system tumors.

Page **11** of **19** Cervical Spine MRI

POLICY HISTORY

| Date | Summary |
|-----------------|---|
| <u>Dec 2023</u> | Conservative treatment language updated in body and background |
| May 2023 | Updated references |
| | Updated background section |
| | Clarified pathological reflexes |
| | Added trigeminal neuralgia |
| | Added "Further evaluation of indeterminate or questionable findings on |
| | prior imaging": |
| | Clarified cerebellar ataxia in gait table |
| | Removed Additional Resources |
| | General Information moved to beginning of guideline with added |
| | statement on clinical indications not addressed in this guideline |
| | Added statement regarding clinical indications not addressed in the guideline. |
| March 2022 | Added |
| | Combination request for overlapping body part statement |
| | Clarified muscle weakness no related to plexopathy or peripheral |
| | neuropathy |
| | Clarified bowel and bladder dysfunction – not related to an inherent |
| | bowel or bladder problem |
| | Clarified isolated neck pain in pediatric patient |
| | Clarified combination MS for cervical and/or thoracic spine combination requests |
| | Added subsection for cervical and thoracic spine section for syrinx and syringomyelia |
| | Descriptions for tethered cord |
| | Background section of Drop Metastases |
| | Background section of Leptomeningeal Carcinomatosis |
| | Clarified toe walking in pediatric patient with myelopathy for cervical |
| | spine |
| | Removed |
| | Removed from combination section syrinx and syringomyelia and added |
| | subsection for cervical and thoracic spine section |
| | Removed pediatric back pain from the total spine combination section |
| | |

 $\ensuremath{\mathbb{C}}$ 1997-2024 National Imaging Associates, Inc., All Rights Reserved.

Page **12** of **19**

REFERENCES

1. Acharya AB, Fowler JB. Chaddock Reflex. StatPearls Publishing. Updated June 27, 2022. Accessed December 1, 2022. https://www.ncbi.nlm.nih.gov/books/NBK519555/

2. Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis and Treatment of Cervical Radiculopathy from Degenerative Disorders. North American Spine Society (NASS). Updated 2010. Accessed December 1, 2022.

https://www.spine.org/Portals/0/Assets/Downloads/ResearchClinicalCare/Guidelines/CervicalRadiculo pathy.pdf

3. Bono CM, Ghiselli G, Gilbert TJ, et al. An evidence-based clinical guideline for the diagnosis and treatment of cervical radiculopathy from degenerative disorders. *Spine J*. Jan 2011;11(1):64-72. doi:10.1016/j.spinee.2010.10.023

4. Stolper K, Haug JC, Christensen CT, Samsey KM, April MD. Prevalence of thoracic spine lesions masquerading as cauda equina syndrome: yield of a novel magnetic resonance imaging protocol. *Intern Emerg Med*. Dec 2017;12(8):1259-1264. doi:10.1007/s11739-016-1565-9

 Teoli D, Rocha Cabrero F, Ghassemzadeh S. Lhermitte Sign. StatPearls Publishing. Updated September 21, 2022. Accessed Decmber 1, 2022. https://www.ncbi.nlm.nih.gov/books/NBK493237/
 Albert TJ, Murrell SE. Surgical management of cervical radiculopathy. *J Am Acad Orthop Surg*. Nov-Dec 1999;7(6):368-76. doi:10.5435/00124635-199911000-00003

7. Allegri M, Montella S, Salici F, et al. Mechanisms of low back pain: a guide for diagnosis and therapy. *F1000Res*. 2016;5doi:10.12688/f1000research.8105.2

8. American Association of Neurological Surgeons, Congress of Neurological Surgeons. Five things physicians and patients should question: Don't obtain imaging (plain radiographs, magnetic resonance imaging, computed tomography [CT], or other advanced imaging) of the spine in patients with non-specific acute low back pain and without red flags. Choosing Wisely Initiative ABIM Foundation. Updated 2014. Accessed November 19, 2022. https://www.choosingwisely.org/clinician-

lists/american-association-neurological-surgeons-imaging-for-nonspecific-acute-low-back-pain/
Jarvik JG, Gold LS, Comstock BA, et al. Association of early imaging for back pain with clinical outcomes in older adults. *Jama*. Mar 17 2015;313(11):1143-53. doi:10.1001/jama.2015.1871
Eubanks JD. Cervical radiculopathy: nonoperative management of neck pain and radicular

symptoms. Am Fam Physician. Jan 1 2010;81(1):33-40.

11. North American Spine Society. Five things physicians and patients should question: Don't use electromyography (EMG) and nerve conduction studies (NCS) to determine the cause of axial lumbar, thoracic or cervical spine pain. Choosing Wisely Initiative ABIM Foundation. Updated 2019. Accessed December 1, 2022. https://www.choosingwisely.org/clinician-lists/nass-emg-nerve-conduction-studies-to-determine-cause-of-spine-pain/

12. American College of Radiology. ACR Appropriateness Criteria[®] Cervical Neck Pain or Cervical Radiculopathy. American College of Radiology. Updated 2018. Accessed December 1, 2022. https://acsearch.acr.org/docs/69426/Narrative/

Page 13 of 19

13. American College of Radiology. ACR Appropriateness Criteria[®]Back Pain–Child. American College of Radiology (ACR). Updated 2016. Accessed December 1, 2022.

https://acsearch.acr.org/docs/3099011/Narrative/

14. Bernstein RM, Cozen H. Evaluation of back pain in children and adolescents. *Am Fam Physician*. Dec 1 2007;76(11):1669-76.

15. Feldman DS, Straight JJ, Badra MI, Mohaideen A, Madan SS. Evaluation of an algorithmic approach to pediatric back pain. *J Pediatr Orthop*. May-Jun 2006;26(3):353-7.

doi:10.1097/01.bpo.0000214928.25809.f9

16. Rao D, Scuderi G, Scuderi C, Grewal R, Sandhu SJ. The Use of Imaging in Management of Patients with Low Back Pain. *J Clin Imaging Sci.* 2018;8:30. doi:10.4103/jcis.JCIS_16_18

17. Corona-Cedillo R, Saavedra-Navarrete MT, Espinoza-Garcia JJ, Mendoza-Aguilar AN, Ternovoy SK, Roldan-Valadez E. Imaging Assessment of the Postoperative Spine: An Updated Pictorial Review of Selected Complications. *Biomed Res Int*. 2021;2021:9940001. doi:10.1155/2021/9940001

18. Fisher BM, Cowles S, Matulich JR, Evanson BG, Vega D, Dissanaike S. Is magnetic resonance imaging in addition to a computed tomographic scan necessary to identify clinically significant cervical spine injuries in obtunded blunt trauma patients? *Am J Surg*. Dec 2013;206(6):987-93; discussion 993-4. doi:10.1016/j.amjsurg.2013.08.021

19. Choi BW, Song KJ, Chang H. Ossification of the posterior longitudinal ligament: a review of literature. *Asian Spine J*. Dec 2011;5(4):267-76. doi:10.4184/asj.2011.5.4.267

20. American College of Radiology. ACR Appropriateness Criteria® Myelopathy. American College of Radiology (ACR). Updated 2020. Accessed January 29, 2023.

https://acsearch.acr.org/docs/69484/Narrative/

21. Behrbalk E, Salame K, Regev GJ, Keynan O, Boszczyk B, Lidar Z. Delayed diagnosis of cervical spondylotic myelopathy by primary care physicians. *Neurosurg Focus*. Jul 2013;35(1):E1. doi:10.3171/2013.3.Focus1374

22. Davies BM, Mowforth OD, Smith EK, Kotter MR. Degenerative cervical myelopathy. *Bmj*. Feb 22 2018;360:k186. doi:10.1136/bmj.k186

23. Sarbu N, Lolli V, Smirniotopoulos JG. Magnetic resonance imaging in myelopathy: a pictorial review. *Clin Imaging*. Sep-Oct 2019;57:56-68. doi:10.1016/j.clinimag.2019.05.002

24. de Oliveira Vilaça C, Orsini M, Leite MA, et al. Cervical Spondylotic Myelopathy: What the Neurologist Should Know. *Neurol Int*. Nov 2 2016;8(4):6330. doi:10.4081/ni.2016.6330

25. Consortium of Multiple Sclerosis Centers. 2018 MRI Protocol and Clinical Guidelines for MS. Consortium of Multiple Sclerosis Centers (CMSC). Updated May 22, 2018. Accessed January 23, 2023. https://www.mscare.org/page/MRI_protocol

26. Filippi M, Rocca MA, Ciccarelli O, et al. MRI criteria for the diagnosis of multiple sclerosis: MAGNIMS consensus guidelines. *Lancet Neurol*. Mar 2016;15(3):292-303. doi:10.1016/s1474-4422(15)00393-2

27. Kaunzner UW, Gauthier SA. MRI in the assessment and monitoring of multiple sclerosis: an update on best practice. *Ther Adv Neurol Disord*. Jun 2017;10(6):247-261. doi:10.1177/1756285617708911

28. Barakat N, Gorman MP, Benson L, Becerra L, Borsook D. Pain and spinal cord imaging measures in children with demyelinating disease. *Neuroimage Clin*. 2015;9:338-47. doi:10.1016/j.nicl.2015.08.019



29. Wingerchuk DM, Banwell B, Bennett JL, et al. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology*. Jul 14 2015;85(2):177-89.

doi:10.1212/wnl.000000000001729

30. American College of Radiology. ACR Appropriateness Criteria[®] Suspected Spine Trauma American College of Radiology. Updated 2018. Accessed December 1, 2022.

https://acsearch.acr.org/docs/69359/Narrative/

31. American College of Radiology. ACR Appropriateness Criteria[®] Inflammatory Back Pain: Known or Suspected Axial Spondyloarthritis. American College of Radiology (ACR). Updated 2021. Accessed December 1, 2022. https://acsearch.acr.org/docs/3094107/Narrative/

32. Koivikko MP, Koskinen SK. MRI of cervical spine injuries complicating ankylosing spondylitis. *Skeletal Radiol*. Sep 2008;37(9):813-9. doi:10.1007/s00256-008-0484-x

33. Taljanovic MS, Hunter TB, Wisneski RJ, et al. Imaging characteristics of diffuse idiopathic skeletal hyperostosis with an emphasis on acute spinal fractures: review. *AJR Am J Roentgenol*. Sep 2009;193(3 Suppl):S10-9, Quiz S20-4. doi:10.2214/ajr.07.7102

34. Alexandru D, So W. Evaluation and management of vertebral compression fractures. *Perm J*. Fall 2012;16(4):46-51. doi:10.7812/tpp/12-037

35. Kim YS, Han IH, Lee IS, Lee JS, Choi BK. Imaging findings of solitary spinal bony lesions and the differential diagnosis of benign and malignant lesions. *J Korean Neurosurg Soc*. 2012;52(2):126-132. doi:10.3340/jkns.2012.52.2.126

36. Roberts CC, Daffner RH, Weissman BN, et al. ACR appropriateness criteria on metastatic bone disease. *J Am Coll Radiol*. Jun 2010;7(6):400-9. doi:10.1016/j.jacr.2010.02.015

37. American College of Radiology. ACR Appropriateness Criteria[®] Neck Mass/Adenopathy. American College of Radiology. Updated 2018. Accessed January 23, 2023.

https://acsearch.acr.org/docs/69504/Narrative/

38. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Central Nervous System Cancers Version 2.2022. National Comprehensive Cancer Network (NCCN). Updated September 29, 2022. Accessed January 23, 2023. https://www.nccn.org/professionals/physician_gls/pdf/cns.pdf

39. Ziu E, Viswanathan VK, Mesfin FB. Spinal Metastasis. StatPearls Publishing. Updated August 22, 2022. Accessed December 1, 2022. https://www.ncbi.nlm.nih.gov/books/NBK441950/

40. Kowalski TJ, Berbari EF, Huddleston PM, Steckelberg JM, Osmon DR. Do follow-up imaging examinations provide useful prognostic information in patients with spine infection? *Clin Infect Dis*. Jul 15 2006;43(2):172-9. doi:10.1086/505118

41. Bond A, Manian FA. Spinal Epidural Abscess: A Review with Special Emphasis on Earlier Diagnosis. *Biomed Res Int*. 2016;2016:1614328. doi:10.1155/2016/1614328

42. Berbari EF, Kanj SS, Kowalski TJ, et al. 2015 Infectious Diseases Society of America (IDSA) Clinical Practice Guidelines for the Diagnosis and Treatment of Native Vertebral Osteomyelitis in Adults. *Clin Infect Dis*. Sep 15 2015;61(6):e26-46. doi:10.1093/cid/civ482

43. Colebatch AN, Edwards CJ, Østergaard M, et al. EULAR recommendations for the use of imaging of the joints in the clinical management of rheumatoid arthritis. *Ann Rheum Dis*. Jun 2013;72(6):804-14. doi:10.1136/annrheumdis-2012-203158

44. Mańczak M, Gasik R. Cervical spine instability in the course of rheumatoid arthritis - imaging methods. *Reumatologia*. 2017;55(4):201-207. doi:10.5114/reum.2017.69782



45. Gillick JL, Wainwright J, Das K. Rheumatoid Arthritis and the Cervical Spine: A Review on the Role of Surgery. *Int J Rheumatol*. 2015;2015:252456. doi:10.1155/2015/252456

46. Henderson FC, Sr., Austin C, Benzel E, et al. Neurological and spinal manifestations of the Ehlers-Danlos syndromes. *Am J Med Genet C Semin Med Genet*. Mar 2017;175(1):195-211. doi:10.1002/ajmg.c.31549

47. Nagashima H, Yamane K, Nishi T, Nanjo Y, Teshima R. Recent trends in spinal infections: retrospective analysis of patients treated during the past 50 years. *Int Orthop*. Mar 2010;34(3):395-9. doi:10.1007/s00264-009-0741-1

48. American College of Radiology. ACR Appropriateness Criteria[®] Suspected Spine Infection. American College of Radiology (ACR). Updated 2021. Accessed December 1, 2022.

https://acsearch.acr.org/docs/3148734/Narrative/

49. Zalatimo O. Tethered Spinal Cord Syndrome. American Association of Neurological Surgeons (AANS). Accessed December 1, 2022. https://www.aans.org/Patients/Neurosurgical-Conditions-and-Treatments/Tethered-Spinal-Cord-Syndrome

50. Düz B, Gocmen S, Secer HI, Basal S, Gönül E. Tethered cord syndrome in adulthood. *J Spinal Cord Med*. 2008;31(3):272-8. doi:10.1080/10790268.2008.11760722

51. Milhorat TH, Bolognese PA, Nishikawa M, et al. Association of Chiari malformation type I and tethered cord syndrome: preliminary results of sectioning filum terminale. *Surg Neurol*. Jul 2009;72(1):20-35. doi:10.1016/j.surneu.2009.03.008

52. Whitson WJ, Lane JR, Bauer DF, Durham SR. A prospective natural history study of nonoperatively managed Chiari I malformation: does follow-up MRI surveillance alter surgical decision making? *J Neurosurg Pediatr*. Aug 2015;16(2):159-66. doi:10.3171/2014.12.Peds14301

53. Legare JM. Achondroplasia. University of Washington, Seattle. Updated January 6, 2022. Accessed December 1, 2022. https://www.ncbi.nlm.nih.gov/books/NBK1152/

54. White KK, Bompadre V, Goldberg MJ, et al. Best practices in the evaluation and treatment of foramen magnum stenosis in achondroplasia during infancy. *Am J Med Genet A*. Jan 2016;170a(1):42-51. doi:10.1002/ajmg.a.37394

55. Timpone VM, Patel SH. MRI of a syrinx: is contrast material always necessary? *AJR Am J Roentgenol*. May 2015;204(5):1082-5. doi:10.2214/ajr.14.13310

56. Shim YW, Paeng SH, Lee KS, Kim ST, Lee WH. Trigeminal Neuralgia Resulting from Delayed Cervical Cord Compression after Acute Traumatic Fracture of Odontoid Process. *Korean J Neurotrauma*. Apr 2019;15(1):38-42. doi:10.13004/kjnt.2019.15.e10

57. American College of Radiology. ACR Appropriateness Criteria[®] Headache. American College of Radiology. Updated 2022. Accessed January 23, 2023. https://acsearch.acr.org/docs/69482/Narrative/ 58. American College of Radiology. ACR Appropriateness Criteria[®] Headache-Child. American College of Radiology. Updated 2017. Accessed December 1, 2022.

https://acsearch.acr.org/docs/69439/Narrative/

59. Strahle J, Smith BW, Martinez M, et al. The association between Chiari malformation Type I, spinal syrinx, and scoliosis. *J Neurosurg Pediatr*. Jun 2015;15(6):607-11. doi:10.3171/2014.11.Peds14135 60. Juvenile Scoliosis. Scoliosis Research Society (SRS). Accessed December 1, 2022.

https://www.srs.org/professionals/online-education-and-resources/conditions-and-treatments/juvenile-scoliosis



61. American College of Radiology. ACR Appropriateness Criteria[®] Scoliosis-Child. American College of Radiology. Updated 2018. Accessed December 1, 2022.

https://acsearch.acr.org/docs/3101564/Narrative/

62. Trenga AP, Singla A, Feger MA, Abel MF. Patterns of congenital bony spinal deformity and associated neural anomalies on X-ray and magnetic resonance imaging. *J Child Orthop*. Aug 2016;10(4):343-52. doi:10.1007/s11832-016-0752-6

63. Ozturk C, Karadereler S, Ornek I, Enercan M, Ganiyusufoglu K, Hamzaoglu A. The role of routine magnetic resonance imaging in the preoperative evaluation of adolescent idiopathic scoliosis. *Int Orthop*. Apr 2010;34(4):543-6. doi:10.1007/s00264-009-0817-y

64. Strahle J, Muraszko KM, Kapurch J, Bapuraj JR, Garton HJ, Maher CO. Chiari malformation Type I and syrinx in children undergoing magnetic resonance imaging. *J Neurosurg Pediatr*. Aug 2011;8(2):205-13. doi:10.3171/2011.5.Peds1121

65. Radic JAE, Cochrane DD. Choosing Wisely Canada: Pediatric Neurosurgery Recommendations. *Paediatr Child Health*. Sep 2018;23(6):383-387. doi:10.1093/pch/pxy012

66. Hertzler DA, 2nd, DePowell JJ, Stevenson CB, Mangano FT. Tethered cord syndrome: a review of the literature from embryology to adult presentation. *Neurosurg Focus*. Jul 2010;29(1):E1. doi:10.3171/2010.3.Focus1079

67. Shah LM, Salzman KL. Imaging of spinal metastatic disease. *Int J Surg Oncol*. 2011;2011:769753. doi:10.1155/2011/769753

68. Last AR, Hulbert K. Chronic low back pain: evaluation and management. *Am Fam Physician*. Jun 15 2009;79(12):1067-74.

69. American College of Radiology. ACR Appropriateness Criteria[®] Low Back Pain. American College of Radiology (ACR). Updated 2021. Accessed January 29, 2023.

https://acsearch.acr.org/docs/69483/Narrative/

70. Chhetri SK, Gow D, Shaunak S, Varma A. Clinical assessment of the sensory ataxias; diagnostic algorithm with illustrative cases. *Pract Neurol*. Aug 2014;14(4):242-51. doi:10.1136/practneurol-2013-000764

71. Foster H, Drummond P, Jandial S, Clinch J, Wood M, Driscoll S. Evaluation of gait disorders in children. BMJ Best Practice. Updated February 23, 2021. Accessed January 23, 2023. https://bestpractice.bmj.com/topics/en-us/709

72. Standford Medicine. Gait Abnormalities. Stanford University. Accessed January 23, 2023. https://stanfordmedicine25.stanford.edu/the25/gait.html

73. Haynes KB, Wimberly RL, VanPelt JM, Jo CH, Riccio AI, Delgado MR. Toe Walking: A Neurological Perspective After Referral From Pediatric Orthopaedic Surgeons. *J Pediatr Orthop*. Mar 2018;38(3):152-156. doi:10.1097/bpo.00000000001115

74. Marshall FJ. Approach to the elderly patient with gait disturbance. *Neurol Clin Pract*. Jun 2012;2(2):103-111. doi:10.1212/CPJ.0b013e31825a7823

75. Pirker W, Katzenschlager R. Gait disorders in adults and the elderly : A clinical guide. *Wien Klin Wochenschr*. Feb 2017;129(3-4):81-95. doi:10.1007/s00508-016-1096-4

76. Borofsky S, Levy LM. Neurofibromatosis: types 1 and 2. *AJNR Am J Neuroradiol*. Dec 2013;34(12):2250-1. doi:10.3174/ajnr.A3534

77. Evans DGR, Salvador H, Chang VY, et al. Cancer and Central Nervous System Tumor Surveillance in Pediatric Neurofibromatosis 2 and Related Disorders. *Clin Cancer Res*. Jun 15 2017;23(12):e54-e61. doi:10.1158/1078-0432.Ccr-17-0590

78. Krueger DA, Northrup H. Tuberous sclerosis complex surveillance and management:

recommendations of the 2012 International Tuberous Sclerosis Complex Consensus Conference. *Pediatr Neurol*. Oct 2013;49(4):255-65. doi:10.1016/j.pediatrneurol.2013.08.002

79. Varshney N, Kebede AA, Owusu-Dapaah H, Lather J, Kaushik M, Bhullar JS. A Review of Von Hippel-Lindau Syndrome. J Kidney Cancer VHL. 2017;4(3):20-29. doi:10.15586/jkcvhl.2017.88

80. Comi AM. Presentation, diagnosis, pathophysiology, and treatment of the neurological features of Sturge-Weber syndrome. *Neurologist*. Jul 2011;17(4):179-84. doi:10.1097/NRL0b013e318220c5b6

81. Ahmed A. MRI features of disseminated 'drop metastases'. *S Afr Med J.* Jul 2008;98(7):522-3.

82. Batool A, Kasi A. Leptomeningeal Carcinomatosis. StatPearls Publishing

Copyright © 2022, StatPearls Publishing LLC. Updated April 5, 2022. Accessed December 1, 2022. https://www.ncbi.nlm.nih.gov/books/NBK499862/





Reviewed / Approved by NIA Clinical Guideline Committee

Disclaimer: National Imaging Associates, Inc. (NIA) authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Evolent Health LLC subsidiaries including, but not limited to, National Imaging Associates ("NIA"). The policies constitute only the reimbursement and coverage guidelines of NIA. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. NIA reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of regulations.

