

<u>Subject:</u>	<u>Minimally Invasive Ablative Procedures for Epilepsy</u>	<u>Publish Date:</u>	<u>12/12/2018</u>
<u>Guideline #:</u>	<u>CG-SURG-91</u>	<u>Last Review Date:</u>	<u>11/08/2018</u>
<u>Status:</u>	<u>New</u>		

**Description**

**This document addresses minimally invasive ablative procedures used in the treatment of medically refractory epilepsy in individuals with symptomatic localized epilepsy. Minimally invasive procedures have been proposed as a means to minimize or eliminate major craniotomy and bone flap incisions, decrease pain and down-time, preserve tissue and decrease neurocognitive adverse effects. These procedures utilize laser, radiofrequency, or cryotherapy techniques, in combination with stereotactic magnetic resonance imaging (MRI) guidance, for targeted ablation of the epileptogenic foci.**

**Note: Please see the following related documents for other treatments of epilepsy:**

- BEH.00002 Transcranial Magnetic Stimulation
- CG-ANC-03 Acupuncture
- CG-MED-05 Ketogenic Diet for Treatment of Intractable Seizures
- MED.00057 MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications
- SURG.00007 Vagus Nerve Stimulation
- SURG.00026 Deep Brain, Cortical, and Cerebellar Stimulation

**Clinical Indications****Metrically Necessary:**

**This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.**

**Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.**

**No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.**

**The treatment of medically refractory epilepsy using stereotactic laser techniques (MRI-guided laser interstitial thermal ablation [MRIgLITT]), including stereotactic laser amygdalohippocampotomy (SLAH), is considered medically necessary when the following criteria are met:**

- **Documented disabling seizures despite the use of two or more tolerated antiepileptic drug regimens; and**
- **Documented presence of two or fewer well delineated epileptogenic foci accessible by laser.**

**The use of stereotactic radiofrequency thermocoagulation (RF-TC) in the treatment of hypothalamic hamartomas is considered medically necessary.**

## **Not Medically Necessary:**

**The treatment of medically refractory epilepsy using stereotactic laser techniques or stereotactic radiofrequency thermocoagulation is considered not medically necessary when the criteria above have not been met.**

**Other minimally invasive procedures to treat medically intractable epilepsy are considered not medically necessary, including but not limited to stereotactic radiofrequency amygdalohippocampectomy, SEEG-guided radiofrequency thermocoagulation or stereotactic cryosurgery.**

## **Coding**

***The following codes for treatments and procedures applicable to this guideline are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.***

**CPT  
64999**

**Unlisted procedure, nervous system [when specified as minimally invasive surgery such as MRI-guided laser interstitial thermal ablation, radiofrequency thermal coagulation]**

**This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.**

**Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.**

**No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.**

## ICD-10 Diagnosis

D33.0

G40.001-G40.919

Benign neoplasm of brain, supratentorial [specified as hypothalamic hamartoma]

Epilepsy and recurrent seizures

## Discussion/General Information

Approximately 1.2% of the U.S population, or 3.4 million individuals have active epilepsy (CDC, 2018).

Active epilepsy is defined as being physician diagnosed and having had one or more seizures in the past year or taking medication to control it, or both (CDC, 2018). Mesial temporal lobe epilepsy is a common type of epilepsy, representing approximately one-quarter of all cases; one-third of these individuals are considered medication refractory (Gross, 2018). Other types of epilepsy have varying degrees of success with pharmacotherapy. The failure of 2 anti-seizure medications is approximately 97% reliable in identifying drug resistance (Kwan, 2010). The International League against Epilepsy defines drug resistant epilepsy as:

failure of adequate trials of two tolerated and appropriately chosen and used AED schedules (whether as monotherapies or in combination) to achieve sustained seizure freedom (Kwan, 2010).

While pharmacotherapy is the first line of treatment to control seizures, it does not affect complete seizure control in one-third of the individuals (LaRiviere, 2016). For this population, open procedures have been the gold standard in achieve lasting seizure control. These procedures, such as anterior temporal lobectomy (ATL) or selective amygdalohippocampotomy (SAH) involve localizing and excising the epileptogenic zones (Shukla, 2017). These open procedures typically report a 60-80% long term seizure-free rate with 95% of individuals reporting an improvement (Chang, 2015; Kang, 2016). However, concerns about the invasiveness of the procedure and the effect on neuropsychological function has limited the number of individuals utilizing this option. Approximately 2% of those individuals considered candidates undergo open surgery every year (Kang, 2016). Minimally invasive therapies have been explored as alternatives to open surgery. These therapies are proposed to be viable options for those with epileptogenic foci located near deep, eloquently brain structures, multiple comorbidities or those who are poor surgical candidates (Shukla, 2017).

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

**Stereotactic LITT, typically MRI- guided, uses low-voltage laser energy delivered via optical fiber through a burr hole to destroy the targeted tissue while minimizing injury to the surrounding tissue. Laser energy is converted to thermal energy, which, when applied to tissue leads to coagulative necrosis. Use with MRI guidance allows for monitoring of both the device tip and thermal damage in real time (Gross, 2016; Shukla, 2017). However, LITT is not without drawbacks. Ablative therapies do not allow for tissue to be obtained for pathology, and Kang (2016) noted a technical limitation “The curvature of the hippocampus and the presence of potential heat sinks (i.e., blood and cerebral spinal fluid) may prevent adequate ablation of the epileptogenic network in some individuals.**

### Laser interstitial therapy

**Currently, there are two Food and Drug Administration (FDA) cleared MRgLITT systems, Visualase® thermal therapy (Medtronic, MN) and NeuroBlate® Systems (Monteris Medical, Minneapolis, MN). The Visualase device, approved in 2007, has been used to treat epilepsy since 2012 (Kang, 2016). The NeuroBlate device, approved in 2009, is not commonly used to treat epilepsy, but is typically used to treat brain tumors. The use of NeuroBlate to treat epilepsy is currently undergoing a FDA approved trial.**

**Gross and colleagues (2018) evaluated outcomes of 58 individuals with mesial temporal lobe epilepsy (MTLE) with or without mesial temporal sclerosis who had undergone stereotactic laser amygdalohippocampotomy (SLAH). Analysis was limited to those with documented outcomes of 1 year or longer. For those for whom the initial SLAH procedure was ineffective, a repeat SLAH procedure was offered when the postoperative MRI scan showed a remnant region thought to be responsible for the ongoing procedures. Following SLAH, 53.4% (31/58) were considered free from disabling seizures, and 22 of the individuals achieved Engel 1A response at 12 months or longer, including 3 individuals who had undergone a repeat SLAH. An additional 9 individuals achieved Engel 1B-D responses. During the same period 22 individuals underwent open temporal lobe resections, with 50% (11/22) of these individuals considered seizure free at one year. Complications of SLAH include visual field defects in five individuals, with one being persistent and symptomatic. Other complications were transient and treated with resolution of symptoms. The authors reported that 8.2% of the individuals experienced decline in verbal memory, compared to rates of 30 to 60% decline reported for open resection. The authors note:**

**This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.**

**Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.**

**No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.**

**Thus, on balance, although SLAH may yield a marginal decrease in the chance of freedom from disabling seizures, and possibly in the rate of being completely seizure-free, as compared to open resection, the cognitive benefits and safety profile of SLAH make it an attractive alternative, especially when considering surgery on the dominant hemisphere.**

**The outcomes of LITT to treat 43 consecutive individuals with MTLE who underwent unilateral therapy was reported by Donos and associates (2018). In addition to seizure free rates, cognitive function was assessed prior to and following the procedure, although 8 individuals did not return for follow-up testing. At 6 months following their procedure, 79.5% of the individuals had achieved an Engel class I outcome. Additional 5.1% achieved Engel class II and 15.4% were classified as having an Engel III outcome. In later follow-up (mean interval of 20.3 months), 67.4% of the individuals achieved an Engel class I outcome and an additional 16.3% of the individuals each achieved Engel class II and III. There were declines in some aspects of memory following the surgery, but changes observed at the group level were not small and did not reach the level of significance. The authors noted that the follow-up testing was performed at 6 months, and as functional recovery continued, a further improvement in cognitive performance was likely to occur. The results of the study were similar to those of previous studies, the authors reported slightly higher rates of Engel I outcomes.**

**Kang and associates (2016) described the outcomes of 20 individuals with medically intractable MTLE who underwent MRgLITT between 2011 and 2014. Assessed outcomes included mesial temporal lobe ablated volumes, verbal memory, and surgical outcomes at 6 months, 1 year, 2 years and most recent visit. At 6 months, 8/15 individuals (53%, 95% CI 30.1-75.2%) were free of consciousness impairing seizures. At 1 and 2 years, 4/11 individuals (36.4%, 95% CI 14.9-64.8%) and 3/5 individuals (60%, 95% CI 22.9-88.4%) were free of consciousness impairing seizures respectively. None of the participants reported a statistically reliable decline in logical memory performance. Reported efficacy rates are modestly lower than those reported after anterior temporal lobectomy (ATL). However, verbal learning and memory performance was better preserved in the MRgLITT group compared to the ATL group. In addition, the hospital stay postoperative pain and recovery period was minimal, and individuals were immediately able to return to work. The authors note that this finding might be significant for those with dominant mesial temporal foci, as**

**This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.**

**Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.**

**No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.**

**individuals who undergo ATL in the dominant hemisphere are at higher risk for declines in language function and verbal memory.**

**In 2015, Waseem and colleagues reported on the outcomes of 7 individuals over the age of 50 who underwent MRgLITT for medically refractory MTLE. The outcomes were compared to 7 individuals who had undergone selective anterior mesial temporal lobe (AMTL) resection. All 7 individuals in the AMTL group were seizure free at 12 months. In the MRgLITT group, 4 individuals were classified as seizure free, 2 individuals reported rare seizures (almost seizure free) and 2 individuals had not reached the 12 month follow-up. Complications at 30 days included one case of aseptic meningitis that resolved with therapy (ATML), 2 cases of postoperative partial visual field deficits (MRgLITT) and 1 case postoperative seizure requiring readmission (MRgLITT). There was no significant differences in neuropsychological outcomes between the two groups. The authors note that MRgLITT might be an acceptable alternative for those individuals who cannot or is not willing to undergo an open surgical intervention.**

**The current body of evidence is limited to case review and series, and prospective and retrospective studies. While randomized, controlled studies comparing minimally invasive procedures to the gold standard of ATL provides a high level of evidence, this type of study is not feasible for LITT in the current setting. The preponderance of evidence suggests that the efficacy of LITT is comparable to open procedures and has improved safety and tolerance outcomes (Curry, 2012; Drane, 2015; Hawasli, 2013; Lewis, 2015; McCracken, 2016; Patel, 2016; Perry, 2017; Tao, 2018; Tandonb, 2018; Wilfong, 2013; Willie, 2015; Youngerman, 2018).**

### **Radiofrequency therapies**

**Stereoelectroencephalography-guided radiofrequency thermocoagulation (SEEG-guided RF-TC), also called thermo-SEEG, involves the use of implanted SEEG electrodes to both identify the location of seizure-onset zones and to perform multiple stereotactic lesioning of the identified areas. SEEG-guided RF-TC is proposed as alternative in a select population when resective surgery is not feasible, such as when the ictal onset zone size is limited or when the zone is located in a highly functional area (Moles, 2018). Moles (2018) notes the following advantages to SEEG-guided RF-TC:**

**This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.**

**Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.**

**No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.**

- (i) there is no additional bleeding risk when compared to a conventional stereotactic procedure as the same electrode is used for both SEEG and RF-TC (all bleedings reported in the literature were related to removal of SEEG electrodes, and these were not those used for RF-TC [9]),
- (ii) it allows a very accurate targeting of the seizure-onset zone, previously delineated by intracranial recordings,
- (iii) multiple lesions can be performed, instead of the single or double lesions usually performed in a conventional stereotactic lesioning procedure, and
- (iv) a functional mapping, through direct electrical stimulation on SEEG electrodes, is done prior to any lesion being made, thus allowing to anticipate the possible adverse effects in detail.

In a retrospective review comparing the outcomes of SEEG-guided RF-TC (n=21) and ATL (n=49) in temporal lobe epilepsy, Moles and associates (2018) reported significantly worse outcomes for the SEEG-guided RF-TC group. At 12 months, none of the SEEG-guided RF-TC group were seizure free while 75.5% of the ATL group were seizure free. Of the 21 individuals in the SEEG-guided RF-TC group, 19 individuals subsequently underwent ATL, 1 individual was waiting to undergo ATL, and the last individual was seizure-free for 2 months before being classified as a responder (at least 50% reduction in seizure frequency).

Bourdillon and associates (2017) evaluated the outcomes 162 individuals who underwent SEEG-guided RF-TC to treat drug-resistant focal epilepsy. The primary outcome was identified as seizure frequency at 2 months and one year following SEEG-guided RF-TC. In addition, the rate of responders, defined as individuals with at least a 50% decrease in seizure frequency compared the 3 months immediately preceding the procedure, was the secondary outcome. At 2 months, 25% (n=41) of the individuals were seizure free and 67% (n=108) were considered responders. At 12 months, 7% (n=11) of the individuals were seizure free and 48% (n=78) of the individuals were considered responders. A total of 50 individuals underwent a second procedure within the year following the procedure, 2 of those individuals had a second SEEG-guided RF-TC. While the results of this study might suggest that SEEG-guided RF-TC has a favorable risk-benefit ratio compared to conventional surgery and other palliative treatments, the significance of these finding was limited by the prospective nature of the study and the fact this single center study was performed over a 10 year period and included modifications in practice over this time.

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

**Radiofrequency ablation has been studied as a means of eliminating epileptogenic zones without performing a craniotomy. While early results were disappointing, later studies have reported improved outcomes over the earlier data (LaRiviere, 2016). However, these later studies are small, lower quality studies (Guénot, 2004; Guénot, 2011; Krámská, 2017; Malikova, 2014; Wu, 2014). Additional study with long-term follow-up is needed to address question regarding efficacy, safety and durability of outcomes compared to current standard treatments.**

## **Hypothalamic hamartomas (HH)**

**HH is a rare congenital disorder, affecting an estimated 1 in every 200,000 individuals. While HH can present in a number of ways, intractable gelastic seizures are a common symptom (Kameyama, 2016). In addition to the gelastic seizures, approximately half of the affected individuals also have epileptic encephalopathy, which is exhibited as cognitive impairment and behavioral disorders. Gelastic seizures typically do not respond to pharmacotherapy, treatment options include microscopic surgery, endoscopic disconnection, stereotactic radiofrequency thermocoagulation (SRT), laser interstitial thermal therapy, and Gamma Knife surgery (Tandon, 2018a). The location of HH lesions, near of the midline of the skull base, make the lesions difficult to reach with microsurgical approaches and too invasive to provide total resection (Kameyama, 2016). The success rate of microscopic surgery has been reported as 5-60% with higher rates of morbidity and mortality (Tandon, 2018). In addition to treating gelastic seizures, both laser and radiofrequency procedures have been associated with an improvement in epileptic encephalopathy symptoms (Sonoda, 2017).**

**Sonoda and associates (2017) evaluated cognitive function outcomes in 88 individuals who underwent SRT as a treatment of drug-resistant gelastic seizures (GS). A total of 87 individuals underwent post-operative follow-up for an average of 3.3 years. At the final hospital visit, 85.2% (75/87) had achieved GS remission. For those individuals in gelastic seizure remission there was a significant post-operative improvement in full-scale intelligence quotient (FSIQ) compared to preoperative scores (preoperative  $72.8 \pm 25.1$  versus postoperative at the last visit:  $81.2 \pm 26.7$ ;  $p<0.001$ ; paired t-test). Individuals who did not achieve GS remission did not achieve a significant improvement in FSIQ scores (preoperative:  $68.8 \pm 23.3$  versus postoperative at the last visit  $71.1 \pm 24.1$ ;  $p=0.36$ ; paired t-test).**

---

**This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.**

**Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.**

**No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.**

**Kameyama and colleagues (2016) reported on the outcomes of 100 consecutive pediatric (n=70) and adult (n=30) individuals with intractable gelastic seizures who underwent MRI-guided stereotactic radiofrequency thermocoagulation (SRT). In addition to gelastic seizures, 90 of the individuals had other types of seizures including complex partial, secondarily generalized tonic-clonic, tonic, atonic, myoclonic and epileptic. All individuals had SRT as the sole treatment for hypothalamic hamartomas, although 32 individuals underwent multiple (up to 4) SRT procedures for residual disease. Individuals were followed-up between 1 and 17 years. A reported 86% of the individuals achieved freedom from gelastic seizures and 78.9% of the individuals with concurrent non-gelastic seizures also achieved seizure freedom.**

**Several other studies, including case series, retrospective reviews, reported on the outcomes of either stereotactic laser ablation or stereotactic radiofrequency thermocoagulation used to treat HH lesions have been published. Both laser and SRT trials have reported outcomes superior to microscopic surgery with limited morbidity (Curry, 2018; Kameyama, 2016; Southwell, 2018; Tandon, 2018; Wei, 2018).**

### ***Other minimally invasive therapies***

**To date, there is a paucity of studies regarding other minimally invasive treatments of epilepsy. Chkhenkeli and colleagues (2013) reported on the results of 21 individuals who underwent stereotactic cryosurgery to treat intractable bitemporal epilepsy. However, there have been no additional published regarding the use of cryosurgery for the treatment of epilepsy.**

### **Definitions**

**Anterior temporal lobectomy (ATL): Surgical resection of the mesial-basal temporal lobe structures. A limitation of ATL is that access to these structures requires resection of the temporal neocortex which affects cognitive and neuropsychological abilities.**

**Medically intractable epilepsy: The failure of at least two separate drug regimens to control seizure activity; also known as drug resistant epilepsy.**

**This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.**

**Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.**

**No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.**

**Selective amygdalohippocampectomy (SAH): Surgical resection of mesial-basal temporal lobe structures which does not involve neocortical resection.**

## References

### Peer Reviewed Publications:

1. Al Sufiani F, Ang LC. Neuropathology of temporal lobe epilepsy. *Epilepsy Res Treat*. 2012;624519.
2. Bourdillon P, Isnard J, Catenoix H, et al. Stereo electroencephalography-guided radiofrequency thermocoagulation (SEEG-guided RF-TC) in drug-resistant focal epilepsy: results from a 10-year experience. *Epilepsia*. 2017; 58(1):85-93.
3. Buckley R, Estronza-Ojeda S, Ojemann JG. Laser Ablation in Pediatric Epilepsy. *Neurosurg Clin N Am*. 2016; 27(1):69-78.
4. Chang EF, Englot DJ, Vadera S. Minimally invasive surgical approaches for temporal lobe epilepsy. *Epilepsy Behav*. 2015; 47:24-33.
5. Chkhenkeli SA, Šramka M, Rakviashvili TN, et al. Bitemporal intractable epilepsy: could it be surgically treatable? *Stereotact Funct Neurosurg*. 2013; 91(2):104-112.
6. Curry DJ, Gowda A, McNichols RJ, Wilfong AA. MR-guided stereotactic laser ablation of epileptogenic foci in children. *Epilepsy Behav*. 2012; 24(4):408-414.
7. Drane DL, Loring DW, Voets NL, et al. Better object recognition and naming outcome with MRI-guided stereotactic laser amygdalohippocampotomy for temporal lobe epilepsy. *Epilepsia*. 2015; 56(1):101-113.
8. Engel J Jr. What can we do for people with drug-resistant epilepsy? The 2016 Wartenberg Lecture. *Neurology*. 2016; 87(23):2483-2489.
9. Grewal SS, Zimmerman RS, Worrell G, et al. Laser ablation for mesial temporal epilepsy: a multi-site, single institutional series. *J Neurosurg*. 2018;1-8.
10. Gross RE, Stern MA, Willie JT, et al. Stereotactic laser amygdalohippocampotomy for mesial temporal lobe epilepsy. *Ann Neurol*. 2018; 83(3):575-587.
11. Guénöt M, Isnard J, Catenoix H, Mauguière F, Sindou M. SEEG-guided RF-thermocoagulation of epileptic foci: a therapeutic alternative for drug-resistant non-operable partial epilepsies. *Adv Tech Stand Neurosurg*. 2011; 36:61-78.

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

12. Guénöt M, Isnard J, Ryvlin P, et al. **SEEG-guided RF thermocoagulation of epileptic foci: feasibility, safety, and preliminary results.** *Epilepsia.* 2004; 45(11):1368-1374.
13. Hawasli AH, Bagade S, Shimony JS, et al. **Magnetic resonance imaging-guided focused laser interstitial thermal therapy for intracranial lesions: single-institution series.** *Neurosurgery.* 2013; 73(6):1007-1017.
14. Kang JY, Wu C, Tracy J, et al. **Laser interstitial thermal therapy for medically intractable mesial temporal lobe epilepsy.** *Epilepsia.* 2016; 57(2):325-334.
15. Lagman C, Chung LK, Pelargos PE, et al. **Laser neurosurgery: a systematic analysis of magnetic resonance-guided laser interstitial thermal therapies.** *J Clin Neurosci.* 2017; 36:20-26.
16. LaRiviere MJ, Gross RE. **Stereotactic laser ablation for medically intractable epilepsy: the next generation of minimally invasive epilepsy surgery.** *Front Surg.* 2016; 3:64.
17. Lewis EC, Weil AG, Duchowny M, et al. **MR-guided laser interstitial thermal therapy for pediatric drug-resistant lesional epilepsy.** *Epilepsia.* 2015; 56(10):1590-1598.
18. Malikova H, Kramská L, Vojtech Z, et al. **Different surgical approaches for mesial temporal epilepsy: resection extent, seizure, and neuropsychological outcomes.** *Stereotact Funct Neurosurg.* 2014; 92(6):372-380.
19. Moles A, Guénöt M, Rheims S, et al. **SEEG-guided radiofrequency coagulation (SEEG-guided RF-TC) versus anterior temporal lobectomy (ATL) in temporal lobe epilepsy.** *J Neurol.* 2018; 265(9):1998-2004.
20. Patel P, Patel NV, Danish SF. **Intracranial MR-guided laser-induced thermal therapy: single-center experience with the Visualase thermal therapy system.** *J Neurosurg.* 2016; 125(4):853-860.
21. Perry MS, Donahue DJ, Malik SI, et al. **Magnetic resonance imaging-guided laser interstitial thermal therapy as treatment for intractable insular epilepsy in children.** *J Neurosurg Pediatr.* 2017; 20(6):575-582.
22. Shukla ND, Ho AL, Pendharkar AV, et al. **Laser interstitial thermal therapy for the treatment of epilepsy: evidence to date.** *Neuropsychiatr Dis Treat.* 2017; 13:2469-2475.
23. Spencer D, Burchiel K. **Selective amygdalohippocampectomy.** *Epilepsy Res Treat.* 2012; 2012:382095.
24. Tandon V, Chandra PS, Doddamani RS, et al. **Stereotactic radiofrequency thermocoagulation of hypothalamic hamartoma using robotic guidance (ROSA) coregistered with O-arm guidance- preliminary technical note.** *World Neurosurg.* 2018a; 112:267-274.
25. Tandon V, Lang M, Chandra PS, et al. **Is edema a matter of concern after laser ablation of epileptogenic focus?** *World Neurosurg.* 2018b; 113:366-372.e3.

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

26. Tao JX, Wu S, Lacy M, et al. Stereotactic EEG-guided laser interstitial thermal therapy for mesial temporal lobe epilepsy. J Neurol Neurosurg Psychiatry. 2018; 89(5):542-548.
27. Waseem H, Osborn KE, Schoenberg MR, et al. Laser ablation therapy: an alternative treatment for medically resistant mesial temporal lobe epilepsy after age 50. Epilepsy Behav. 2015; 51:152-157.
28. Waseem H, Vivas AC, Vale FL. MRI-guided laser interstitial thermal therapy for treatment of medically refractory non-lesional mesial temporal lobe epilepsy: outcomes, complications, and current limitations: a review. J Clin Neurosci. 2017; 38:1-7.
29. Wei PH, An Y, Fan XT, et al. Stereoelectroencephalography-guided radiofrequency thermocoagulation for hypothalamic hamartomas: preliminary evidence. World Neurosurg. 2018; 114:e1073-e1078.
30. Wilfong AA, Curry DJ. Hypothalamic hamartomas: optimal approach to clinical evaluation and diagnosis. Epilepsia. 2013; 54 Suppl 9:109-114.
31. Wu Z, Zhao Q, Tian Z, et al. Efficacy and safety of a new robot-assisted stereotactic system for radiofrequency thermocoagulation in patients with temporal lobe epilepsy. Exp Ther Med. 2014; 7(6):1728-1732.
32. Youngerman BE, Oh JY, Anbarasan D, et al; Columbia Comprehensive Epilepsy Center Co-Authors. Laser ablation is effective for temporal lobe epilepsy with and without mesial temporal sclerosis if hippocampal seizure onsets are localized by stereoelectroencephalography. Epilepsia. 2018; 59(3):595-606.

**Government Agency, Medical Society, and Other Authoritative Publications:**

1. Centers for Disease Control and Prevention (CDC). Epilepsy. Updated June 7, 2018. Available at: <https://www.cdc.gov/epilepsy/data/index.html>. Accessed on November 8, 2018.
2. Kwan P, Arzimanoglou A, Berg AT, et al. Definition of drug resistant epilepsy: consensus proposal by the ad hoc Task Force of the ILAE Commission on Therapeutic Strategies. Epilepsia. 2010; 51(6):1069-1077.
3. Laser Interstitial Thermal Therapy for Treating Laser Interstitial Thermal Therapy for Treating Intracranial Lesions and Epilepsy. A Health Technology Assessment and Policy Analysis. January 27, 2016. Available at: <https://obrieniph.ucalgary.ca/system/files/litt-jan-27th-2016-references-fixed.pdf>. Accessed on November 8, 2018.
4. U.S. Food & Drug Administration.

---

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

5. **NeuroBlate® System (Monteris Medical; Plymouth, MN). 510(K) Number: K173305. November 17, 2017. Available at: [https://www.accessdata.fda.gov/cdrh\\_docs/pdf17/K173305.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf17/K173305.pdf). Accessed on November 8, 2018.**
6. **Visualase® Thermal Therapy System (Medtronic; Minneapolis, MN). 510(K) Number: K053087. March 1, 2006. Available at: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/pmn.cfm?ID=K053087>. Accessed on November 8, 2018.**

## W e b s i t e s f o r A d d i t i o n a l I n f o r m a t i o n

1. **National Institute of Health. National Institute of Neurological Disorders and Stroke. Epilepsy Information Page. Last updated June 18, 2018. Available at: <https://www.ninds.nih.gov/disorders/all-disorders/epilepsy-information-page>. Accessed on November 8, 2018.**
2. **National Organization for Rare Disorders (NORD). Hypothalamic Hamartoma. Last published 2017. Available at: <https://rarediseases.org/rare-diseases/hypothalamic-hamartoma/>. Accessed on November 8, 2018.**
3. **The Epilepsy Foundation. About Epilepsy: The Basics. Available at: <https://www.epilepsy.com/learn/about-epilepsy-basics>. Accessed on November 8, 2018.**

## I n d e x

**The use of specific product names is illustrative only. It is not intended to be a recommendation of one product over another, and is not intended to represent a complete listing of all products available.**

## H i s t o r y

<u>Status</u>	<u>Date</u>	<u>Action</u>
<u>New</u>	<u>11/08/2018</u>	<u>Medical Policy &amp; Technology Assessment Committee (MPTAC) review.</u> <u>Initial document development.</u>

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

DRAFT

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to implement a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.