

Louisiana Medicaid
Acne Agents, Topical

Formatted: Font: 12 pt

The *Louisiana Uniform Prescription Drug Prior Authorization Form* should be utilized to request clinical authorization for acne agents, topical (preferred and non-preferred). Criteria for approval of tazarotene for psoriasis is found on the next page.

For Acne

Approval Criteria

- The recipient is less than 21 years of age on the date of the request; **AND**
- For trifarotene, the recipient is at least 9 years of age on the date of the request; AND
- The recipient has a diagnosis of acne; AND
- The severity is stated on the request and is either Grade 3 moderately severe nodulocystic acne (numerous papules and pustules; the occasional inflamed nodule; the back and chest may also be affected) or Grade 4 severe nodulocystic acne (numerous large, painful pustules and nodules; inflammation); **AND**
- For tazarotene used for acne, the product requested is either 0.1% cream or 0.1% gel; **AND**
- If the request is for a non-preferred agent - **ONE** of the following is required:
 - The recipient has had a *treatment failure* with at least one preferred product; **OR**
 - The recipient has had an *intolerable side effect* to at least one preferred product; **OR**
 - The recipient has a *documented contraindication(s)* to all of the preferred products that are appropriate to use for the condition being treated; **OR**
 - There is *no preferred product that is appropriate* to use for the condition being treated; **AND**
- By submitting the authorization request, the prescriber attests to the following:
 - The prescribing information for the requested medication has been thoroughly reviewed, including any Black Box Warning, Risk Evaluation and Mitigation Strategy (REMS), contraindications, minimum age requirements, recommended dosing, and prior treatment requirements; **AND**
 - All laboratory testing and clinical monitoring recommended in the prescribing information have been completed as of the date of the request and will be repeated as recommended; **AND**
 - The recipient has no concomitant drug therapies or disease states that would limit the use of the requested medication and will not be receiving the requested medication in combination with any medication that is contraindicated or not recommended per FDA labeling.

Reauthorization criteria for a diagnosis of acne

- The recipient continues to meet all initial criteria with improved disease severity; **AND**
- The request states the current acne severity, which is an improvement from baseline.

Formatted: Font: Bold

Formatted: Font: 12 pt

Duration of initial and reauthorization approval for acne: 12 months (or up to the recipient's 21st birthday, whichever is less)

Formatted: Font: 12 pt

For Tazarotene Cream or Gel for Psoriasis

Criteria for Both Initial Approval and Reauthorization

- The recipient has a diagnosis of psoriasis; AND
- If the request is for a non-preferred agent - **ONE** of the following is required:
 - The recipient has had a *treatment failure* with at least one preferred topical antipsoriatic product (see Dermatology – Antipsoriatics, Topical on PDL); **OR**
 - The recipient has had an *intolerable side effect* to at least one preferred topical antipsoriatic product (see Dermatology – Antipsoriatics, Topical on PDL); **OR**
 - The recipient has a *documented contraindication(s)* to all of the preferred topical antipsoriatic products that are appropriate to use for the condition being treated (see Dermatology – Antipsoriatics, Topical on PDL); **OR**
 - There is *no preferred topical antipsoriatic product that is appropriate* to use for the condition being treated (see Dermatology – Antipsoriatics, Topical on PDL); **AND**
- By submitting the authorization request, the prescriber attests to the following:
 - The prescribing information for the requested medication has been thoroughly reviewed, including any Black Box Warning, Risk Evaluation and Mitigation Strategy (REMS), contraindications, minimum age requirements, recommended dosing, and prior treatment requirements; **AND**
 - All laboratory testing and clinical monitoring recommended in the prescribing information have been completed as of the date of the request and will be repeated as recommended; **AND**
 - The recipient has no concomitant drug therapies or disease states that would limit the use of the requested medication and will not be receiving the requested medication in combination with any medication that is contraindicated or not recommended per FDA labeling.

Duration of initial and reauthorization approval for tazarotene cream or gel for psoriasis: 12 months

Formatted: Font: 12 pt

References

Aklief (trifarotene) [package insert]. Fort Worth, TX: GALDERMA LABORATORIES, L.P.; <https://www.galderma.com/us/sites/g/files/jcdfhc341/files/2019-10/10-2-2019%20Revised%20PI%20NDA%20211527.pdf>

Formatted: Font: 12 pt

Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; Retrieved from <https://www.clinicalkey.com/pharmacology/>

Formatted: No underline, Font color: Auto

DiPiro JT, Talbert RL, Yee GC, Matzke GR, Wells BG, Posey L. eds. *Pharmacotherapy: A Pathophysiologic Approach*, 10e New York, NY: McGraw-Hill; Retrieved from <https://accesspharmacy.mhmedical.com/book.aspx?bookid=1861>

Tazorac gel (tazarotene) [package insert]. Madison, NJ: Allergan USA, Inc.; 2018. Retrieved from https://www.allergan.com/assets/pdf/tazorac_gel_pi

Tazorac cream (tazarotene) [package insert]. Irvine, CA: Allergan; 2017. Retrieved from https://www.allergan.com/assets/pdf/tazorac_cream_pi

Formatted: No underline, Font color: Auto

Revision	Date
Removed POS information, formatting changes, updated references	July 2020
Added Akleif®, formatting changes	<u>TBD</u>

Formatted: No underline, Font color: Auto