

## Leqvio (inclisiran) (for Louisiana Only)

Policy Number: [CSLA2022D00101A](#)

Effective Date: TBD

[Instructions for Use](#)

Table of Contents	Page
<a href="#">Application</a>	1
<a href="#">Coverage Rationale</a>	2
<a href="#">Applicable Codes</a>	2
<a href="#">Background</a>	2
<a href="#">Clinical Evidence</a>	3
<a href="#">U.S. Food and Drug Administration</a>	43
<a href="#">References</a>	43
<a href="#">Policy History/Revision Information</a>	43
<a href="#">Instructions for Use</a>	43

### Application

This Medical Benefit Drug Policy only applies to the state of Louisiana.

### Coverage Rationale

**Leqvio (inclisiran) is proven and medically necessary for the treatment of primary hypercholesterolemia (including heterozygous familial and non-familial) patients who meet all of the following criteria:<sup>1,2</sup>**

- **For initial therapy, all of the following:**
  - Diagnosis of primary hypercholesterolemia (including heterozygous familial and non-familial); and
  - Prescribed by, a lipid specialist (e.g., cardiologist, endocrinologist, lipid specialist/lipidologist); and
  - **One of the following:**
    - Despite adherence to PCSK9 therapy (defined by at least 12 consecutive weeks of use), one of the following:
      - **Both of the following:**
        - Patient has clinical ASCVD; and
        - Patient failed to achieve LDL-C goal of <70 mg/dL
      - or
      - **Both of the following:**
        - Patient has severe primary hypercholesterolemia (pre-treatment LDL-C ≥190 mg/dL); and
        - Patient failed to achieve LDL-C goal of <100 mg/dL
    - or
    - Patient has a history of intolerance or contraindication to PCSK9 therapy;
  - and**
    - Patient will continue other traditional low-density lipoprotein-cholesterol (LDL-C) lowering therapies (e.g., maximally tolerated statins, ezetimibe) in combination with Leqvio; and
    - Leqvio will not be used in combination with PCSK9 therapy; and
    - Leqvio dosing is in accordance with the United States Food and Drug Administration approved labeling; and
    - Initial authorization will be for no more than 12 months
  - **For continuation of therapy, all of the following:**
    - Documentation of a positive clinical response to therapy from pre-treatment baseline (e.g., achieved LDL-C goal of <100 mg/dL or achieved a 50% reduction in LDL-C levels); and

- Patient continues treatment with other traditional low-density lipoprotein-cholesterol (LDL-C) lowering therapies (e.g., statin, ezetimibe) in combination with Leqvio; **and**
- Leqvio will not be used in combination with PCSK9 therapy; **and**
- Leqvio dosing is in accordance with the United States Food and Drug Administration approved labeling; **and**
- Reauthorization will be for no more than 12 months.

## Definitions

For the purposes of this policy, **major ASCVD events** are defined as:

- Recent acute coronary syndrome (within the past 12 months)
- History of myocardial infarction (other than recent acute coronary syndrome event listed above)
- History of ischemic stroke
- Symptomatic peripheral arterial disease (history of claudication with ankle brachial index <0.85, or previous revascularization or amputation)

**High risk conditions** are defined as:

- Age ≥65 years
- Heterozygous familial hypercholesterolemia
- History of prior coronary artery bypass surgery or PCI outside of the major ASCVD event(s)
- Diabetes Mellitus
- Hypertension
- Chronic kidney disease (eGFR 15-59 mL/min/1.73 m<sup>2</sup>)
- Current smoking
- Persistently elevated LDL-C (LDL-C ≥100 mg/dL (≥2.6 mmol/L)) despite maximally tolerated statin therapy and ezetimibe
- History of congestive heart failure

**Very High Risk** is defined as a history of multiple major ASCVD events or 1 major ASCVD event and multiple high-risk conditions.

## Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

HCPCS Code	Description
C9399	Unclassified drugs or biologicals
J3490	Unclassified drugs
J3590	Unclassified biologics

Diagnosis Code	Description
E78.01	Familial hypercholesterolemia
Z83.42	Family history of FH

## Background

Atherosclerosis is an accumulation of lipids (mostly low density lipoprotein cholesterol [LDL-C]) in the inner lining of the arteries over time. An atherosclerotic cardiovascular event (such as heart attack or stroke) can be caused by an unexpected rupture of the atherosclerotic plaque. Proprotein convertase subtilisin/kexin type 9 (PCSK9), which is synthesized primarily in hepatocytes, enters circulation, and binds to hepatic LDL receptors, targeting the LDL receptors for degradation. In turn, this process reduces the capacity of the liver to bind and remove LDL-C, resulting in increased LDL-C levels. The binding of PCSK9 by monoclonal antibodies has been shown to reduce LDL-C levels by more than 50%.<sup>3,4,5,6</sup>

Inclisiran is a cholesterol-lowering double-stranded small interfering ribonucleic acid (siRNA), conjugated on the sense strand with triantennary N-acetylgalactosamine (GalNAc), to facilitate uptake by hepatocytes. Utilizing the RNA interference mechanism, inclisiran directs catalytic breakdown of mRNA in hepatocytes for PCSK9. This increases LDL-C receptor recycling and expression, therefore increasing LDL-C uptake and reducing LDL-C levels in circulation.<sup>1</sup>

## Clinical Evidence

Inclisiran is indicated for the treatment in adults with primary hypercholesterolemia (including heterozygous familial and non-familial) or mixed dyslipidemia, as an adjunct to diet in combination with a statin or statin with other lipid-lowering therapies in patients unable to reach LDL-C goals with the maximum tolerated dose of a statin, or alone or in combination with other lipid lowering therapies in patients who are statin-intolerant, or for whom a statin is contraindicated.<sup>1</sup>

ORION-9 (NCT03397121) was a phase 3, randomized, double-blind, placebo controlled trial, that evaluated the use of inclisiran in adult patients with heterozygous familial hypercholesterolemia (HeFH) who have been treated with a maximally tolerated dose of statin therapy. The study randomly assigned in a 1:1 ratio, 242 patients to receive inclisiran and 240 to receive placebo. 25% of patients had preexisting coronary artery disease and 10% had diabetes. The mean baseline LDL-C level was 153.1 mg/dL ( $\pm$ 54 mg/dL). 90% of patients were receiving statins, including 75% who were on a high intensity statin. More than 50% were also receiving ezetimibe. The primary end points were the percent change from baseline in the LDL-C level at day 510 and time adjusted percent change from baseline in the LDL-C level between day 90 and day 540. 91.7% of patients in the inclisiran group completed the trial activities through day 540. Secondary endpoints included mean absolute change from baseline in LDL-C at day 510, time-adjusted absolute reduction from baseline between day 90 and day 540, and changes in levels of PCSK9, total cholesterol, apolipoprotein B, and non-high-density lipoprotein (HDL) cholesterol. Prespecified exploratory end points included the proportion of patients who met lipid targets for their level of cardiovascular risk and treatment response according to genotype of familial hypercholesterolemia. Study results showed at day 510, the percent change in LDL-C level was a reduction of 39.7% (95% CI -43.7 to -35.7) in the inclisiran group and an increase of 8.2% (95% CI, 4.3 to 12.2) in the placebo group; the between-group difference was -47.9 percentage points (95% CI, -53.5 to -42.3;  $P < 0.001$ ). The time-averaged percent change in the LDL cholesterol level between day 90 and day 540 was a reduction of 38.1% (95% CI, -41.1 to -35.1) in the inclisiran group and an increase of 6.2% (95% CI, 3.3 to 9.2) in the placebo group, for a between-group difference of -44.3 percentage points (95% CI, -48.5 to -40.1;  $P < 0.001$ ). Secondary endpoint analysis showed the mean absolute change from baseline in the LDL-C level at day 510 had a between-group difference of -68.9 mg/dL (95% CI, -77.1 to -60.7;  $P < 0.001$ ). Additionally, the time-averaged observed difference in LDL cholesterol levels between day 90 and day 540 showed a between-group difference of -62.6 mg/dL ( $P < 0.001$ ). At day 510, a reduction from baseline in the mean LDL cholesterol level of 50% or more was reported in 38% of patients in the inclisiran group (compared to 0.8% in the placebo group;  $P < 0.001$ ). 65.3% of patients achieved an LDL-C level of less than 100 mg/dL. The authors concluded that among adults with HeFH, those who received inclisiran had significantly lower levels of LDL-C, than those who received placebo.<sup>6</sup>

Two randomized, double-blind, placebo-controlled, parallel-group phase 3 trials, ORION-10 (NCT03399370) (n= 1561) and ORION-11 (NCT03400800) (n= 1617), were conducted to assess the efficacy, safety, and adverse-event profile of inclisiran over a period of 19 months in patients at high risk for cardiovascular disease in whom LDL-C levels remained elevated, despite use of a maximally tolerated statin therapy with or without additional lipid-lowering therapy. Randomization was stratified according to background use of statins, where patients were assigned 1:1 to receive either inclisiran or placebo on days 1, 90, 270, and 450. The primary endpoints in each trial were placebo-corrected percent change in LDL-C level from baseline to day 510 and time-adjusted percent change in LDL-C level from baseline after day 90 and up to day 540. Secondary endpoints included mean absolute change from baseline in LDL-C at day 510, time-adjusted absolute reduction from baseline between day 90 and day 540, and changes in levels of PCSK9, total cholesterol, apolipoprotein B, and non-high-density lipoprotein (HDL) cholesterol. The mean LDL-C level at baseline was 104.7 $\pm$ 38.3 mg/dL (ORION-10) and 105.5 $\pm$ 39.1 mg/dL (ORION-11). Additionally 68% of patients were receiving high-intensity statins. The primary endpoint analysis showed at day 510, inclisiran reduced LDL-C by 52.3% (95% CI, 48.8 to 55.7) in the ORION-10 trial and by 49.9% (95% CI, 46.6 to 53.1) in the ORION-11 trial, with corresponding time-adjusted reductions of 53.8% (95% CI, 51.3 to 56.2) and 49.2% (95% CI, 46.8 to 51.6) ( $P < 0.001$  for all comparisons vs. placebo). Authors concluded that reductions in LDL-C levels of approximately 50% were obtained with inclisiran, when administered every 6 months.

## Professional Societies

The American College of Cardiology/American Heart Association Task Force published their clinical practice guidelines for the management of blood cholesterol in 2018. In regards to those with severe hypercholesterolemia (LDL-C  $\geq$ 190 mg/dL), the guideline recommends:<sup>5</sup>

In patients 20 to 75 years of age with an LDL-C level of 190 mg/dL or higher ( $\geq 4.9$  mmol/L) maximally tolerated statin therapy is recommended (Level I; B-R)

- In patients 20 to 75 years of age with an LDL-C level of 190 mg/dL or higher ( $\geq 4.9$  mmol/L) who achieve less than a 50% reduction in LDL-C while receiving maximally tolerated statin therapy and/or have an LDL-C level of 100 mg/dL or higher ( $\geq 2.6$  mmol/L) ezetimibe therapy is reasonable (Level IIa; B-R)
- In patients 20 to 75 years of age with a baseline LDL-C level 190 mg/dL or higher ( $\geq 4.9$  mmol/L), who achieve less than a 50% reduction in LDL-C levels and have fasting triglycerides 300 mg/dL or lower ( $\leq 3.4$  mmol/L) while taking maximally tolerated statin and ezetimibe therapy, the addition of a bile acid sequestrant may be considered (Level IIb; B-R)
- In patients 30 to 75 years of age with heterozygous FH and with an LDL-C level of 100 mg/dL or higher ( $\geq 2.6$  mmol/L) while taking maximally tolerated statin and ezetimibe therapy, the addition of a PCSK9 inhibitor may be considered (Level IIb; B-R)
- In patients 40 to 75 years of age with a baseline LDL-C level of 220 mg/dL or higher ( $\geq 5.7$  mmol/L) and who achieve an on-treatment LDL-C level of 130 mg/dL or higher ( $\geq 3.4$  mmol/L) while receiving maximally tolerated statin and ezetimibe therapy, the addition of a PCSK9 inhibitor may be considered (Level IIb; C-LD)

## U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Leqvio is indicated for the treatment in adults with primary hypercholesterolemia (including heterozygous familial and non-familial) or mixed dyslipidemia, as an adjunct to diet in combination with a statin or statin with other lipid-lowering therapies in patients unable to reach LDL-C goals with the maximum tolerated dose of a statin, or alone or in combination with other lipid lowering therapies in patients who are statin-intolerant, or for whom a statin is contraindicated.<sup>1</sup>

## References

1. Leqvio [package insert]. East Hanover, NJ: Novartis; December 2021.
2. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APHA/ASPC/NLA/PCNA Guideline on the Management of Blood Cholesterol: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines.
3. Banerjee Y, Shah K, Al-Rasadi K. Effect of a monoclonal antibody to PCSK9 on LDL cholesterol. N Engl J Med. 2012;366(25):2425-2426.
4. Ray KK, Landmesser U, Leiter LA, et al. Inclisiran in Patients at High Cardiovascular Risk with Elevated LDL Cholesterol. N Engl J Med. 2017;376(15):1430-1440.
5. Ray KK, Wright RS, Kallend D, et al. Two Phase 3 Trials of Inclisiran in Patients with Elevated LDL Cholesterol. N Engl J Med. 2020;382(16):1507-1519.
6. Raal FJ, Kallend D, Ray KK, et al. Inclisiran for the Treatment of Heterozygous Familial Hypercholesterolemia. N Engl J Med. 2020;382(16):1520-1530.

## Policy History/Revision Information

<u>Date</u>	<u>Summary of Changes</u>
<u>TBD</u>	<u>New policy</u>

## Instructions for Use

This Medical Benefit Drug Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plan may differ from the standard plan. In the event of a conflict, the member specific benefit plan document governs. Before using this policy, please check the member specific benefit plan document and any applicable federal or state mandates. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Benefit Drug Policy is provided for informational purposes. It does not constitute medical advice.

This Medical Benefit Drug Policy may also be applied to Medicare Advantage plans in certain instances. In the absence of a Medicare National Coverage Determination (NCD), Local Coverage Determination (LCD), or other Medicare coverage guidance, CMS allows a Medicare Advantage Organization (MAO) to create its own coverage determinations, using objective evidence-based rationale relying on authoritative evidence (Medicare IOM Pub. No. 100-16, Ch. 4, §90.5).

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. UnitedHealthcare Medical Benefit Drug Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

DRAFT