Medical Drug Clinical Criteria

Subject: Istodax (romidepsin)

Document #: CC-0100 **Publish Date:** 09/23/2024/10/01/2025

Status: Revised **Last Review Date:** 08/16/202408/15/2025

Table of Contents

<u>Overview</u> <u>Coding</u> <u>References</u>

<u>Clinical criteria</u> <u>Document history</u>

Overview

This document addresses the use of romidepsin, an intravenously administered histone deacetylase (HDAC) inhibitor. HDAC inhibitors are useful as antineoplastic agents as they cause accumulation of acetylated histones, inducing cell cycle arrest and/or apoptosis of some transformed cells. It is used to treat a certain subset of non-hodgkin lymphoma (NHL) known as T-Cell lymphomas. T-cell lymphomas account for approximately 15% of all non-hodgkin lymphoma in the United States.

NHLs are a broad and diverse group of malignancies affecting both B- and T-lymphocytes. T-Cell Lymphomas can broadly be classified as cutaneous or non-cutaneous. Romidepsin is currently FDA approved for cutaneous T-cell lymphoma in patients that have received at least one prior therapy. The National Comprehensive Cancer Network® (NCCN) provides additional recommendations with a category 2A level of evidence for the use of romidepsin for Mycosis Fungoides/Sezary Syndrome, anaplastic large cell lymphoma (ALCL) (including breast implant-associated), extranodal NK/T-Cell lymphomas, hepatosplenic T-Cell Lymphoma, and peripheral T-cell lymphomas (PTCL).

Definitions and Measures

Mycosis fungoides/ Sézary Syndrome (MF/SS): Cutaneous T-cell Lymphomas (CTCLs) are a group of NHLs of mature T-cells that primarily present in the skin, and at times progress to involve lymph nodes, blood, and visceral organs. MF is the most common subtype with primary cutaneous involvement and SS is an erythrodermic, leukemic variant of CTCL that is characterized by significant blood involvement and lymphadenopathy.

Refractory Disease: Illness or disease that does not respond to treatment.

Relapse or recurrence: After a period of improvement, during which time a disease (for example, cancer) could not be detected, the return of signs and symptoms of illness or disease. For cancer, it may come back to the same place as the original (primary) tumor or to another place in the body.

Clinical Criteria

When a drug is being reviewed for coverage under a member's medical benefit plan or is otherwise subject to clinical review (including prior authorization), the following criteria will be used to determine whether the drug meets any applicable medical necessity requirements for the intended/prescribed purpose.

Istodax (romidepsin)

Requests for Istodax (romidepsin) may be approved if the following criteria are met:

- I. Individual has a diagnosis of cutaneous T-cell lymphoma; AND
 - A. Individual is using for relapsed or refractory disease following at least one prior systemic therapy;

OR

Individual has a diagnosis of one of the following (NCCN 2A):

- A. Primary cutaneous anaplastic large cell lymphoma (ALCL) with multifocal lesions; OR
- B. Cutaneous ALCL with regional node (N1);

Formatted: Font: Not Bold

#.	AND	
A.III.	Individual has relapsed or refractory disease; AND	
B.IV.	Using romidepsin as monotherapy;	
OR.		
III. —	Individual has a diagnosis of cutaneous ALCL with regional node (N1); AND	
	A. Individual has relapsed or refractory disease; AND	
	B. Using romidepsin as monotherapy;	
OR	In dividual has a discussion of population like the Tool Legachage (NOON OA).	
<u>V.</u>	Individual has a diagnosis of panniculitis-like T-cell Lymphoma (NCCN 2A);	
OB		
<u>OR</u> VI.	Individual has a diagnosis of Mycosis Fungoides or Sézary Syndrome (NCCN 2A);	
<u>VI.</u>	individual has a diagnosis of mycosis rungoldes of Sezary Syndrome (NCCN 2A),	
OR		
VII.	Individual has a diagnosis of one of the following (NCCN 2A):	
<u>v 11.</u>	A. Peripheral T-cell lymphoma; OR	
	B. Relapsed or refractory Breast Implant-associated ALCL; OR	
	C. Refractory Hepatosplenic T-cell Lymphoma after two first-line therapy regimens;	
	AND	
VIII.	Using romidepsin as monotherapy or in combination with duvelisib;	
OR		
IX.	Individual has a diagnosis of Extranodal NK/T-cell lymphomas (NCCN 2A); AND	
Χ.	Individual has relapsed or refractory disease; AND	
IV. XI.	Using romidepsin as monotherapy.	
		
Reques	sts for Istodax (romidepsin) may not be approved when the above criteria are not met and for all other	

Requests for Istodax (romidepsin) may not be approved when the above criteria are not met and for all other indications.

Coding

The following codes for treatments and procedures applicable to this document are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

HCPCS

J9318	Injection, romidepsin, non-lyophilized, 0.1 mg
J9319	Injection, romidepsin, lyophilized, 0.1 mg [Istodax]

ICD-10 Diagnosis

C84.00	Mycosis fungoides, unspecified site
C84.01	Mycosis fungoides, lymph nodes of head, face, and neck
C84.02	Mycosis fungoides, intrathoracic lymph nodes
C84.03	Mycosis fungoides, intra-abdominal lymph nodes
C84.04	Mycosis fungoides, lymph nodes of axilla and upper limb
C84.05	Mycosis fungoides, lymph nodes of inguinal region and lower limb
C84.06	Mycosis fungoides, intrapelvic lymph nodes
C84.07	Mycosis fungoides, spleen
C84.08	Mycosis fungoides, lymph nodes of multiple sites
C84.09	Mycosis fungoides, extranodal and solid organ sites

Formatted: No bullets or numbering

Formatted: Numbered + Level: 1 + Numbering Style: I, II, III, ... + Start at: 1 + Alignment: Right + Aligned at: 0.25" + Indent at: 0.5"

Formatted: Don't add space between paragraphs of the same style, Line spacing: single

Formatted: Don't add space between paragraphs of the same style

Formatted: Font: (Default) Arial, 9 pt, Bold

Formatted: Normal, Indent: Left: 0.06", Don't add space between paragraphs of the same style, No bullets or numbering

Formatted: Don't add space between paragraphs of the same style

Formatted: Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

Formatted: Font: (Default) Arial, 9 pt, Bold, Font color: Black

Formatted: Normal, Indent: Left: 0.06", Don't add space between paragraphs of the same style, No bullets or numbering

Formatted: Don't add space between paragraphs of the same style

Formatted: Don't add space between paragraphs of the same style

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Indent: Left: 0.5", Don't add space between paragraphs of the same style, No bullets or

Formatted: Don't add space between paragraphs of the same style

Formatted: Font: (Default) Arial, 9 pt, Font color: Black

Formatted: Normal, Don't add space between paragraphs of the same style, No bullets or numbering

Formatted: Font: (Default) Arial, 9 pt, Bold, Font color:

Black

Formatted

Formatted

Formatted: Font: Bold

Formatted: Font: Bold

C84.10	Sezary disease, unspecified site
C84.11	Sezary disease, lymph nodes of head, face, and neck
C84.12	Sezary disease, intrathoracic lymph nodes
C84.13	Sezary disease, intra-abdominal lymph nodes
C84.14	Sezary disease, lymph nodes of axilla and upper limb
C84.15	Sezary disease, lymph nodes of inguinal region and lower limb
C84.16	Sezary disease, intrapelvic lymph nodes
C84.17	Sezary disease, spleen
C84.18	Sezary disease, lymph nodes of multiple sites
C84.19	Sezary disease, extranodal and solid organ sites
C84.40	Peripheral T-cell lymphoma, not elsewhere classified, unspecified site
C84.41	Peripheral T-cell lymphoma, not elsewhere classified, lymph nodes of head,
C84.42	face, and neck Peripheral T-cell lymphoma, not elsewhere classified, intrathoracic lymph nodes
C84.43	Peripheral T-cell lymphoma, not elsewhere classified, intra-abdominal lymph
C84.44	nodes Peripheral T-cell lymphoma, not elsewhere classified, lymph nodes of axilla and upper limb
C84.45	Peripheral T-cell lymphoma, not elsewhere classified, lymph nodes of inguinal region and lower limb
C84.46	Peripheral T-cell lymphoma, not elsewhere classified, intrapelvic lymph nodes
C84.47	Peripheral T-cell lymphoma, not elsewhere classified, spleen
C84.48	Peripheral T-cell lymphoma, not elsewhere classified, lymph nodes of multiple sites
C84.49 C84.60	Peripheral T-cell lymphoma, not elsewhere classified, extranodal and solid organ sites Anaplastic large cell lymphoma, ALK-positive, unspecified site
C84.61	Anaplastic large cell lymphoma, ALK-positive, lymph nodes of head, face, and
	neck
C84.62	Anaplastic large cell lymphoma, ALK-positive, intrathoracic lymph nodes
C84.63	Anaplastic large cell lymphoma, ALK-positive, intra-abdominal lymph nodes
C84.64	Anaplastic large cell lymphoma, ALK-positive, lymph nodes of axilla and upper limb
C84.65	Anaplastic large cell lymphoma, ALK-positive, lymph nodes of inguinal region and lower limb
C84.66	Anaplastic large cell lymphoma, ALK-positive, intrapelvic lymph nodes
C84.67	Anaplastic large cell lymphoma, ALK-positive, spleen
C84.68	Anaplastic large cell lymphoma, ALK-positive, lymph nodes of multiple sites
C84.69	Anaplastic large cell lymphoma, ALK-positive, extranodal and solid organ sites
C84.70	Anaplastic large cell lymphoma, ALK-negative, unspecified site
C84.71	Anaplastic large cell lymphoma, ALK-negative, lymph nodes of head, face, and neck
C84.72	Anaplastic large cell lymphoma, ALK-negative, intrathoracic lymph nodes
C84.73	Anaplastic large cell lymphoma, ALK-negative, intra-abdominal lymph nodes
C84.74	Anaplastic large cell lymphoma, ALK-negative, lymph nodes of axilla and upper limb
C84.75	Anaplastic large cell lymphoma, ALK-negative, lymph nodes of inguinal region and lower limb Anaplastic large cell lymphoma, ALK-negative, intrapelyic lymph nodes
C84.76 C84.77	Anaplastic large cell lymphoma, ALK-negative, intrapelvic lymph nodes
684.77	Anaplastic large cell lymphoma, ALK-negative, spleen

C84.79	Anaplastic large cell lymphoma, ALK-negative, extranodal and solid organ sites
C84.90	Mature T/NK-cell lymphomas, unspecified site
C84.91	Mature T/NK-cell lymphomas, lymph nodes of head, face, and neck
C84.92	Mature T/NK-cell lymphomas, intrathoracic lymph nodes
C84.93	Mature T/NK-cell lymphomas, intra-abdominal lymph nodes
C84.94	Mature T/NK-cell lymphomas, lymph nodes of axilla and upper limb
C84.95	Mature T/NK-cell lymphomas, lymph nodes of inquinal region and lower limb
C84.96	Mature T/NK-cell lymphomas, intrapelvic lymph nodes
C84.97	Mature T/NK-cell lymphomas, spleen
C84.98	Mature T/NK-cell lymphomas, lymph nodes of multiple sites
C84.99	Mature T/NK-cell lymphomas, extranodal and solid organ sites
C84.7A	Anaplastic large cell lymphoma, ALK-negative, breast
C84.A0-C84.A9	Cutaneous T-cell lymphoma, unspecified
C84.Z0	Other mature T/NK-cell lymphomas, unspecified site
C84.Z1	Other mature T/NK-cell lymphomas, lymph nodes of head, face, and neck
C84.Z2	Other mature T/NK-cell lymphomas, intrathoracic lymph nodes
C84.Z3	Other mature T/NK-cell lymphomas, intra-abdominal lymph nodes
C84.Z4	Other mature T/NK-cell lymphomas, lymph nodes of axilla and upper limb
C84.Z5	Other mature T/NK-cell lymphomas, lymph nodes of inguinal region and lower limb
C84.Z6	Other mature T/NK-cell lymphomas, intrapelvic lymph nodes
C84.Z7	Other mature T/NK-cell lymphomas, spleen
C84.Z8	Other mature T/NK-cell lymphomas, lymph nodes of multiple sites
C84.Z9	Other mature T/NK-cell lymphomas, extranodal and solid organ sites
C86.0	Extranodal NK/T-cell lymphoma, nasal type
C86.1	Hepatosplenic T-cell lymphoma
C86.2	Enteropathy-type (intestinal) T-cell lymphoma
C86.3	Subcutaneous panniculitis-like T-cell lymphoma
C86.5	Angioimmunoblastic T-cell lymphoma
C86.6	Primary cutaneous CD30-positive T-cell proliferations
Z85.72	Personal history of non-Hodgkin lymphomas
C84.00-C84.09	Mycosis fungoides
C84.10-C84.19	Sezary disease
C84.40-C84.49	Peripheral T-cell lymphoma, not elsewhere classified
C84.60-C84.69	Anaplastic large cell lymphoma, ALK-positive
C84.70-C84.79	Anaplastic large cell lymphoma, ALK-negative

Cutaneous T-cell lymphoma, unspecified

Anaplastic large cell lymphoma, ALK-negative, lymph nodes of multiple sites

C84.78

C84.A0-C84.A9

Formatted Table

C84.Z0-C84.Z9 Other mature T/NK-cell lymphomas C84.90-C84.99 Mature T/NK-cell lymphomas, unspecified C86.00 Extranodal NK/T-cell lymphoma, nasal type not having achieved remission C86.10 Hepatosplenic T-cell lymphoma not having achieved remission C86.20 Enteropathy-type (intestinal) T-cell lymphoma not having achieved remission C86.30 Subcutaneous panniculitis-like T-cell lymphoma not having achieved remission Angioimmunoblastic T-cell lymphoma not having achieved remission C86.50 C86.60 Primary cutaneous CD30-positive T-cell proliferations not having achieved remission Z85.72 Personal history of non-Hodgkin lymphomas

Document History

Revised: 08/15/2025 Document History:

- 08/15/2025 Annual Review: Add use in panniculitis-like T-cell lymphoma, breast implant-associated ALCL, extranodal NK/T-cell lymphomas, hepatosplenic T-cell Lymphoma, and peripheral T-cell lymphomas (PTCL) per NCCN. Coding Reviewed: Updated description for HCPCS J9319. Consolidated ICD-10-CM codes into ranges C84.00-C84.09, C84.10-C84.19, C84.40-C84.49, C84.60-C84.69, C84.70-C84.79, C84.90-C84.99, C84.Z0-C84.Z9. Removed ICD-10-CM C86.0, C86.1, C86.2, C86.3, C86.5, C86.6 and added C86.00, C86.10, C86.20, C86.30, C86.50, C86.60.
- 08/16/2024 Annual Review: Wording and formatting; include brand name. Coding Reviewed: Add ICD-10-CM C84.90, C84.91, C84.92, C84.93, C84.94, C84.95, C84.96, C84.97, C84.98, C84.99, C84.Z0, C84.Z1, C84.Z2, C84.Z3, C84.Z4, C84.Z5, C84.Z6, C84.Z7, C84.Z8, C84.Z9.
- 08/18/2023 Annual Review: Add NCCN 2A recommendation for use in primary cutaneous ALCL or cutaneous ALCL with regional node disease. Wording and formatting. Romidepsin is generically available. Coding Reviewed: Removed the name Istodax from HCPCS J9318, J9319.
- 11/18/2022 Select Review: Update criteria to only include cutaneous T-cell lymphomas per label. Coding Reviewed: No changes.
- 08/19/2022 Annual Review: Remove requirement for "cutaneous" T-cell lymphoma to allow other types per NCCN. Coding Reviewed: Removed ICD-10-CM C84.A0-C84.A9., C84.00-C84.09, C84.10-C84.19, C84.40-C84.49, C84.60-C84.79. Added ICD-10-CM C84.00, C84.01, C84.02, C84.03, C84.04, C84.05, C84.06, C84.07, C84.08, C84.09, C84.11, C84.11, C84.12, C84.13, C84.14, C84.15, C84.16, C84.17, C84.18, C84.19, C84.40, C84.41, C84.42, C84.43, C84.44, C84.45, C84.46, C84.47, C84.48, C84.49. C84.60, C84.61, C84.62, C84.63, C84.63, C84.65, C84.66, C84.67, C84.68, C84.69, C84.70, C84.71, C84.72, C84.73, C84.74, C84.75, C84.76, C84.77, C84.78, C84.79. Effective 10/1/2022 Added ICD-10-CM C84.A0-C84.A9, C86.0, C86.1, C86.3.
- 09/13/2021 Select Review: Update criteria to specify cutaneous T-cell lymphomas to align with updated FDA indication. Coding reviewed: Added HCPCS J9318, J9319. Removed HCPCS J9315, C9065.
- 08/20/2021 Annual Review: No Changes. Coding reviewed: No changes.
- 08/21/2020 Annual Review: No Changes. Coding reviewed: Added HCPCS C9065. Effective 6/9/2021 Do not terminate C9065. Removed HCPCS J9314 per Medicare update 7/1/2021.
- 11/15/2019 Annual Review: No Changes. Coding reviewed: No changes.
- 05/17/2019 Annual Review: First review of Istodax clinical criteria. Add reference for off-label criteria.
 Coding reviewed: No changes.

References

- DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. http://dailymed.nlm.nih.gov/dailymed/about.cfm. Accessed: June 30, 2024.
- 2. DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.
- 3. Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2024; Updated periodically.
- Coiffier B, Pro B, Prince HM, et al. Results from a Pivotal, Open-Label, Phase II Study of Romidepsin in Relapsed or Refractory Peripheral T-Cell Lymphoma after Prior Systemic Therapy. J Clin Oncol 2012; 30: 631-636.

- 5. Bachy E, Camus V, Thieblemont C, et al. Romidepsin Plus CHOP Versus CHOP in Patients with Previously Untreated Peripheral T-Cell Lymphoma: Results of the Ro-CHOP Phase III Study (Conducted by LYSA). J Clin Oncol 2022; 40:242-251.
- NCCN Clinical Practice Guidelines in Oncology™. © 2025 National Comprehensive Cancer Network, Inc. For additional information visit the NCCN website: http://www.nccn.org/index.asp. Accessed on June 16, 2025.

 a. Primary Cutaneous Lymphomas. V3.2025. Revised June 10, 2025.

 - b. T-Cell Lymphomas. V2.2025. Revised May 28, 2025.

Federal and state laws or requirements, contract language, and Plan utilization management programs or polices may take precedence over the application of this clinical criteria.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

© CPT Only - American Medical Association