

## Clinical Policy: Nivolumab (Opdivo)

Reference Number: LA.PHAR.121

Effective Date: ~~04-21~~

Last Review Date: ~~06-23~~04-22

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

**\*\*Please note: This policy is for medical benefit\*\***

### Description

Nivolumab (Opdivo®) is a programmed death receptor-1 (PD-1) blocking antibody.

### FDA Approved Indication(s)

Opdivo is indicated for the treatment of:

- **Melanoma**

- ~~Adult and pediatric (12 years and older) p~~Patients with unresectable or metastatic melanoma, as a single agent or in combination with ipilimumab.
- ~~Adult and pediatric (12 years and older) p~~Patients with melanoma with lymph node involvement or metastatic disease who have undergone complete resection, in the adjuvant setting.

- **Non-small cell lung cancer (NSCLC)**

- ~~Adult patients with resectable (tumors > 4 cm or node positive) NSCLC in the neoadjuvant setting, in combination with platinum-doublet chemotherapy~~
- Adult patients with metastatic NSCLC expressing PD-L1 (≥1%) as determined by an FDA-approved test, with no EGFR or ALK genomic tumor aberrations, as first-line treatment in combination with ipilimumab.
- Adult patients with metastatic or recurrent NSCLC with no EGFR or ALK genomic tumor aberrations as first-line treatment, in combination with ipilimumab and 2 cycles of platinum-doublet chemotherapy.
- ~~Adult p~~Patients with metastatic NSCLC and progression on or after platinum-based chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving Opdivo.

- **Malignant pleural mesothelioma**

- Adult patients with unresectable malignant pleural mesothelioma, as first-line treatment in combination with ipilimumab.

- **Renal cell carcinoma (RCC)**

- Adult patients with advanced renal cell carcinoma (RCC) who have received prior antiangiogenic therapy.
- Adult patients with advanced renal cell carcinoma, as a first-line treatment in combination with cabozantinib.
- Adult patients with intermediate or poor risk advanced RCC, as a first-line treatment in combination with ipilimumab.

## CLINICAL POLICY

### Nivolumab



- **Classical Hodgkin lymphoma (cHL)**
  - Adult patients with cHL that has relapsed or progressed after: \*
    - autologous hematopoietic stem cell transplantation (HSCT) and brentuximab vedotin, or
    - 3 or more lines of systemic therapy that includes autologous HSCT.
- **Squamous cell carcinoma of the head and neck (SCCHN)**
  - **Adult p**Patients with recurrent or metastatic SCCHN with disease progression on or after a platinum-based therapy.
- **Urothelial carcinoma (UC)**
  - Adjuvant treatment of adult patients with UC who are at high risk of recurrence after undergoing radical resection of UC.
  - **Adult p**Patients with locally advanced or metastatic UC who: \*
    - have disease progression during or following platinum-containing chemotherapy, or
    - have disease progression within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy.
- **Colorectal cancer**
  - Adult and pediatric (12 years and older) patients with microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) metastatic colorectal cancer (CRC) that has progressed following treatment with a fluoropyrimidine, oxaliplatin, and irinotecan, as a single agent or in combination with ipilimumab.\*
- **Hepatocellular carcinoma (HCC)**
  - **Adult p**Patients with HCC who have been previously treated with sorafenib in combination with ipilimumab.\*
- **Esophageal cancer**
  - As adjuvant treatment in adult patients with completely resected esophageal or gastroesophageal junction cancer with residual pathologic disease who have received neoadjuvant chemoradiotherapy (CRT).
  - In combination with fluoropyrimidine- and platinum-containing chemotherapy for the first-line treatment of adult patients with unresectable advanced or metastatic esophageal squamous cell carcinoma (ESCC).
  - In combination with ipilimumab for the first-line treatment of adult patients with unresectable advanced or metastatic ESCC.
  - **Adult p**Patients with unresectable advanced, recurrent or metastatic ESCC after prior fluoropyrimidine- and platinum-based chemotherapy.
- **Gastric cancer, gastroesophageal junction cancer, and esophageal adenocarcinoma**
  - **Adult p**Patients with advanced or metastatic gastric cancer, gastroesophageal junction cancer, and esophageal adenocarcinoma in combination with fluoropyrimidine- and platinum-containing chemotherapy.

\*This indication is approved under accelerated approval based on overall or tumor response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trials.

#### Policy/Criteria

*Provider must submit documentation (including such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of Louisiana Healthcare Connections that Opdivo is **medically necessary** when the following criteria are met:

**I. Initial Approval Criteria**

**A. Melanoma** (must meet all):

1. Diagnosis of unresectable, metastatic, or lymph node positive melanoma;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq 12$  years;
4. Request meets one of the following (a, b, or c):\*
  - a. If prescribed as monotherapy (unresectable or metastatic disease, or adjuvant treatment), dose does not exceed any of the following (i or ii):
    - i. Adult and pediatric members weighing  $\geq 40$  kg: 240 mg every 2 weeks or 480 mg every 4 weeks;
    - ii. Pediatric members weighing  $< 40$  kg: 3 mg/kg every 2 weeks or 6 mg/kg every 4 weeks (see Appendix E for dose rounding guidelines);
  - b. If prescribed in combination with Yervoy® (unresectable or metastatic disease), dose does not exceed any of the following (i or ii; see Appendix E for dose rounding guidelines):
    - i. Adult and pediatric members weighing  $\geq 40$  kg: 1 mg/kg every 3 weeks for 4 doses, following by 240 mg every 2 weeks or 480 mg every 4 weeks (see Appendix E for dose rounding guidelines);
    - ii. Pediatric members weighing  $< 40$  kg: 1 mg/kg every 34 weeks for 4 doses, followed by 3 mg/kg every 2 weeks or 6 mg/kg every 4 weeks
  - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

\*Prescribed regimen must be FDA-approved or recommended by NCCN

**Approval duration: 6 months**

**B. Non-Small Cell Lung Cancer** (must meet all):

1. Diagnosis of resectable, recurrent, advanced, or metastatic NSCLC;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq 18$  years;
4. Member has not previously progressed on a PD-1/PD-L1 inhibitor (e.g., Keytruda®, Tecentriq®, Imfinzi®);
5. For resectable NSCLC: Both of the following are met (a and b):
  - a. Opdivo is prescribed as neoadjuvant treatment;
  - b. Tumors  $\geq 4$  cm or node positive disease;
- 4.6. For recurrent, advanced, or metastatic NSCLC: Opdivo is prescribed in one of the following ways (a, b, or c):
  - a. For use as a single agent, and disease has progressed on or after systemic therapy;
  - b. For use as a single agent or in combination with Yervoy for tumors positive for the Tumor Mutation Burden (TMB) biomarker;
  - c. For use in combination with Yervoy, and both of the following (i and ii):
    - i. Request meets one of the following (a, b, or c):

- a) Disease mutation status is unknown or negative for EGFR, ALK, ROS1, BRAF, MET exon 14 skipping, and RET, and member has not received prior systemic therapy for advanced disease;
- b) Disease mutation status is positive for EGFR, ALK, ROS1, BRAF, MET exon 14 skipping, RET, or NTRK gene fusion, and member has received mutation-specific treatment;
- c) Disease is positive for a RET rearrangement;
- ii. Request meets one of the following (a or b):
  - a) Member has PD-L1 tumor expression of  $\geq 1\%$ ;
  - b) Opdivo is being used in combination with Yervoy  $\pm$  a platinum-based regimen (*see Appendix B*);

*\*Prior authorization may be required for Yervoy*

5-7. Request meets one of the following (a, b, c, d, or e):\*

- a. Monotherapy: Dose does not exceed 240 mg every 2 weeks or 480 mg every 4 weeks;
- b. In combination with Yervoy: Dose does not exceed 3 mg/kg every 2 weeks (*see Appendix E for dose rounding guidelines*);
- c. In combination with Yervoy and platinum-doublet chemotherapy: Dose does not exceed 360 mg every 3 weeks;
- d. In combination with platinum-doublet chemotherapy, both of the following are met (i and ii):
  - i. Dose does not exceed 360 mg every 3 weeks
  - ii. Request does not exceed 3 cycles
- e. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months** (9 weeks for neoadjuvant NSCLC)

**C. Malignant Pleural Mesothelioma** (must meet all):

- 1. Diagnosis of unresectable malignant pleural mesothelioma;
- 2. Prescribed by or in consultation with an oncologist;
- 3. Age  $\geq 18$  years;
- 4. Prescribed in one of the following ways (a or b):
  - a. As first-line therapy in combination with Yervoy;
  - b. If not administered first-line, as subsequent therapy in combination with Yervoy or as a single agent;
- 5. Request meets one of the following (a or b):\*
  - a. Dose does not exceed 360 mg every 3 weeks;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

**D. Renal Cell Carcinoma** (must meet all):

- 1. Diagnosis of RCC;
- 2. Prescribed by or in consultation with an oncologist;
- 3. Age  $\geq 18$  years;

4. Request meets one of the following (a, b, or c):\*
  - a. Monotherapy or in combination with cabozantinib: Dose does not exceed 240 mg every 2 weeks or 480 mg every 4 weeks;
  - b. In combination with Yervoy: Dose does not exceed 3 mg/kg every 3 weeks for 4 doses, followed by 240 mg every 2 weeks or 480 mg every 4 weeks (*see Appendix E for dose rounding guidelines*);
  - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

\*Prescribed regimen must be FDA-approved or recommended by NCCN

**Approval duration: 6 months**

**E. Classical Hodgkin Lymphoma (must meet all):**

1. Diagnosis of relapsed, refractory or progressive cHL;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq 18$  years;
4. Prescribed as subsequent therapy;
5. Request meets one of the following (a or b):\*
  - a. Dose does not exceed 240 mg every 2 weeks or 480 mg every 4 weeks;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

\*Prescribed regimen must be FDA-approved or recommended by NCCN

**Approval duration: 6 months**

**F. Squamous Cell Carcinoma of the Head and Neck (must meet all):**

1. Diagnosis of SCCHN;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq 18$  years;
- 3-4. Prescribed as a single agent;
- 4-5. Disease has progressed on or after a platinum-containing regimen (e.g., cisplatin, carboplatin);
- 5-6. Request meets one of the following (a or b):\*
  - a. Dose does not exceed 240 mg every 2 weeks or 480 mg every 4 weeks;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

\*Prescribed regimen must be FDA-approved or recommended by NCCN

**Approval duration: 6 months**

**G. Urothelial Carcinoma (must meet all):**

1. Diagnosis of UC;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq 18$  years;
4. One of the following (a, b, or c):
  - a. Failure of a platinum-containing regimen (e.g., cisplatin, carboplatin), unless clinically significant adverse effects are experienced or all are contraindicated;
  - b. Prescribed as adjuvant treatment and member is at high risk of recurrence after undergoing resection of UC;

- c. Member is at high risk of recurrence and did not previously receive a platinum-containing regimen;
- 5. Request meets one of the following (a or b):\*
  - a. Dose does not exceed 240 mg every 2 weeks or 480 mg every 4 weeks;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

**H. Colorectal Cancer (must meet all):**

- 1. Diagnosis of unresectable or metastatic CRC;
- 2. Tumor is characterized as MSI-H or dMMR;
- 3. Prescribed by or in consultation with an oncologist;
- 4. Age  $\geq$  12 years;
- 5. Dose does not exceed one of the following (a, b, or c):\*
  - a. Monotherapy: 240 mg every 2 weeks;
  - b. In combination with Yervoy: 3 mg/kg every 3 weeks for 4 doses, then 240 mg every 2 weeks or 480 mg every 4 weeks (*see Appendix E for dose rounding guidelines*);
  - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

**I. Hepatocellular Carcinoma (must meet all):**

- 1. Diagnosis of HCC;
- 2. Prescribed by or in consultation with an oncologist;
- 3. Age  $\geq$  18 years;
- 4. Member has had disease progression following treatment with Nexavar<sup>®</sup>, ~~Lenvima<sup>®</sup>~~, ~~Tecentriq<sup>®</sup>~~, bevacizumab (Mvasi<sup>®</sup> and Zirabev<sup>™</sup> are preferred), or Imfinzi<sup>®</sup>;  
*\*Prior authorization may be required for Nexavar, Lenvima, Tecentriq, bevacizumab, and Imfinzi.*  
*\*Prior authorization may be required for Nexavar or Lenvima*
- 5. ~~Member has not had previous treatment with a checkpoint inhibitor (e.g., Yervoy, Keytruda, Tecentriq, Imfinzi);~~ Prescribed in combination with Yervoy
- 6. ~~Request is for use in combination Yervoy and~~ Documentation supports of Child-Pugh Class A status;
- 7. Dose does not exceed one of the following (a or b):\*
  - a. In combination with Yervoy: 1 mg/kg every 3 weeks for 4 doses, then 240 mg every 2 weeks or 480 mg every 4 weeks (*see Appendix E for dose rounding guidelines*);
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

Formatted: Font: 10 pt, Italic

**J. Esophageal Cancer** (must meet all):

1. Diagnosis of one of the following (a or b):
  - a. Completely resected esophageal cancer or gastroesophageal junction (esophagogastric junction; EGJ) cancer;
  - b. Unresectable advanced, recurrent, or metastatic ESCC;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq$  18 years;
4. For completely resected esophageal cancer or EGJ cancer, member meets both of the following (a and b):
  - a. Member has residual pathologic disease;
  - b. Member has previously received CRT;

**5. For ESCC, one of the following (a or b):**

a. For unresectable advanced or metastatic disease: Prescribed in combination with Yervoy or with fluoropyrimidine- and platinum-containing chemotherapy;

e-b. For unresectable advanced, recurrent, or metastatic ~~ESCC~~ disease: Member has had previous treatment with a fluoropyrimidine-based (e.g., 5-fluorouracil, capecitabine) and platinum-based (e.g., carboplatin, cisplatin, oxaliplatin) chemotherapy;

**5-6.** Request meets one of the following (a, b, or c):\*

- a. ESCC in combination with Yervoy: Dose does not exceed 3 mg/kg every 2 weeks or 360 mg every 3 weeks;
- b. Other indications: Dose does not exceed 240 mg every 2 weeks or 480 mg every 4 weeks;
- c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

**K. Gastric and Esophageal Adenocarcinomas** (must meet all):

1. Diagnosis of gastric cancer, EGJ cancer, or esophageal adenocarcinoma;
2. Member meets one of the following (a or b):
  - a. Disease is advanced, recurrent, or metastatic;
  - b. For EGJ cancer or esophageal adenocarcinoma: member meets one of the following (i or ii):
    - i. Member is post-operative following chemoradiation;
    - ii. Disease is advanced, recurrent, or metastatic;
3. Prescribed by or in consultation with an oncologist;
4. Age  $\geq$  18 years;
5. For advanced, recurrent, or metastatic disease: both of the following are met (a and b):
  - a. Prescribed in combination with a fluoropyrimidine- (e.g., 5-fluorouracil, capecitabine) and platinum-containing (e.g., carboplatin, cisplatin, oxaliplatin) chemotherapy;
  - b. Disease is HER2-negative;
6. Request meets one of the following (a or b):\*
  - a. Dose does not exceed 240 mg every 2 weeks or 360 mg every 3 weeks;

- b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

**L. Off-label NCCN Compendium Recommended Indications (must meet all):**

1. Diagnosis of one of the following (a-~~o~~k):
  - a. Squamous cell anal carcinoma that is metastatic;
  - b. Merkel cell carcinoma;
  - c. Gestational trophoblastic neoplasia;
  - d. Uveal melanoma that is metastatic;
  - e. Small bowel adenocarcinoma that is advanced or metastatic;
  - f. Extranodal NK/T-cell lymphoma, nasal type, that is relapsed or refractory;
  - g. Pediatric Hodgkin lymphoma, as subsequent therapy;
  - h. Vulvar cancer - HPV-related advanced, recurrent, or metastatic disease, as second-line treatment;
  - i. Cervical cancer;
  - j. Endometrial carcinoma that is recurrent or metastatic;
  - ~~k.~~ Small cell lung cancer, as subsequent therapy;
  - l. Bone cancer (e.g., Ewing Sarcoma, chordoma, osteosarcoma, chondrosarcoma);
  - m. Central nervous system (CNS) cancer (e.g., brain metastases);
  - n. Pediatric primary mediastinal large B-cell lymphoma;
  - ~~k-o.~~ Pediatric diffuse high-grade gliomas;
2. Prescribed by or in consultation with an oncologist;
- ~~3. Age > 18 years~~
- ~~4-3.~~ For anal carcinoma: prescribed as second line or subsequent therapy (examples of prior therapy include 5-FU/cisplatin, carboplatin/paclitaxel, FOLFOX, FOLFCIS);
- ~~5-4.~~ For gestational trophoblastic neoplasia: prescribed as a single agent for multi-agent chemotherapy-resistant disease (*see Appendix B*) in one of the following settings (a or b):
  - a. Recurrent or progressive intermediate trophoblastic tumor following treatment with a platinum-containing regimen (e.g., cisplatin, carboplatin);
  - b. High-risk disease (*see Appendix D*);
5. For pediatric primary mediastinal large B-cell lymphoma: prescribed as one of the following (a or b):
  - a. As a single agent as second line therapy after failure of induction therapy/initial treatment (*see appendix B*);
  - b. Combination with brentuximab vedotin as consolidation/additional therapy;
6. For pediatric diffuse high-grade gliomas: prescribed as a single agent for adjuvant therapy or for recurrent/progressive disease;
7. For uveal melanoma, bone cancer, CNS cancer: prescribed as a single agent or in combination with Yervoy;  
*\*Prior authorization may be required for Yervoy.*
8. For cervical cancer: prescribed as second line or subsequent therapy for PD-L1 tumor expression of  $\geq 1\%$ ;



9. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).\*

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

**M. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53
- ~~1. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.~~

**II. Continued Therapy**

**A. All Indications in Section I (must meet all):**

1. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving Opdivo for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a, b, c, d, e, or f ~~or e~~):\*
  - a. NSCLC in combination with Yervoy: New dose does not exceed 3 mg/kg every 2 weeks;
  - b. Malignant pleural mesothelioma in combination with Yervoy, and gastric and esophageal adenocarcinomas: New dose does not exceed 360 mg every 3 weeks;
  - c. ESCC in combination with Yervoy: New dose does not exceed 3 mg/kg every 2 weeks or 360 mg every 3 weeks;
  - d. Melanoma (i or ii):
    - i. If prescribed as monotherapy (unresectable or metastatic disease, or adjuvant treatment), new dose does not exceed any of the following (a or b):
      - a) Adult and pediatric members weighing  $\geq$  40 kg: 240 mg every 2 weeks or 480 mg every 4 weeks;
      - b) Pediatric members weighing  $<$  40 kg: 3 mg/kg every 2 weeks or 6 mg/kg every 4 weeks;
    - ii. If prescribed in combination with Yervoy (unresectable or metastatic disease), new dose does not exceed any of the following (a or b):
      - a) Adult and pediatric members weighing  $\geq$  40kg: 1 mg/kg every 3 weeks for 4 doses, followed by 240 mg every 2 weeks or 480 mg every 4 weeks;
      - b) Pediatric members weighing  $<$  40 kg: 1 mg/kg every 3 weeks for 4 doses, followed by 3 mg/kg every 2 weeks or 6 mg/kg every 4 weeks;
  - e-e. Other indications: New dose does not exceed 240 mg every 2 weeks or 480 mg every 4 weeks;

4.f. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 12 months**

**B. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53
- ~~1. Currently receiving medication via Louisiana Healthcare Connections benefit and documentation supports positive response to therapy. Approval duration: Duration of request or 6 months (whichever is less); or~~
- ~~2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.~~

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy LA.PMN.53 ~~for Medicaid, or evidence of coverage documents.~~

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

ALK: anaplastic lymphoma kinase	receptor-2
BRAF: B-Raf proto-oncogene, serine/threonine kinase	HSCT: hematopoietic stem cell transplantation
CHL: classic Hodgkin lymphoma	MET: mesenchymal-epithelial transition
CNS: central nervous system	MSI-H: microsatellite instability-high
CRC: colorectal cancer	NSCLC: non-small cell lung cancer
dMMR: mismatch repair deficient	PD-1: programmed death receptor-1
EGFR: epidermal growth factor receptor	PD-L1: programmed death-ligand 1
EGJ: esophagogastric junction	RCC: renal cell carcinoma
ESCC: esophageal squamous cell carcinoma	ROS1: ROS proto-oncogene 1
FDA: Food and Drug Administration	SCLC: small cell lung cancer
HCC: hepatocellular carcinoma	TMB: Tumor Mutational Burden
HER-2: human epidermal growth factor	UC: urothelial carcinoma

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.*

**CLINICAL POLICY**  
Nivolumab



Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Nexavar (sorafenib)	HCC: 400 mg PO BID until clinical benefit ceases or unacceptable toxicity occurs	800 mg/day
Lenvima (lenvatinib)	HCC: 12 mg PO QD (patients $\geq$ 60 kg) or 8 mg PO QD (patients < 60 kg) until disease progression or unacceptable toxicity	12 mg/day
<u>Tecentriq (atezolizumab) + bevacizumab (Avastin<sup>®</sup>, Mvasi, Zirabev)</u>	<u>HCC</u> <u>Tecentriq: 840 mg IV every 2 weeks, 1,200 mg IV every 3 weeks, or 1,680 mg IV every 4 weeks</u> <u>Bevacizumab: 15 mg/kg IV every 3 weeks</u>	<u>See regimen</u>
<u>Imfinzi (durvalumab)*</u>	<u>HCC</u> <u>Varies</u>	<u>Varies</u>
<u>Cisplatin or carboplatin-containing chemotherapy</u>	<u>SCLC, UC, SCCHN: Varies</u>	<u>Varies</u>
First-line therapies (e.g., 5-FU/cisplatin, carboplatin/paclitaxel, FOLFOX, FOLFICIS)	Metastatic anal carcinoma: Varies	Varies
First-line therapies (e.g., platinum/etoposide-containing regimen)	Gestational trophoblastic neoplasia: Varies	Varies
platinum-containing regimens	NSCLC – squamous cell carcinoma: paclitaxel + carboplatin dose varies  NSCLC – nonsquamous cell carcinoma: pemetrexed + [carboplatin or cisplatin] dose varies  UC, SCCHN: Varies	Varies
Multiagent chemotherapy regimens examples: EMA/CO (etoposide, methotrexate, dactinomycin/cyclophosphamide, vincristine), EMA/EP (etoposide, methotrexate, dactinomycin/etoposide, cisplatin)	Gestational Trophoblastic Neoplasia: Varies	Varies

**CLINICAL POLICY**  
Nivolumab



Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
<u>Dose-adjusted-EPOCH-R, R-CHOP with radiation therapy, or LMB-modified B/C chemotherapy with rituximab</u>	<u>Pediatric primary mediastinal large B-cell lymphoma:</u> <u>Varies</u>	<u>Varies</u>
<u>Yervoy (ipilimumab)</u>	<u>Melanoma, HCC: 3 mg/kg IV every 3 weeks for a maximum of 4 doses</u>  <u>RCC, CRC: 1 mg/kg IV every 3 weeks for a maximum of 4 doses</u>  <u>NSCLC, malignant pleural mesothelioma, ESCC: 1 mg/kg IV every 6 weeks</u>	<u>See regimen</u>

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

\*Off-label

**Appendix C: Contraindications/Boxed Warnings**

None reported

**Appendix D: General Information**

- High-risk disease in gestational trophoblastic neoplasia is defined as having a FIGO stage IV-I to III and or a prognostics score  $\geq 7$ .~~prognostic score or stage IV.~~

- FIGO staging system:

Stage	Criteria
I	Tumor confined to uterus
II	Tumor extends to other genital structures (ovary, tube, vagina, broad ligaments) by metastasis or direct extension
III	Lung metastasis
IV	All other distant metastases

- Prognostic Scoring Index

- The total score is obtained by adding the individual scores for each prognostic factor (low risk is indicated by a score  $< 7$  and high risk is indicated by a score  $\geq 7$ )

Prognostic factor	Risk score			
	0	1	2	4
Age (years)	$< 40$	$\geq 40$	--	--
Antecedent pregnancy	Hydatidiform mole	Abortion	Term pregnancy	--
Interval from index	$< 4$	4 to 6	7 to 12	$> 12$

**CLINICAL POLICY**  
Nivolumab



Prognostic factor	Risk score			
	0	1	2	4
pregnancy (months)				
Pretreatment hCG (IU/L)	< 10 <sup>3</sup>	10 <sup>3</sup> to < 10 <sup>4</sup>	10 <sup>4</sup> to 10 <sup>5</sup>	≥ 10 <sup>5</sup>
Largest tumor size, including uterus (cm)	< 3	3 to 5	> 5	
Site of metastases	Lung	Spleen, kidney	Gastrointestinal tract	Brain, liver
Number of metastases identified	0	1 to 4	5 to 8	> 8
Previous failed chemotherapy	--	--	Single drug	Two or more drugs
Total score	--	--	--	--

*Appendix E: Dose Rounding Guidelines\**

Weight-based Dose Range	Vial Quantity Recommendation
≤ 41.99 mg	1 vial of 40 mg/4 mL
42 mg-104.99 mg	1 vial of 100 mg/10 mL
105 mg-146.99 mg	1 vial of 40 mg/4 mL and 100 mg/10 mL
147 mg-209.99 mg	2 vials of 100 mg/10 mL
210 mg-251.99 mg	1 vial of 240 mg/24 mL
260 mg-293.99 mg	1 vial of 40 mg/4 mL and 240 mg/24 mL
294 mg-356.99 mg	1 vial of 100 mg/4 mL and 240 mg/24 mL
357 mg-503.99 mg	2 vials of 240 mg/24 mL

*\*This is part of a dose rounding guideline on select drug classes as part of an initiative conducted on a larger scale with multiple references and prescriber feedback.*

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
Melanoma (unresectable or metastatic)	<p>Monotherapy: <del>240 mg IV every 2 weeks or 480 mg IV every 4 weeks</del></p> <ul style="list-style-type: none"> <li>• <u>Adult and pediatric patients weighing ≥ 40 kg: 240 mg IV every 2 weeks or 480 mg IV every 4 weeks</u></li> <li>• <u>Pediatric patients weighing &lt; 40 kg: 3 mg/kg IV every 2 weeks or 6 mg/kg IV every 4 weeks</u></li> </ul> <p><u>With ipilimumab: 1 mg/kg IV, followed by ipilimumab on the same day, every 3 weeks for</u></p>	<del>480mg/dose</del> See <a href="#">regimen</a>

**CLINICAL POLICY**  
Nivolumab



Indication	Dosing Regimen	Maximum Dose
	<p><del>4 doses, then nivolumab 240 mg IV every 2 weeks or 480 mg IV every 4 weeks</del></p> <ul style="list-style-type: none"> <li>Adult and pediatric patients weighing <math>\geq 40</math> kg: 1 mg/kg IV, followed by ipilimumab on the same day, every 3 weeks for 4 doses, then nivolumab 240 mg IV every 2 weeks or 480 mg IV every 4 weeks</li> <li>Pediatric patients weighing <math>&lt; 40</math> kg: 1 mg/kg IV, followed by ipilimumab on the same day, every 3 weeks for 4 doses, then nivolumab 3 mg/kg IV every 3 weeks or 6 mg/kg mg IV every 6 weeks</li> </ul>	
Melanoma (adjuvant treatment) <del>RCC - advanced with previous anti-angiogenic therapy, cHL, SCCHN, UC</del>	<ul style="list-style-type: none"> <li>Adult and pediatric patients weighing <math>\geq 40</math> kg: 240 mg IV every 2 weeks or 480 mg IV every 4 weeks</li> <li>Pediatric patients weighing <math>&lt; 40</math> kg: 3 mg/kg IV every 2 weeks or 6 mg/kg IV every 4 weeks</li> </ul> <p><del>240 mg IV every 2 weeks or 480 mg IV every 4 weeks</del></p>	480 mg/dose <del>See regimen</del>
<del>RCC - advanced with previous anti-angiogenic therapy, cHL, SCCHN, UC</del>	<del>240 mg IV every 2 weeks or 480 mg IV every 4 weeks</del>	<del>480 mg/dose</del>
MSI-H/dMMR CRC	<p>Monotherapy: 240 mg IV every 2 weeks or 480 mg IV every 4 weeks</p> <p>With ipilimumab: 3 mg/kg IV, followed by ipilimumab 1 mg/kg on the same day every 3 weeks for 4 doses, then nivolumab 240 mg IV every 2 weeks or 480 mg IV every 4 weeks</p>	<p>Monotherapy: 480 mg/dose</p> <p>With ipilimumab: 3 mg/kg/dose</p>
RCC - advanced previously untreated	<p>Monotherapy or with cabozantinib: 240 mg IV every 2 weeks or 480 mg every 4 weeks</p> <p>With ipilimumab: 3 mg/kg IV, followed by ipilimumab 1 mg/kg IV on the same day every 3 weeks for 4 doses, then nivolumab 240 mg IV every 2 weeks or 480 mg IV every 4 weeks</p>	480 mg/dose
HCC	With ipilimumab: nivolumab 1 mg/kg IV, followed by ipilimumab 3 mg/kg IV on the same day, every 3 weeks for a maximum of 4 doses, then as single-agent nivolumab 240 mg IV every 2 weeks or 480 mg IV every 4 weeks until disease progression or unacceptable toxicity	480 mg/dose
NSCLC	Monotherapy: 240 mg IV every 2 weeks or 480 mg IV every 4 weeks until disease progression or unacceptable toxicity	Monotherapy: 480 mg/dose

**CLINICAL POLICY**  
Nivolumab



Indication	Dosing Regimen	Maximum Dose
	<p>With ipilimumab: nivolumab 3 mg/kg IV every 2 weeks and ipilimumab 1 mg/kg IV every 6 weeks until disease progression, unacceptable toxicity, or for up to 2 years in patients without disease progression</p> <p>With ipilimumab and platinum-doublet chemotherapy: nivolumab 360 mg IV every 3 weeks and ipilimumab 1 mg/kg IV every 6 weeks and histology-based platinum-doublet chemotherapy every 3 weeks for 2 cycles until disease progression, unacceptable toxicity, or up to 2 years in patients without disease progression</p> <p><u>With platinum-doublet chemotherapy: nivolumab 360 mg IV every 3 weeks with platinum-doublet chemotherapy on the same day every 3 weeks for 3 cycles</u></p>	<p>With ipilimumab: 3 mg/kg/dose</p> <p>With <del>ipilimumab</del> and platinum doublet <u>with or without ipilimumab</u>: 360 mg/dose</p>
Esophageal cancer	<p><u>Adjuvant treatment of resected esophageal or GEJ cancer: 240 mg IV every 2 weeks or 480 mg IV every 4 weeks for a total treatment duration of 1 year</u></p> <p><u>ESCC: until disease progression, unacceptable toxicity, or up to 2 years:</u></p> <p><u>As a single agent or in combination with fluoropyrimidine- and platinum-containing chemotherapy: 240 mg every 2 weeks or 480 mg every 4 weeks</u></p> <ul style="list-style-type: none"> <li><u>In combination with ipilimumab: nivolumab 3 mg/kg every 2 weeks or 360 mg every 3 weeks with ipilimumab 1 mg/kg every 6 weeks</u> <del>240 mg IV every 2 weeks or 480 mg IV every 4 weeks until disease progression or unacceptable toxicity</del></li> </ul>	<p><del>480 mg/dose</del> See regimen</p>
Gastric cancer, EGJ cancer, and esophageal adenocarcinoma	240 mg every 2 weeks or 360 mg every 3 weeks	360 mg/dose

Indication	Dosing Regimen	Maximum Dose
Malignant pleural mesothelioma	With ipilimumab: nivolumab 360 mg every 3 weeks and ipilimumab 1 mg/kg every 6 weeks	With ipilimumab: 360 mg/dose

#### VI. Product Availability

Single-dose vials: 40 mg/4 mL, 100 mg/10 mL, 120 mg/12 mL, 240 mg/24 mL

#### VII. References

1. Opdivo Prescribing Information. Princeton, NJ: Bristol-Myers Squibb; ~~September-February 2023~~. Available at <https://www.opdivo.com/>. Accessed ~~November-March 16, 2023~~.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at <http://www.nccn.org>. Accessed ~~November-April 22~~13, 2023.
3. National Comprehensive Cancer Network. Non-Small Cell Lung Cancer Version ~~5.7.2022~~. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/nscl.pdf](https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf). Accessed November ~~04~~22, 2022.
4. National Comprehensive Cancer Network. Kidney Cancer, Version 3.2023. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/kidney.pdf](https://www.nccn.org/professionals/physician_gls/pdf/kidney.pdf). Accessed November ~~04~~22, 2022.
5. Hellman MD, Paz-Ares L, Bernabe Caro R, et al. Nivolumab plus ipilimumab in advanced non-small-cell lung cancer. N Engl J Med. 2019 November; 381(21):2020-2031.
6. National Comprehensive Cancer Network. Bladder Cancer, Version ~~2.5.2022~~. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/bladder.pdf](https://www.nccn.org/professionals/physician_gls/pdf/bladder.pdf). Accessed November ~~04~~22, 2022.
7. National Comprehensive Cancer Network. Hepatobiliary Cancers, Version ~~3.5.2022~~. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/hepatobiliary.pdf](https://www.nccn.org/professionals/physician_gls/pdf/hepatobiliary.pdf). Accessed November ~~4~~22, 2022. updated criteria for melanoma to reflect FDA approved pediatric age extension; updated Appendix B. National Comprehensive Cancer Network. Esophageal and Esophagogastric Junction Cancers, Version 4.2022. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/esophageal.pdf](https://www.nccn.org/professionals/physician_gls/pdf/esophageal.pdf). Accessed November 4, 2022.
8. National Comprehensive Cancer Network. Pediatric Central Nervous System Cancers, Version 2.2023. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/ped\\_cns.pdf](https://www.nccn.org/professionals/physician_gls/pdf/ped_cns.pdf). Accessed November 4, 2022.
9. National Comprehensive Cancer Network. Central Nervous System Cancers, Version 2.2022. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/cns.pdf](https://www.nccn.org/professionals/physician_gls/pdf/cns.pdf). Accessed November 4, 2022.
10. National Comprehensive Cancer Network. Pediatric Aggressive Mature B-Cell Lymphomas, Version 3.2022. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/ped\\_b-cell.pdf](https://www.nccn.org/professionals/physician_gls/pdf/ped_b-cell.pdf). Accessed November 4, 2022.
11. National Comprehensive Cancer Network. Bone Cancer, Version 2.2023. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/bone.pdf](https://www.nccn.org/professionals/physician_gls/pdf/bone.pdf). Accessed November 4, 2022.
12. National Comprehensive Cancer Network. Melanoma: Cutaneous, Version 2.2023. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/cutaneous\\_melanoma.pdf](https://www.nccn.org/professionals/physician_gls/pdf/cutaneous_melanoma.pdf). Accessed April 13, 2023.



## CLINICAL POLICY

### Nivolumab



7.13.

#### Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPSC Codes	Description
J9299	Injection, nivolumab, 1 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy	01.21	04.21
FDA approved malignant pleural mesothelioma added. Per FDA/NCCN as follows: for melanoma, unresectable, metastatic, or lymph node positive disease added; for NSCLC, single-agent therapy for TMB positive tumor added, combination therapy for RET rearrangement added, combination therapy changed from Yervoy and platinum doublet therapy to Yervoy plus/minus a platinum based regimen; for cHL, relapsed, refractory or progressive disease added, post HSCT replaced with prescribed as subsequent therapy; for HCC, Lenvima added as a prior therapy option, added documentation of Child-Pugh class status; off-label pediatric Hodgkin lymphoma and vulvar cancer added; SCLC criteria per label update; added new FDA approved indication of use in combination with cabozantinib as first-line therapy for advanced RCC; Added new FDA-approved indications of gastric cancer, gastroesophageal junction cancer, and esophageal adenocarcinoma; Added new FDA-approved indication of completely resected esophageal or gastroesophageal junction cancer; Per updated prescribing information removed use in HCC as a single agent; for UC added indication for adjuvant treatment; updates made per NCCN: for urothelial carcinoma removed requirement for resection to be radical as NCCN also supports partial resection prior to adjuvant therapy and added treatment option of highrisk recurrence as an optional criterion; added cervical cancer as off-label indication; updated gestational trophoblastic neoplasia treatment settings; added criterion for use as singleagent therapy for SCCHN; clarified uveal melanoma to be metastatic; removed "metastatic" designation for Merkel cell carcinoma; clarified small bowel adenocarcinoma be advanced or metastatic; small cell lung cancer indication added; clarified extranodal NK/T-cell lymphoma to be relapsed or refractory.	04.22	07.23.22
<a href="#">Added new FDA-approved indication of neoadjuvant use in NSCLC. Criteria added for new FDA approved indication for first-line use in ESCC in combination with Yervoy or with fluoropyrimidine- and</a>	06.27.23	

**CLINICAL POLICY**  
Nivolumab



Reviews, Revisions, and Approvals	Date	LDH Approval Date
<a href="#">platinum-containing chemotherapy; for HCC, added additional options for prior use of Tecentriq+bevacizumab or Imfinzi and removed requirement for no previous treatment with a checkpoint inhibitor per latest NCCN guidelines.</a> <a href="#">Added off-label criteria for bone cancer, central nervous system cancers, pediatric primary mediastinal large Bcell lymphoma, pediatric diffuse high-grade gliomas per NCCN 2A recommendations; removed age restriction from off-label criteria; updated Appendix D to simplify definition of high-risk disease in GTN to mirror the 2023 NCCN GTN guidelines.</a> <a href="#">Template changes applied to other diagnoses/indications.</a> <a href="#">References reviewed and updated.</a> <a href="#">Added blurb this for medical benefit only.</a> <a href="#">Updated criteria for melanoma to reflect FDA approved pediatric age extension; updated Appendix B.</a>		

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

## CLINICAL POLICY

### Nivolumab



This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2023 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.