

Clinical Policy: Cipaglucosidase Alfa-atga + Miglustat (Pombiliti + Opfolda)

Reference Number: LA.PHAR.567

Effective Date:

Last Review Date: 03.28.24

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

****Please note: This policy is for medical benefit****

Description

Cipaglucosidase alfa-atga + miglustat (Pombiliti™ + Opfolda™) is a combination therapy of hydrolytic lysosomal glycogen-specific recombinant human α -glucosidase (rhGAA) enzyme (cipaglucosidase alfa-atga) with an enzyme stabilizer (miglustat).

FDA Approved Indication(s)

Pombiliti is indicated for use in combination with Opfolda for the treatment of adult patients with late-onset Pompe disease (lysosomal acid alpha-glucosidase [GAA] deficiency) weighing \geq 40 kg and who are not improving on their current enzyme replacement therapy (ERT).

Opfolda is indicated for use in combination with Pombiliti for the treatment of adult patients with late-onset Pompe disease (lysosomal GAA deficiency) weighing \geq 40 kg and who are not improving on their current ERT.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that Pombiliti + Opfolda are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Pompe Disease (must meet all):

1. Diagnosis of late-onset Pompe disease confirmed by one of the following (a or b):
 - a. Enzyme assay confirming low GAA activity;
 - b. DNA testing;
2. Age \geq 18 years;
3. Member weighs \geq 40 kg;
4. Pombiliti and Opfolda are prescribed together;
5. Pombiliti and Opfolda are not prescribed concurrently with Lumizyme® or Nexvazyme®;
6. Dose does not exceed any of the following (a or b):
 - a. Members weighing \geq 50 kg: Pombiliti 20 mg/kg + Opfolda 260 mg (or 4 capsules) every other week;

- b. Members weighing ≥ 40 kg to < 50 kg: Pombiliti 20 mg/kg + Opfolda 195 mg (or 3 capsules) every other week.

Approval duration: 6 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the PDL, the no coverage criteria policy: LA.PMN.255 for Medicaid; or
 - b. For drugs NOT on the PDL, the non-formulary policy: LA.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

II. Continued Therapy

A. Pompe Disease (must meet all):

1. Member is currently receiving medication via Louisiana Healthcare Connections benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy as evidenced by improvement in the individual member's Pompe disease manifestation profile (*see Appendix D for examples*);
3. Pombiliti and Opfolda are prescribed together;
4. Pombiliti and Opfolda are not prescribed concurrently with Lumizyme[®] or Nexvazyme[®];
5. If request is for a dose increase, new dose does not exceed any of the following (a or b):
 - a. Members weighing ≥ 50 kg: Pombiliti 20 mg/kg + Opfolda 260 mg (or 4 capsules) every other week;
 - b. Members weighing ≥ 40 kg to < 50 kg: Pombiliti 20 mg/kg + Opfolda 195 mg (or 3 capsules) every other week.

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the PDL, the no coverage criteria policy: LA.PMN.255 for Medicaid; or
 - b. For drugs NOT on the PDL, the non-formulary policy: LA.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND

criterion 1 above does not apply, refer to the off-label use policy: LA.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – LA.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

6MWT: 6-minute walk test

ERT: enzyme replacement therapy

FDA: Food and Drug Administration

GAA: acid alpha-glucosidase

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): pregnancy
- Boxed warning(s): (*Pombiliti only*) severe hypersensitivity reactions, infusion-associated reactions, and risk of acute cardiorespiratory failure in susceptible patients

Appendix D: Measures of Therapeutic Response

- Pompe disease manifests as a clinical spectrum that varies with respect to age at onset*, rate of disease progression, and extent of organ involvement. Patients can present with a variety of signs and symptoms, which can include cardiomegaly, cardiomyopathy, hypotonia, muscle weakness, respiratory distress (eventually requiring assisted ventilation), and skeletal muscle dysfunction.
- While there is not one generally applicable set of clinical criteria that can be used to determine appropriateness of continued therapy, clinical parameters that can indicate therapeutic response to Pombiliti + Opfolda include improved or maintained forced vital capacity, and improved or maintained 6-minute walk test (6MWT) distance.

**Although infantile-onset disease typically presents in the first year of life, age of onset alone does not necessarily distinguish between infantile- and late-onset disease since juvenile-onset disease can present prior to 12 months of age.*

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Pompe disease	<ul style="list-style-type: none"> • Members weighing ≥ 50 kg: Pombiliti 20 mg/kg IV + Opfolda 260 mg (or 4 capsules) PO every other week • Members weighing ≥ 40 kg to < 50 kg: Pombiliti 20 mg/kg IV + Opfolda 195 mg (or 3 capsules) PO every other week 	Pombiliti 20 mg/kg and Opfolda 260 mg every other week

VI. Product Availability

Drug Name	Availability
cipaglucosidase alfa-atga (Pombiliti)	Vial with lyophilized powder for reconstitution: 105 mg
miglustat (Opfolda)	Oral capsule: 65 mg

VII. References

1. Pombiliti Prescribing Information. Philadelphia, PA: Amicus Therapeutics US, LLC; September 2023. Available at: <https://amicusrx.com/pi/pombiliti.pdf>. Accessed October 3, 2023.
2. Opfolda Prescribing Information. Philadelphia, PA: Amicus Therapeutics US, LLC; September 2023. Available at: <https://amicusrx.com/pi/opfolda.pdf>. Accessed October 3, 2023.
3. Schoser B, Roberts M, Byrne BJ, et al. Safety and efficacy of cipaglucosidase alfa plus miglustat versus alglucosidase alfa plus placebo in late-onset Pompe disease (PROPEL): an international, randomised, double-blind, parallel-group, phase 3 trial. *Lancet Neurology* 2021;20:1027-37.
4. Cupler EJ, Berger KI, Leshner RT, et al. Consensus treatment recommendations for late-onset Pompe disease. *Muscle Nerve* 2012;45:319-33.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
G0138	Intravenous infusion of cipaglucosidase alfa-atga, including provider/supplier acquisition and clinical supervision of oral administration of miglustat in preparation of receipt of cipaglucosidase alfa-atga
J1202	Miglustat, oral, 65 mg
J1203	Injection, cipaglucosidase alfa-atga, 5 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy.	03.28.24	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. Louisiana Healthcare Connections makes no

CLINICAL POLICY

Cipaglucosidase Alfa-atga + Miglustat



representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Louisiana Healthcare Connections administrative policies and procedures.

This clinical policy is effective as of the date determined by Louisiana Healthcare Connections. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. Louisiana Healthcare Connections retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom Louisiana Healthcare Connections has no control or right of control. Providers are not agents or employees of Louisiana Healthcare Connections.

This clinical policy is the property of Louisiana Healthcare Connections. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members, and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2024 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.