

Clinical Policy: Paclitaxel, Protein-Bound (Abraxane)

Reference Number: LA.PHAR.176

Effective Date:

Last Review Date: 04.22

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Protein-bound paclitaxel (Abraxane®) is microtubule inhibitor.

FDA Approved Indication(s)

Abraxane is indicated for the treatment of:

- Metastatic breast cancer, after failure of combination chemotherapy for metastatic disease or relapse within 6 months of adjuvant chemotherapy. Prior therapy should have included an anthracycline unless clinically contraindicated
- Locally advanced or metastatic non-small cell lung cancer (NSCLC), as first-line treatment in combination with carboplatin, in patients who are not candidates for curative surgery or radiation therapy
- Metastatic adenocarcinoma of the pancreas as first-line treatment, in combination with gemcitabine

Policy/Criteria

Prior authorization is required. Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connection that Abraxane is medically necessary when the following criteria are met:

I. Initial Approval Criteria

A. Breast Cancer (must meet all):

1. Diagnosis of breast cancer;
2. Disease is recurrent, metastatic, or unresponsive to preoperative systemic therapy;
3. Prescribed by or in consultation with an oncologist;
4. Age ≥ 18 years;
5. For non-triple negative breast cancer: Prior therapy* included an anthracycline (e.g., doxorubicin, pegylated liposomal doxorubicin, epirubicin), unless all are contraindicated;
**Prior authorization may be required for prior therapies*
6. Request meets one of the following (a or b):*
 - a. Dose does not exceed 260 mg/m² every 3 weeks;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

CLINICAL POLICY

Paclitaxel, Protein-Bound

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

B. Non-Small Cell Lung Cancer (must meet all):

1. Diagnosis of recurrent, advanced, or metastatic NSCLC;
2. Prescribed by or in consultation with an oncologist;
3. Age \geq 18 years;
4. Member must use paclitaxel, unless contraindicated or clinically significant adverse effects are experienced;
5. Request meets one of the following (a or b):*
 - a. Dose does not exceed 100 mg/m² IV on Days 1, 8, and 15 of each 21-day cycle;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

C. Adenocarcinoma of the Pancreas (must meet all):

1. Diagnosis of adenocarcinoma of the pancreas;
2. Prescribed by or in consultation with an oncologist;
3. Age \geq 18 years;
4. Abraxane will be used in combination with gemcitabine*;
**Gemcitabine may require prior authorization*
5. Disease is metastatic, unresectable, or borderline resectable;
6. Request meets one of the following (a or b):*
 - a. Dose does not exceed 125 mg/m² on Days 1, 8 and 15 of each 28-day cycle;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

D. Additional NCCN Recommended Uses (off-label) (must meet all):

1. Prescribed for one of the following NCCN categories 1 and 2A supported indications (a - f):
 - a. AIDS-related Kaposi sarcoma;
 - b. Cutaneous or uveal melanoma, prescribed as a single agent;
 - c. Endometrial carcinoma, prescribed as a single agent;
 - d. Cholangiocarcinoma or gallbladder cancer, and member meets both of the following (i and ii):
 - i. Disease is unresectable or metastatic;
 - ii. Abraxane is prescribed in combination with gemcitabine;
 - e. Relapsed ovarian cancer;
 - f. Advanced or metastatic small bowel adenocarcinoma;
2. Prescribed by or in consultation with an oncologist;
3. Age \geq 18 years;

CLINICAL POLICY

Paclitaxel, Protein-Bound

4. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).*

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration: 6 months

E. Other diagnoses/indications

1. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connection benefit, or documentation supports that member is currently receiving Abraxane for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, meets one of the following (a or b):*
 - a. New dose does not exceed one of the following (i, ii, or iii):
 - i. For breast cancer: 260 mg/m² IV every 3 weeks;
 - ii. For NSCLC: 100 mg/m² IV on Days 1, 8, and 15 of each 21-day cycle;
 - iii. For adenocarcinoma of the pancreas: 125 mg/m² on Days 1, 8 and 15 of each 28-day cycle
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Louisiana Healthcare Connection benefit and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – LA.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

EGFR: epidermal growth factor
receptor

FDA: Food and Drug Administration

CLINICAL POLICY

Paclitaxel, Protein-Bound

HER2: human epidermal growth factor receptor 2

NSCLC: non-small cell lung cancer

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may require prior authorization.

<u>Drug Name</u>	<u>Dosing Regimen</u>	<u>Dose Limit/Maximum Dose</u>
<u>anthracyclines (e.g., doxorubicin, pegylated liposomal doxorubicin, epirubicin)</u>	<u>For breast cancer:</u> <u>Refer to prescribing information</u>	<u>Refer to prescribing information</u>
<u>paclitaxel (Taxol®)</u>	<u>For NSCLC:</u> <u>Various combinations</u>	<u>250 mg/m² every 3 weeks</u>
<u>gemcitabine (Gemzar®)</u>	<u>For adenocarcinoma of the pancreas:</u> <u>1,000 mg/m² IV over 30 to 40 minutes on days 1, 8, and 15 preceded by nab-paclitaxel (125 mg/m² IV over 30 to 40 minutes on days 1, 8, and 15) every 28 days</u>	<u>1000 mg/m² once weekly for up to 7 consecutive weeks</u>

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- **Contraindication(s): neutrophil counts of < 1,500 cells/mm³, severe hypersensitivity**
- **Boxed warning(s): neutropenia**

V. Dosage and Administration

<u>Indication</u>	<u>Dosing Regimen</u>	<u>Maximum Dose</u>
<u>Metastatic breast cancer</u>	<u>260 mg/m² IV every 3 weeks</u>	<u>260 mg/m²</u>
<u>Non-small cell lung cancer</u>	<u>100 mg/m² IV on days 1, 8, and 15 of each 21-day cycle</u>	<u>260 mg/m²</u>
<u>Metastatic adenocarcinoma of the pancreas</u>	<u>125 mg/m² IV on days 1, 8 and 15 of each 28-day cycle</u>	<u>260 mg/m²</u>

VI. Product Availability

Injectable suspension: lyophilized powder containing 100 mg of paclitaxel formulated as albumin-bound particles in single-use vial for reconstitution

CLINICAL POLICY

Paclitaxel, Protein-Bound

VII. References

1. Abraxane Prescribing Information. Summit, NJ: Celgene Corporation; August 2020. Available at <http://www.abraxane.com/>. Accessed February 21, 2022.
2. Paclitaxel, albumin bound. In: National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed February 21, 2022.
3. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc. Updated periodically. Accessed February 20, 2022.

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2021, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

The following is a list of procedures codes for which coverage may be provided when billed with a diagnosis code(s) that supports coverage criteria (see list of ICD codes supporting coverage criteria further below).

<u>CPT® /HCPCS Codes</u>	<u>Description</u>
<u>96413</u>	<u>Chemotherapy administration, intravenous infusion technique, up to 1 hour, single or initial substance/drug</u>
<u>96415</u>	<u>Chemotherapy administration; each additional hour (List separately in addition to code for primary procedure)</u>
<u>J9264</u>	<u>Injection, paclitaxel protein-bound particles, 1 mg</u>

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

The following is a list of diagnosis codes that support coverage for the applicable covered procedure code(s).

<u>ICD-10-CM Code</u>	<u>Description</u>
<u>C17.0</u>	<u>Malignant neoplasm of duodenum</u>
<u>C17.1</u>	<u>Malignant neoplasm of jejunum</u>
<u>C17.2</u>	<u>Malignant neoplasm of ileum</u>
<u>C17.3</u>	<u>Meckel's diverticulum, malignant</u>
<u>C17.8</u>	<u>Malignant neoplasm of overlapping sites of small intestine</u>
<u>C17.9</u>	<u>Malignant neoplasm of small intestine, unspecified</u>
<u>C22.1</u>	<u>Intrahepatic bile duct carcinoma</u>
<u>C25.0-C25.9</u>	<u>Malignant neoplasm of pancreas</u>
<u>C34.00-C34.02</u>	<u>Malignant neoplasm of main bronchus</u>
<u>C34.10-C34.12</u>	<u>Malignant neoplasm of upper lobe, bronchus or lung</u>

CLINICAL POLICY

Paclitaxel, Protein-Bound

<u>ICD-10-CM Code</u>	<u>Description</u>
<u>C34.2</u>	<u>Malignant neoplasm of middle lobe, bronchus or lung</u>
<u>C34.30-C34.32</u>	<u>Malignant neoplasm of lower lobe, bronchus or lung</u>
<u>C34.80-C34.82</u>	<u>Malignant neoplasm of overlapping sites of bronchus or lung</u>
<u>C34.90-C34.92</u>	<u>Malignant neoplasm of unspecified part of unspecified bronchus or lung</u>
<u>C43.0-C43.8</u>	<u>Melanoma and other malignant neoplasms of skin</u>
<u>C46.0-C46.9</u>	<u>Kaposi's sarcoma</u>
<u>C48.1</u>	<u>Malignant neoplasm of specified parts of peritoneum</u>
<u>C48.2</u>	<u>Malignant neoplasm of peritoneum, unspecified</u>
<u>C48.8</u>	<u>Malignant neoplasm of overlapping sites of retroperitoneum and peritoneum</u>
<u>C50.011-C50.012</u>	<u>Malignant neoplasm of nipple and areola, female</u>
<u>C50.021-C50.022</u>	<u>Malignant neoplasm of nipple and areola, male</u>
<u>C50.111-C50.112</u>	<u>Malignant neoplasm of central portion of breast, female</u>
<u>C50.121-C50.122</u>	<u>Malignant neoplasm of central portion of breast, male</u>
<u>C50.211-C50.212</u>	<u>Malignant neoplasm of upper-inner quadrant of breast, female</u>
<u>C50.221-C50.222</u>	<u>Malignant neoplasm of upper-inner quadrant of breast, male</u>
<u>C50.311-C50.312</u>	<u>Malignant neoplasm of lower-inner quadrant of breast, female</u>
<u>C50.321-C50.322</u>	<u>Malignant neoplasm of lower-inner quadrant of breast, male</u>
<u>C50.411-C50.412</u>	<u>Malignant neoplasm of upper-outer quadrant of breast, female</u>
<u>C50.421-C50.422</u>	<u>Malignant neoplasm of upper-outer quadrant of breast, male</u>
<u>C50.511-C50.512</u>	<u>Malignant neoplasm of lower-outer quadrant of breast, female</u>
<u>C50.521-C50.522</u>	<u>Malignant neoplasm of lower-outer quadrant of breast, male</u>
<u>C50.611-C50.612</u>	<u>Malignant neoplasm of axillary tail of breast, female</u>
<u>C50.621-C50.622</u>	<u>Malignant neoplasm of axillary tail of breast, male</u>
<u>C50.811-C50.812</u>	<u>Malignant neoplasm of overlapping sites of breast, female</u>
<u>C50.821-C50.822</u>	<u>Malignant neoplasm of overlapping sites of breast, male</u>
<u>C50.911-C50.912</u>	<u>Malignant neoplasm of breast of unspecified site, female</u>
<u>C50.921-C50.922</u>	<u>Malignant neoplasm of breast of unspecified site, male</u>
<u>C54.1</u>	<u>Malignant neoplasm of endometrium</u>
<u>C56.1-C56.2</u>	<u>Malignant neoplasm of ovary</u>
<u>C57.01-C57.02</u>	<u>Malignant neoplasm of fallopian tube</u>
<u>C57.11-C57.12</u>	<u>Malignant neoplasm of broad ligament</u>
<u>C57.21-C57.22</u>	<u>Malignant neoplasm of round ligament</u>
<u>C57.3</u>	<u>Malignant neoplasm of parametrium</u>
<u>C57.4</u>	<u>Malignant neoplasm of uterine adnexa, unspecified</u>
<u>C57.7</u>	<u>Malignant neoplasm of other specified female genital organs</u>
<u>C57.8</u>	<u>Malignant neoplasm of overlapping sites of female genital organs</u>
<u>C65.1 – C65.2</u>	<u>Malignant neoplasm of renal pelvis</u>
<u>C67.0 – C67.9</u>	<u>Malignant neoplasm of bladder</u>
<u>C68.0</u>	<u>Malignant neoplasm of urethra</u>
<u>C69.31-C69.32</u>	<u>Malignant neoplasm of choroid</u>

CLINICAL POLICY

Paclitaxel, Protein-Bound

<u>ICD-10-CM Code</u>	<u>Description</u>
<u>C69.41-C69.42</u>	<u>Malignant neoplasm of ciliary body</u>
<u>Z85.05</u>	<u>Personal history of malignant neoplasm of liver</u>
<u>Z85.068</u>	<u>Personal history of other malignant neoplasm of small intestine</u>
<u>Z85.07</u>	<u>Personal history of malignant neoplasm of pancreas</u>
<u>Z85.118</u>	<u>Personal history of other malignant neoplasm of bronchus and lung</u>
<u>Z85.3</u>	<u>Personal history of malignant neoplasm of breast</u>
<u>Z85.42</u>	<u>Personal history of malignant neoplasm of other parts of uterus</u>
<u>Z85.43</u>	<u>Personal history of malignant neoplasm of ovary</u>
<u>Z85.44</u>	<u>Personal history of malignant neoplasm of other female genital organs</u>
<u>Z85.51</u>	<u>Personal history of malignant neoplasm of bladder</u>
<u>Z85.53</u>	<u>Personal history of renal pelvis</u>
<u>Z85.820</u>	<u>Personal history of malignant melanoma of skin</u>
<u>Z85.840</u>	<u>Personal history of malignant neoplasm of eye</u>

<u>Reviews, Revisions, and Approvals</u>	<u>Date</u>	<u>LDH Approval Date</u>
<u>Converted corporate to local policy</u>	<u>04.22</u>	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to

CLINICAL POLICY

Paclitaxel, Protein-Bound

applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2020 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.