

## Clinical Policy: Teclistamab-cqyv (Tecvayli)

Reference Number: LA.PHAR.611

Effective Date:

Last Review Date: 05.01.23

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

**\*\*Please note: This policy is for medical benefit\*\***

### Description

Teclistamab-cqyv (Tecvayli™) is a humanized recombinant immunoglobulin G4-proline, alanine, alanine (IgG4-PAA) antibody, and a bispecific B-cell maturation antigen (BCMA)-directed CD3 T-cell engager.

### FDA Approved Indication(s)

Tecvayli is indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior lines of therapy, including a proteasome inhibitor (PI), an immunomodulatory agent (IMiD), and an anti-CD38 monoclonal antibody.

This indication is approved under accelerated approval based on response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of Louisiana Healthcare Connections® that Tecvayli is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Multiple Myeloma (must meet all):

1. Diagnosis of relapsed or refractory multiple myeloma (RRMM);
2. Prescribed by or in consultation with a hematologist or an oncologist;
3. Age  $\geq$  18 years;
4. Tecvayli is prescribed as monotherapy;
5. Member has measurable disease as evidenced by one of the following assessed within the last 28 days (a, b, or c):
  - a. Serum M-protein  $\geq$  1 g/dL;
  - b. Urine M-protein  $\geq$  200 mg/24 h;
  - c. Serum free light chain (FLC) assay: involved FLC level  $\geq$  10 mg/dL (100 mg/L) provided serum kappa lambda FLC ratio is abnormal;
6. Member has received or has documented intolerance to  $\geq$  4 prior lines of therapy (*see Appendix B for examples*) that include all of the following (a, b, and c):

## CLINICAL POLICY

### Generic Name

- a. One proteasome inhibitor (e.g., bortezomib, Kyprolis®, Ninlaro®);
  - b. One immunomodulatory agent (e.g., Revlimid®, pomalidomide, Thalomid®);
  - c. One anti-CD38 antibody (e.g., Darzalex®/Darzalex Faspro™, Sarclisa®);
- \*Prior authorization may be required*
7. Member does not have a known active central nervous system (CNS) involvement (e.g., seizure, cerebrovascular ischemia) or exhibits clinical signs of meningeal involvement of multiple myeloma;
  8. Member has not previously received treatment with anti-BCMA targeted therapy (e.g., Blenrep™, Abecma®, or Carvykti™);
  9. Dose does not exceed all of the following (a, b, c, d, and e):\*
    - a. Day 1: 0.06 mg/kg;
    - b. Day 4: 0.3 mg/kg;
    - c. Day 7: 1.5 mg/kg;
    - d. Day 8 and thereafter: 1.5 mg/kg per week;
    - e. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label dose use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration:** 6 months

#### **B. Other diagnoses/indications** (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

## II. Continued Therapy

#### **A. Multiple Myeloma** (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving Tecvayli for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):\*
  - a. New dose does not exceed 1.5 mg/kg per week;
  - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label dose use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration:** 12 months

#### **B. Other diagnoses/indications** (must meet 1 or 2):

## CLINICAL POLICY

### Generic Name

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 2 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

### III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off-label use policies – LA.PMN.53 for Medicaid or evidence of coverage documents.
- B. Active or prior history of CNS involvement with myeloma (e.g., seizures, cerebrovascular ischemia) or exhibit clinical signs of meningeal involvement of multiple myeloma.

### IV. Appendices/General Information

#### Appendix A: Abbreviation/Acronym Key

BCMA: B-cell maturation antigen

CNS: central nervous system

CRS: cytokine release syndrome

FDA: Food and Drug Administration

ICANS: immune effector cell-associated neurotoxicity syndrome

IMiD: immunomodulatory drug

MM: multiple myeloma

PI: proteasome inhibitor

RRMM: relapsed or refractory multiple myeloma

#### Appendix B: Therapeutic Alternatives

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
bortezomib/Revlimid® (lenalidomide)/dexamethasone	Varies	Varies
bortezomib/cyclophosphamide/dexamethasone	Varies	Varies
bortezomib/doxorubicin (or liposomal doxorubicin)/dexamethasone	Varies	Varies
Kyprolis® (carfilzomib) Revlimid® (lenalidomide)/dexamethasone	Varies	Varies
Kyprolis® (carfilzomib)/cyclophosphamide/dexamethasone	Varies	Varies
Kyprolis® (carfilzomib – weekly or twice weekly)/dexamethasone	Varies	Varies
Ninlaro® (ixazomib)/Revlimid® (lenalidomide)/dexamethasone	Varies	Varies
Ninlaro® (ixazomib)/dexamethasone	Varies	Varies

## CLINICAL POLICY

### Generic Name

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Ninlaro® (ixazomib)/pomalidomide/dexamethasone	Varies	Varies
bortezomib/dexamethasone	Varies	Varies
bortezomib/Thalomid® (thalidomide)/dexamethasone	Varies	Varies
cyclophosphamide/Revlimid® (lenalidomide)/dexamethasone	Varies	Varies
Revlimid® (lenalidomide)/dexamethasone	Varies	Varies
VTD-PACE (dexamethasone/Thalomid® (thalidomide)/cisplatin/doxorubicin/cyclophosphamide/etoposide/bortezomib)	Varies	Varies
Revlimid® (lenalidomide)/low-dose dexamethasone	Varies	Varies
Darzalex® (daratumumab) or Darzalex Faspro™ (daratumumab/hyaluronidase-fihj)/bortezomib/melphan/prednisone	Varies	Varies
Darzalex® (daratumumab) or Darzalex Faspro™ (daratumumab/hyaluronidase-fihj)/bortezomib/dexamethasone	Varies	Varies
Darzalex® (daratumumab) or Darzalex Faspro™ (daratumumab/hyaluronidase-fihj)/Revlimid® (lenalidomide)/dexamethasone	Varies	Varies
Darzalex® (daratumumab) or Darzalex Faspro™ (daratumumab/hyaluronidase-fihj)	Varies	Varies
Darzalex® (daratumumab) or Darzalex Faspro™ (daratumumab/hyaluronidase-fihj)/pomalidomide/dexamethasone	Varies	Varies
Empliciti® (elotuzumab)/Revlimid® (lenalidomide)/dexamethasone	Varies	Varies
Empliciti® (elotuzumab)/bortezomib/dexamethasone	Varies	Varies
Empliciti® (elotuzumab)/pomalidomide/dexamethasone	Varies	Varies
bendamustine/bortezomib/dexamethasone	Varies	Varies
bendamustine/Revlimid® (lenalidomide)/dexamethasone	Varies	Varies
panobinostat/bortezomib/dexamethasone	Varies	Varies
panobinostat/Kyprolis® (carfilzomib)	Varies	Varies
panobinostat/Revlimid® (lenalidomide)/dexamethasone	Varies	Varies
pomalidomide/cyclophosphamide/dexamethasone	Varies	Varies
pomalidomide/dexamethasone	Varies	Varies
pomalidomide/bortezomib/dexamethasone	Varies	Varies
pomalidomide/Kyprolis® (carfilzomib)/dexamethasone	Varies	Varies
Sarclisa® (isatuximab-irfc)/pomalidomide/dexamethasone	Varies	Varies

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

## CLINICAL POLICY

### Generic Name

#### *Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): None
- Boxed warning(s): Cytokine release syndrome (CRS) with life-threatening and/or fatal reactions, and neurologic toxicity, including immune effector cell-associated neurotoxicity syndrome (ICANS)

#### *Appendix D: General Information*

- Due to the risks of cytokine release syndrome, patients should be hospitalized for 48 hours after administration of all doses within the step-up dosing schedule including the first maintenance dose. Subsequent weekly maintenance doses are managed on outpatient basis according to the Tecvayli REMS program (*see appendix E for more details on REMS Program*).
- In the MajesTEC-1 trial, 100% of enrolled patients reported having an adverse event, of which 94.5% were grade 3 or 4. The most common hematologic adverse events were neutropenia (70.9%), anemia (52.1%), and thrombocytopenia (44.0%). The most common non-hematologic adverse events were diarrhea (28.5%), fatigue (27.9%), and nausea (27.3%).
- In the MajesTEC-1 trial, 72.1% of participants experienced any grade cytokine release syndrome (CRS), and 14.5% of participants experienced any grade immune effector cell-associated neurotoxicity syndrome (ICANS). Both toxicities were managed with supportive measures that included administration of tocilizumab (in 60/119 patients with CRS, and 3/24 patients with ICANS), low-flow oxygen by nasal cannula, glucocorticoids, levetiracetam, and gabapentin.
- In the MajesTEC-1 trial, five deaths were considered to have been related to Tecvayli treatment including one death resulting from progressive multifocal leukoencephalopathy, two deaths related to Covid-19, one death related to hepatic failure, and one death related to streptococcal pneumonia. Subjects positive for hepatitis B, hepatitis C, and/or HIV were excluded from the trial. Prior to treatment with Tecvayli initiation of antiviral prophylaxis to prevent herpes zoster reactivation is recommended.

#### *Appendix E: Tecvayli REMS Program Information*

- Tecvayli is available only through a restricted REMS program due to the risk of cytokine release syndrome and neurologic toxicity, including ICANS.
- Prescribers are required to:
  - 1) obtain certification with the program by enrolling and completing training,
  - 2) counsel patients about the risks associated with Tecvayli therapy,
  - 3) provide patients with patient wallet card.
- Dispensers are required to:
  - 1) obtain certification with the program,
  - 2) verify prescriber certification status with the program prior to dispensing the product.

## V. Dosage and Administration

## CLINICAL POLICY

### Generic Name

Indication	Dosing Regimen	Maximum Dose
Multiple Myeloma	<p>Step-up Dosing Schedule<sup>a</sup>:</p> <ul style="list-style-type: none"> <li>Day 1: 0.06 mg/kg subcutaneously (step-up dose 1)</li> <li>Day 4<sup>b</sup>: 0.3 mg/kg subcutaneously (step-up dose 2)</li> <li>Day 7<sup>c</sup>: 1.5 mg/kg subcutaneously (first treatment dose)</li> </ul> <p>Weekly Dosing Schedule<sup>a</sup>:</p> <ul style="list-style-type: none"> <li>1.5 mg/kg subcutaneously once weekly (one week after first treatment dose and weekly thereafter)</li> </ul>	1.5 mg/kg per week subcutaneously

<sup>a</sup> Refer to prescribing information Table 2 for recommendations on restarting therapy due to dose delays.

<sup>b</sup> Step-up dose 2 may be given between 2 to 4 days after step-up dose 1 and may be given up to 7 days after step-up dose 1 to allow for resolution of adverse reactions.

<sup>c</sup> First treatment dose may be given between 2 to 4 days after step-up dose 2 and may be given up to 7 days after step-up dose 2 to allow for resolution of adverse reactions.

### VI. Product Availability

Solution for subcutaneous injection in a single-dose vial:

- 30 mg/3 mL (10 mg/mL) used for step-up doses 1 and 2
- 153 mg/1.7 mL (90 mg/mL) used for treatment doses

### VII. References

1. Tecvayli Prescribing Information. Horsham, PA: Janssen Biotech, Inc.; October 2022. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/761291s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/761291s000lbl.pdf). Accessed November 10, 2022.
2. ClinicalTrials.gov. A phase 1, first-in-human, open-label, dose escalation study of teclistamab, a humanized BCMA x CD3 bispecific antibody in subjects with relapsed or refractory multiple myeloma. Available at: <https://www.clinicaltrials.gov/ct2/show/NCT03145181>. Accessed November 10, 2022.
3. ClinicalTrials.gov. A phase 1/2, first-in-human, open-label, dose escalation study of teclistamab, a humanized BCMA x CD3 bispecific antibody, in subjects with relapsed or refractory multiple myeloma. Available at: <https://clinicaltrials.gov/ct2/show/NCT04557098>. Accessed November 10, 2022.
4. Girgis S, Lin SXW, Pillarisetti K, et al. Translational modeling predicts efficacious therapeutic dosing range of teclistamab for multiple myeloma. *Target Oncol*. 2022;17(4):433-439.
5. Moreau P, Garfall AL, van de Donk NWCJ, et al. Teclistamab in relapsed or refractory multiple myeloma. *N Engl J Med*. 2022;387(6):495-505.
6. National Comprehensive Cancer Network. Multiple Myeloma Version 2.2023. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/myeloma.pdf](https://www.nccn.org/professionals/physician_gls/pdf/myeloma.pdf). Accessed November 10, 2022.
7. Pillarisetti K, Powers G, Luistro L, et al. Teclistamab is an active T cell-redirecting bispecific antibody against B-cell maturation antigen for multiple myeloma. *Blood Adv*. 2020;4(18):4538-4549.



## CLINICAL POLICY

### Generic Name

8. Usmani SZ, Garfall AL, van de Donk NWCJ, et al. Teclistamab, a B-cell maturation antigen × CD3 bispecific antibody, in patients with relapsed or refractory multiple myeloma (MajesTEC-1): a multicentre, open-label, single-arm, phase 1 study. *Lancet*. 2021;398(10301):665-674.

### Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPSC Codes	Description
J3590	Unclassified biologics
J9999	Not otherwise classified antineoplastic drugs
C9399	Unclassified drugs or biologicals

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Policy created	05.01.23	

### Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between

## CLINICAL POLICY

### Generic Name

the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2023 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.

.