

Clinical Policy: Bortezomib (Velcade)

Reference Number: LA.PHAR.410

Effective Date: 03.16.23

Last Review Date: 06.25.23~~02.23~~

Line of Business: Medicaid

Coding Implications
Revision Log

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

****Please note: This policy is for medical benefit****

Description

Bortezomib (Velcade®) is a proteasome inhibitor.

FDA Approved Indication(s)

Velcade is indicated for treatment of adult patients with:

- Multiple myeloma (MM)
- Mantle cell lymphoma (MCL)*

**The 1 mg and 2.5 mg strengths are indicated specifically for patients who have received at least 1 prior therapy.*

Policy/Criteria

~~Prior authorization is required.~~ Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that Velcade is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Multiple Myeloma and Mantle Cell Lymphoma (must meet all):

1. Diagnosis of one of the following (a or b):
 - a. MM;
 - b. MCL (B-cell lymphoma subtype);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age \geq 18 years;
4. For Velcade requests, member must use bortezomib, if available, unless contraindicated or clinically significant adverse effects are experienced;
5. Request meets one of the following (a or b):*
 - a. Dose does not exceed 1.3 mg/m²;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN.*

Approval duration: 6 months

B. NCCN Recommended Uses (off-label) (must meet all):

1. Diagnosis of one of the following (a-h):
 - a. AIDS-related Kaposi sarcoma (advanced cutaneous, oral, visceral, or nodal disease) - after \geq 2 prior lines of systemic therapy;

CLINICAL POLICY

Bortezomib

- b. Multicentric Castleman's disease (B-cell lymphoma subtype) - as subsequent therapy;
 - c. Systemic light chain amyloidosis;
 - d. Adult T-cell leukemia/lymphoma - as subsequent therapy;
 - e. Waldenström macroglobulinemia/lymphoplasmacytic lymphoma;
 - f. T-cell acute lymphoblastic leukemia (T-ALL) – for relapsed or refractory disease;
 - g. Pediatric acute lymphoblastic leukemia (ALL) - as subsequent therapy;
 - h. Pediatric Hodgkin lymphoma (HL) - as subsequent therapy in combination with ifosafamide and vinorelbine;
2. Prescribed by or in consultation with an oncologist or hematologist;
 3. Age \geq 18 years (all indications except pediatric ALL and HL);
 4. For Velcade requests, member must use bortezomib, if available, unless contraindicated or clinically significant adverse effects are experienced;
 5. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).*

*Prescribed regimen must be FDA-approved or recommended by NCCN.

Approval duration: 6 months

C. Other diagnoses/indications

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53
- ~~1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):~~
 - ~~a. For drugs on the PDL (Medicaid), the no-coverage criteria policy LA.PMN.255 for Medicaid; or~~
 - ~~b. For drugs NOT on the PDL (Medicaid), the non-formulary policy LA.PMN.16 for Medicaid; or 2.~~
- ~~2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53 for Medicaid.~~

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving the requested agent for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;

CLINICAL POLICY

Bortezomib

3. For Velcade requests, member must use bortezomib, if available, unless contraindicated or clinically significant adverse effects are experienced;
4. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 1.3 mg/m²;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN.

Approval duration: 12 months

A. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53
- ~~1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):~~
 - ~~a. For drugs on the PDL (Medicaid), the no coverage criteria policy LA.PMN.255 for Medicaid; or~~
 - ~~b. For drugs NOT on the PDL (Medicaid), the non-formulary policy LA.PMN.16 for Medicaid; or~~
- ~~2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53 for Medicaid.~~

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – LA.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ALL: acute lymphoblastic leukemia
 FDA: Food and Drug Administration
 HL: Hodgkin lymphoma
 MCL: mantle cell lymphoma

MM: multiple myeloma
 NCCN: National Comprehensive Cancer Network
 T-ALL: T-cell acute lymphoblastic leukemia

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):

CLINICAL POLICY

Bortezomib

- Patients with hypersensitivity (not including local reactions) to bortezomib, boron, or mannitol, including anaphylactic reactions
- Contraindicated for intrathecal administration
- Boxed warning(s): none reported

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
MM	<ul style="list-style-type: none"> ● <u>First-line therapy</u>: 1.3 mg/m² as a 3 to 5 second bolus IV injection or SC injection in combination with PO melphalan and PO prednisone for nine 6-week treatment cycles. ● <u>Relapse*</u>: 1.3 mg/m² as a 3 to 5 second bolus IV injection or SC injection as a single agent or in combination with dexamethasone for up to eight 3-week cycles. For therapy beyond eight cycles, see PI for additional dosing options. <i>*If relapse occurs ≥ 6 months after a previous response to Velcade, treatment may be restarted at the last tolerated dose.</i> 	1.3 mg/m ²
MCL	<ul style="list-style-type: none"> ● <u>First-line therapy</u>: 1.3 mg/m² as a 3 to 5 second bolus IV injection or SC injection in combination with IV rituximab, cyclophosphamide, doxorubicin and PO prednisone (VcR-CAP) for up to six 3-week treatment cycles, plus two additional cycles if a positive response. ● <u>Relapse</u>: 1.3 mg/m² as a 3 to 5 second bolus IV injection or SC injection for up to eight 3-week treatment cycles. Therapy may extend beyond eight cycles. 	1.3 mg/m ²

VI. Product Availability*

Single-dose vials for injection:

- Sterile lyophilized powder for reconstitution: 1 mg, 2.5 mg, 3.5 mg
- Solution: 2.5 mg/mL, 3.5 mg/1.4 mL

**The branded product, Velcade, is only available as 3.5 mg sterile lyophilized powder*

VII. References

1. Velcade Prescribing Information. Cambridge, MA: Millennium Pharmaceuticals, Inc.; August 2022. Available at: <https://www.velcade.com> Accessed November 11, 2022.
2. Bortezomib Prescribing Information. Lake Forest, IL: Hospira, Inc.; May 2022. Available at https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/209191s000lbl.pdf. Accessed November 11, 2022.
3. Bortezomib Prescribing Information. Durham, NC: Accord Healthcare; July 2022. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/215441s000lbl.pdf. Accessed November 11, 2022.
4. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed November 11, 2022.

CLINICAL POLICY

Bortezomib

5. National Comprehensive Cancer Network. T-Cell Lymphomas Version 2.2022. Available at: https://www.nccn.org/professionals/physician_gls/pdf/t-cell.pdf. Accessed November 11, 2022.
6. National Comprehensive Cancer Network. Pediatric Acute Lymphoblastic Leukemia Version 1.2023. Available at: https://www.nccn.org/professionals/physician_gls/pdf/ped_all.pdf. Accessed November 11, 2022.
7. National Comprehensive Cancer Network. Acute Lymphoblastic Leukemia Version 1.2023. Available at: https://www.nccn.org/professionals/physician_gls/pdf/all.pdf. Accessed November 11, 2022.
8. National Comprehensive Cancer Network. B-Cell Lymphomas Version 5.2022. Available at: https://www.nccn.org/professionals/physician_gls/pdf/b-cell.pdf. Accessed November 14, 2022.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPSC Codes	Description
J9041	Injection, bortezomib (Velcade), 0.1 mg
J9044	Injection, bortezomib (not otherwise specified), 0.1 mg
<u>J9046</u>	<u>Injection, bortezomib, (dr.reddy's), not therapeutically equivalent to</u>
<u>J9048</u>	<u>Injection, bortezomib (Fresenius kabi), not therapeutically equivalent to J9041, 0.1mg</u>
<u>J9049</u>	<u>Injection, bortezomib (Hospira), not therapeutically equivalent to J9041, 0.1mg</u>

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy.	02.23	03.16.23
<u>Updated criteria for other diagnoses/indications</u> <u>Added HCPSC Codes: J9046, J9048, J9049</u>	<u>06.25.23</u>	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2020 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.