

**Clinical Policy: Ibandronate Injection (Boniva)** 

Reference Number: LA.PHAR.189

Effective Date: <u>09.21</u>

Last Review Date: 073.224 Coding Implications
Line of Business: Medicaid Revision Log

See Important Reminder at the end of this policy for important regulatory and legal information.

## **Description**

Ibandronate injection (Boniva®) is a bisphosphonate.

#### **FDA Approved Indication(s)**

Boniva is indicated for:

• <u>Postmenopausal osteoporosis (PMO):</u> Treatment of osteoporosis in postmenopausal women. In postmenopausal women with osteoporosis, Boniva increases bone mineral density (BMD) and reduces the incidence of vertebral fractures.

**Limitation(s) of use:** Optimal duration of use has not been determined. For patients at low-risk for fracture, consider drug discontinuation after 3 to 5 years of use.

## Policy/Criteria

**Prior authorization is required.** Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that Boniva injection is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

- **A.** Osteoporosis (must meet all):
  - 1. Diagnosis of PMO;
  - 2. Age  $\geq$  18 years or documentation of closed epiphyses on x-ray;
  - 3. Failure of a 12-month oral bisphosphonate\* trial (*Appendix B*) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
    - \*Prior authorization may be required.
  - 4. Dose does not exceed 3 mg (1 syringe) every 3 months.

#### **Approval duration:**

Medicaid – 6 months

## **B.** Other diagnoses/indications

 Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.

#### **II.** Continued Therapy

## louisiana healthcare connections

# CLINICAL POLICY Ibandronate Injection

## **A.** Osteoporosis (must meet all):

- 1. Currently receiving medication via Louisiana Healthcare Connections benefit or member has previously met initial approval criteria;
- 2. Member is responding positively to therapy;
- 3. If request is for a dose increase, new dose does not exceed 3 mg (1 syringe) every 3 months.

# **Approval duration:**

Medicaid – 12 months

## **B. Other diagnoses/indications** (must meet 1 or 2):

- 1. Currently receiving medication via Louisiana Healthcare Connections benefit and documentation supports positive response to therapy.
  - Approval duration: Duration of request or 6 months (whichever is less); or
- 2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.

## III. Diagnoses/Indications for which coverage is NOT authorized:

**A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – LA.PMN.53 for Medicaid or evidence of coverage documents.

#### IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

BMD: bone mineral density

FDA: Food and Drug Administration PMO: postmenopausal osteoporosis

#### Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Oral bisphosphonates		
alendronate (Fosamax <sup>®</sup> )	Treatment/prevention: PMO Treatment: GIO, male osteoporosis Treatment: Paget disease See prescribing information for dose.	Varies
Fosamax <sup>®</sup> Plus D (alendronate / cholecalciferol)	Treatment: PMO, male osteoporosis See prescribing information for dose.	
risedronate (Actonel <sup>®</sup> , Atelvia <sup>®</sup> )	Actonel: Treatment/prevention: PMO, GIO Treatment: male osteoporosis	



# **CLINICAL POLICY**Ibandronate Injection

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	Treatment: Paget disease	
	Atelvia:	
	Treatment: PMO	
	See prescribing information for dose.	
ibandronate (Boniva®)	Treatment/prevention: PMO	
	See prescribing information for dose.	

Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.

## Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): hypocalcemia, hypersensitivity
- Boxed warning(s): none reported

#### V. Dosage and Administration

Indication	<b>Dosing Regimen</b>	Maximum Dose
PMO	3 mg IV every 3 months	3 mg/3 months

## VI. Product Availability

Single-use prefilled syringe: 3 mg/3 mL

#### VII. References

- 1. Boniva Injection Prescribing Information. South San Francisco, CA: Genentech USA, Inc.; April 2019. Available at https://www.gene.com. Accessed SeptemberOctober 1426, 20210.
- 2. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2020. URL: http://www.clinicalpharmacology.com.

## Osteoporosis Diagnosis, Fracture Risk, and Treatment

- 3. Shoback D, Rosen CJ, Black DM, et al. Pharmacological management of osteoporosis in postmenopausal women: an endocrine society guideline update. J Clin Endocrinol Metab; March 2020, 105(3): 587-594.
- 4. Eastell R, Rosen CJ, Black DM, et al. Pharmacological management of osteoporosis in postmenopausal women: An Endocrine Society Clinical Practice Guideline. J Clin Endocrinol Metab; 2019, 104: 1595–1622.
- Camacho PM, Petak SM, Brinkley N et al. AACE/ACE Guidelines- American Association of Clinical Endocrinologists and American College of Endocrinology Clinical Practice Guidelines for Diagnosis and Treatment of Postmenopausal Osteoporosis-2020 update. Endocrine Practice 2020;26(1):1-46Vol 22 (suppl 4) September 2016.
- National Osteoporosis Foundation Clinician's Guide to Prevention and Treatment of Osteoporosis. Osteoporosis International 2014. Available at: <a href="https://cdn.nof.org/wp-content/uploads/2016/01/995.pdf">http://nof.org/wp-content/uploads/2016/01/995.pdf</a>. <a href="https://nof.org/files/nof/public/content/file/2791/upload/919.pdf">http://nof.org/files/nof/public/content/file/2791/upload/919.pdf</a>.
   Accessed September October 1431, 202118.
- 7. Siris ES, Adler R, Bilezikian J, et al. The clinical diagnosis of osteoporosis: a position statement from the National Bone Health Alliance Working Group. Osteoporos Int (2014) 25:1439–1443. DOI 10.1007/s00198-014-2655-z.

## louisiana healthcare connections

# **CLINICAL POLICY**Ibandronate Injection

8. Hodsman AB, Bauder DC, Dempster DW, et al. Parathyroid hormone and teriparatide for the treatment of osteoporosis: a review of the evidence and suggested guidelines for its use. Endocr Rev. 2005 Aug;26(5):688-703. Epub 2005 Mar 15.

## **Coding Implications**

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS	Description
Codes	
J1740	Injection, ibandronate sodium, 1 mg

Reviews, Revisions, and Approvals	Date
Converted corporate to local policy	03.2021
References reviewed and updated.	<u>07.22</u>

## **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or



# CLINICAL POLICY Ibandronate Injection

withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2020 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.