



# Reconciling Monkeypox Vaccines in LINKS

## STEP ONE:

To correct your inventory in LINKS, follow these steps to make the appropriate adjustments. First, login to LINKS. Select your organization and facility. View the left column of the LINKS Home page and click the "Lot Numbers" tab and select "Reconciliation".

Lot Numbers
Reconciliation
Search/Add
Search Results
Detail
Correct Decrementing

## STEP TWO:

The Reconcile Inventory page will open and display all vaccines in your inventory. Scroll down the list of vaccines until you find "Vaccinia, smallpox monkeypox vaccine live, PF ((monkeypox))". You will see the number of doses available in the column "Quantity on Hand" for each LOT Number.

Facility Display Name:

Vaccine	Lot Number	Exp Date	Quantity on Hand	Physical Inventory	Adjustment (+/-)	Category	Reason	Funding Source	Inactive	Add Row
Vaccinia, smallpox monkeypox vaccine live, PF ((monkeypox))	FDP00004	08/31/2023	0		0.0	--No Category Required--	--No Reason Required--	PUB	<input type="checkbox"/>	+
Vaccinia, smallpox monkeypox vaccine live, PF ((monkeypox))	FDP00013	09/30/2024	83		0.0	--No Category Required--	--No Reason Required--	PUB	<input type="checkbox"/>	+

Here you will record the number of physical doses remaining in the "Physical Inventory" blank for the LOT Number you are reconciling.

### Example:

Three doses of LOT# FDP00013 were administered Intradermal injection. You now have two doses wasted with 78 physical doses remaining. 78 is to be recorded as Physical Inventory. The Adjustment (+/-) column will automatically calculate # of doses to be reconciled. (83 Doses – 5 Doses = 78 Doses)

## STEP THREE:

After recording your new inventory count, you must select the "Category" that describes the doses and "Reason" for reconciliation. Click "select" to choose from the options. Remember to "SAVE".

Category	Reason
--select--	--select--
Administered	Broken/Dropped/Spilled
Expired	Dose Count Variance Multi-dose Vial
Recall	Drawn up, not used
Spoiled	Lost and Unaccounted
Transfer	Vaccine Damaged in Transit
Wasted	

### IMPORTANT NOTES:

- ❖ For patients who receive the vaccine by Subcutaneous Injection, 1 dose from the vial must be administered to the patient and 4 doses must be removed from your inventory as "Wasted" due to "dose count variance multi-dose vial".
- ❖ Unused/Wasted doses must be reconciled immediately to keep the MPX inventory correct.
- ❖ Do not change dose volume.

Table 2. Vaccination Schedule and Dosing Regimens for JYNNEOS Vaccine

JYNNEOS vaccine regimen	Route of administration	Injection Volume	Recommended number of doses	Recommended interval between 1st and 2nd dose
<b>Alternative regimen</b>				
People age ≥18 years	ID	0.1 mL	2	28 days
<b>Standard regimen</b>				
People age <18 years	Subcut	0.5 mL	2	28 days
People of any age who have a history of developing keloid scars	Subcut	0.5 mL	2	28 days